

Two-Player Games



Who is this man?

Two-Player Games



Does Garry Kasparov have a winning strategy?

Two-Player Games

Given a propositional formula G with variables $p_1, q_1, \dots, p_n, q_n$.

Two-Player Games

Given a propositional formula G with variables $p_1, q_1, \dots, p_n, q_n$.

There are two players: P and Q .

Two-Player Games

Given a propositional formula G with variables $p_1, q_1, \dots, p_n, q_n$.

There are two players: P and Q .

At step k each player makes a move:

Two-Player Games

Given a propositional formula G with variables $p_1, q_1, \dots, p_n, q_n$.

There are two players: P and Q .

At step k each player makes a move:

1. the player P can choose a boolean value for the variable p_k ;

Two-Player Games

Given a propositional formula G with variables $p_1, q_1, \dots, p_n, q_n$.

There are two players: P and Q .

At step k each player makes a move:

1. the player P can choose a boolean value for the variable p_k ;
2. the player Q can choose a boolean value for the variable q_k .

Two-Player Games

Given a propositional formula G with variables $p_1, q_1, \dots, p_n, q_n$.

There are two players: P and Q .

At step k each player makes a move:

1. the player P can choose a boolean value for the variable p_k ;
2. the player Q can choose a boolean value for the variable q_k .

The player P wins if after n steps the chosen values make the formula G true.

The player Q wins if after n steps the chosen values make the formula G false.

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1
2. $p_1 \rightarrow q_1$

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1
2. $p_1 \rightarrow q_1$
3. $q_1 \rightarrow q_1$

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1
2. $p_1 \rightarrow q_1$
3. $q_1 \rightarrow q_1$ If G is valid, P always wins!

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1
2. $p_1 \rightarrow q_1$
3. $q_1 \rightarrow q_1$ If G is valid, P always wins!
4. $p_1 \wedge \neg p_1$

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1
2. $p_1 \rightarrow q_1$
3. $q_1 \rightarrow q_1$ If G is valid, P always wins!
4. $p_1 \wedge \neg p_1$ If G is unsatisfiable, Q always wins!

Suppose Both Players Make no Errors: Who Wins?

Consider several special cases

1. p_1
2. $p_1 \rightarrow q_1$
3. $q_1 \rightarrow q_1$ If G is valid, P always wins!
4. $p_1 \wedge \neg p_1$ If G is unsatisfiable, Q always wins!
5. $p_1 \leftrightarrow q_1$

Winning Strategy

Problem: does P have a winning strategy?

Winning Strategy

Problem: does P have a winning strategy?

He has a winning strategy if

- ▶ there exists a move for P (a boolean value for p_1) such that

Winning Strategy

Problem: does P have a winning strategy?

He has a winning strategy if

- ▶ there exists a move for P (a boolean value for for p_1) such that
- ▶ for all moves of Q (boolean values for for q_1)

Winning Strategy

Problem: does P have a winning strategy?

He has a winning strategy if

- ▶ there exists a move for P (a boolean value for for p_1) such that
- ▶ for all moves of Q (boolean values for for q_1)
- ▶ there exists a move for P (a boolean value for for p_2) such that

Winning Strategy

Problem: does P have a winning strategy?

He has a winning strategy if

- ▶ there exists a move for P (a boolean value for for p_1) such that
- ▶ for all moves of Q (boolean values for for q_1)
- ▶ there exists a move for P (a boolean value for for p_2) such that
- ▶ for all moves of Q (boolean values for for q_2)

Winning Strategy

Problem: does P have a winning strategy?

He has a winning strategy if

- ▶ there exists a move for P (a boolean value for for p_1) such that
- ▶ for all moves of Q (boolean values for for q_1)
- ▶ there exists a move for P (a boolean value for for p_2) such that
- ▶ for all moves of Q (boolean values for for q_2)
- ▶ ...
- ▶ for all moves of Q (boolean values for for q_n) the formula G becomes true.

Winning Strategy

Problem: does P have a winning strategy?

He has a winning strategy if

- ▶ there exists a move for P (a boolean value for for p_1) such that
- ▶ for all moves of Q (boolean values for for q_1)
- ▶ there exists a move for P (a boolean value for for p_2) such that
- ▶ for all moves of Q (boolean values for for q_2)
- ▶ ...
- ▶ for all moves of Q (boolean values for for q_n) the formula G becomes true.

The existence of a winning strategy can be expressed by a quantified boolean formula $\exists p_1 \forall q_1 \exists p_2 \forall q_2 \dots \exists p_n \forall q_n G$.

Quantified Boolean Formulas

Propositional formula:

- ▶ Every boolean variable is a formula.
- ▶ \top and \perp are formulas.
- ▶ If F_1, \dots, F_n are formulas, where $n \geq 2$, then $(F_1 \wedge \dots \wedge F_n)$ and $(F_1 \vee \dots \vee F_n)$ are formulas.
- ▶ If F is a formula, then $\neg F$ is a formula.
- ▶ If F and G are formulas, then $(F \rightarrow G)$ and $(F \leftrightarrow G)$ are formulas.

Quantified Boolean Formulas

Propositional formula:

- ▶ Every boolean variable is a formula.
- ▶ \top and \perp are formulas.
- ▶ If F_1, \dots, F_n are formulas, where $n \geq 2$, then $(F_1 \wedge \dots \wedge F_n)$ and $(F_1 \vee \dots \vee F_n)$ are formulas.
- ▶ If F is a formula, then $\neg F$ is a formula.
- ▶ If F and G are formulas, then $(F \rightarrow G)$ and $(F \leftrightarrow G)$ are formulas.

Quantified boolean formulas:

- ▶ If p is a boolean variable and F is a formula, then $\forall p F$ and $\exists p F$ are formulas.

Quantifiers

- ▶ \forall is called the universal quantifier.
- ▶ \exists is called the existential quantifier.
- ▶ Read $\forall pF$ as “for all p , F ”.
- ▶ Read $\exists pF$ as “there exists p such that F ” or “for some p , F ”.

Connective	Name	Precedence
\forall	for all	5
\exists	exists	5
\neg	negation	5
\wedge	conjunction	4
\vee	disjunction	3
\rightarrow	implication	2
\leftrightarrow	equivalence	1

New Notation

Define

$$I_p^b(q) \stackrel{\text{def}}{=} \begin{cases} I(q), & \text{if } p \neq q; \\ b, & \text{if } p = q. \end{cases}$$

Example: let $I = \{p \mapsto 1, q \mapsto 0, r \mapsto 1\}$. Then

$$\begin{aligned} I_q^1 &= \{p \mapsto 1, q \mapsto 1, r \mapsto 1\} \\ I_q^0 &= \{p \mapsto 1, q \mapsto 0, r \mapsto 1\} = I \\ I_p^0 &= \{p \mapsto 0, q \mapsto 0, r \mapsto 1\} \end{aligned}$$

Semantics

1. $I(\top) = 1$ and $I(\perp) = 0$.
2. $I(F_1 \wedge \dots \wedge F_n) = 1$ if and only if $I(F_i) = 1$ for all i .
3. $I(F_1 \vee \dots \vee F_n) = 1$ if and only if $I(F_i) = 1$ for some i .
4. $I(\neg F) = 1$ if and only if $I(F) = 0$.
5. $I(F \rightarrow G) = 1$ if and only if $I(F) = 0$ or $I(G) = 1$.
6. $I(F \leftrightarrow G) = 1$ if and only if $I(F) = I(G)$.
7. $I(\forall p F) = 1$ if and only if $I_p^0(F) = 1$ and $I_p^1(F) = 1$.
8. $I(\exists p F) = 1$ if and only if $I_p^0(F) = 1$ or $I_p^1(F) = 1$.

Semantics

1. $I(\top) = 1$ and $I(\perp) = 0$.
2. $I(F_1 \wedge \dots \wedge F_n) = 1$ if and only if $I(F_i) = 1$ for all i .
3. $I(F_1 \vee \dots \vee F_n) = 1$ if and only if $I(F_i) = 1$ for some i .
4. $I(\neg F) = 1$ if and only if $I(F) = 0$.
5. $I(F \rightarrow G) = 1$ if and only if $I(F) = 0$ or $I(G) = 1$.
6. $I(F \leftrightarrow G) = 1$ if and only if $I(F) = I(G)$.
7. $I(\forall p F) = 1$ if and only if $I_p^0(F) = 1$ and $I_p^1(F) = 1$.
8. $I(\exists p F) = 1$ if and only if $I_p^0(F) = 1$ or $I_p^1(F) = 1$.

Evaluating a Formula: and-or trees

Let us evaluate $\forall p \exists q (p \leftrightarrow q)$ on the interpretation $\{p \mapsto 1, q \mapsto 0\}$.

Evaluating a Formula: and-or trees

Let us evaluate $\forall p \exists q (p \leftrightarrow q)$ on the interpretation $\{p \mapsto 1, q \mapsto 0\}$.

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $I_{b_1 b_2}$.

$$I_{10} \models \forall p \exists q (p \leftrightarrow q)$$

Evaluating a Formula: and-or trees

Let us evaluate $\forall p \exists q (p \leftrightarrow q)$ on the interpretation $\{p \mapsto 1, q \mapsto 0\}$.

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $l_{b_1 b_2}$.

$$l_{10} \models \forall p \exists q (p \leftrightarrow q) \quad \Leftrightarrow$$

$l_{00} \models \exists q (p \leftrightarrow q)$	and
$l_{10} \models \exists q (p \leftrightarrow q)$	

Evaluating a Formula: and-or trees

Let us evaluate $\forall p \exists q (p \leftrightarrow q)$ on the interpretation $\{p \mapsto 1, q \mapsto 0\}$.

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $l_{b_1 b_2}$.

$$l_{10} \models \forall p \exists q (p \leftrightarrow q) \quad \Leftrightarrow$$

$$\begin{array}{l} l_{00} \models \exists q (p \leftrightarrow q) \\ l_{10} \models \exists q (p \leftrightarrow q) \end{array} \quad \text{and}$$

\Leftrightarrow

$$\begin{array}{l} l_{00} \models p \leftrightarrow q \\ l_{01} \models p \leftrightarrow q \end{array} \quad \text{or}$$


and

Evaluating a Formula: and-or trees

Let us evaluate $\forall p \exists q (p \leftrightarrow q)$ on the interpretation $\{p \mapsto 1, q \mapsto 0\}$.
Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $l_{b_1 b_2}$.

$$\begin{aligned} l_{10} \models \forall p \exists q (p \leftrightarrow q) &\Leftrightarrow \boxed{\begin{array}{l} l_{00} \models \exists q (p \leftrightarrow q) \\ l_{10} \models \exists q (p \leftrightarrow q) \end{array} \text{ and}} \\ &\Leftrightarrow \boxed{\begin{array}{l} \boxed{\begin{array}{l} l_{00} \models p \leftrightarrow q \\ l_{01} \models p \leftrightarrow q \end{array} \text{ or}} \\ \boxed{\begin{array}{l} l_{10} \models p \leftrightarrow q \\ l_{11} \models p \leftrightarrow q \end{array} \text{ or}} \end{array} \text{ and}} \end{aligned}$$

Evaluating a formula



Evaluating a formula

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $I_{b_1 b_2}$. Use wildcards $*$ to denote “any” boolean value.

$$I_{**} \models \forall p \exists q (p \leftrightarrow q)$$

Evaluating a formula

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $I_{b_1 b_2}$. Use wildcards $*$ to denote “any” boolean value.

$$I_{**} \models \forall p \exists q (p \leftrightarrow q) \quad \Leftrightarrow$$

$I_{0*} \models \exists q (p \leftrightarrow q)$	and
$I_{1*} \models \exists q (p \leftrightarrow q)$	

Evaluating a formula

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $I_{b_1 b_2}$. Use wildcards $*$ to denote “any” boolean value.

$$I_{**} \models \forall p \exists q (p \leftrightarrow q) \quad \Leftrightarrow$$

$$\begin{array}{l} I_{0*} \models \exists q (p \leftrightarrow q) \\ I_{1*} \models \exists q (p \leftrightarrow q) \end{array} \quad \text{and}$$

$$\Leftrightarrow$$

$$\begin{array}{l} I_{00} \models p \leftrightarrow q \\ I_{01} \models p \leftrightarrow q \end{array} \quad \text{or}$$

and

$$\begin{array}{l} I_{10} \models p \leftrightarrow q \\ I_{11} \models p \leftrightarrow q \end{array} \quad \text{or}$$

Evaluating a formula

Denote any interpretation $\{p \mapsto b_1, q \mapsto b_2\}$ by $I_{b_1 b_2}$. Use wildcards $*$ to denote “any” boolean value.

$$\begin{aligned} I_{**} \models \forall p \exists q (p \leftrightarrow q) &\Leftrightarrow \boxed{\begin{array}{l} I_{0*} \models \exists q (p \leftrightarrow q) \\ I_{1*} \models \exists q (p \leftrightarrow q) \end{array} \text{ and}} \\ &\Leftrightarrow \boxed{\begin{array}{l} \boxed{\begin{array}{l} I_{00} \models p \leftrightarrow q \\ I_{01} \models p \leftrightarrow q \end{array} \text{ or}} \\ \boxed{\begin{array}{l} I_{10} \models p \leftrightarrow q \\ I_{11} \models p \leftrightarrow q \end{array} \text{ or}} \end{array} \text{ and}} \end{aligned}$$

The variables p and q are **bound** by quantifiers $\forall p$ and $\exists q$, so the value of the formula does not depend on these variables.

Subformula

Propositional formulas:

- ▶ The formulas F_1, \dots, F_n are the immediate subformulas of the formulas $F_1 \wedge \dots \wedge F_n$ and $F_1 \vee \dots \vee F_n$.
- ▶ The formulas F is the immediate subformula of the formula $\neg F$.
- ▶ The formulas F_1, F_2 are the immediate subformulas of the formulas $F_1 \rightarrow F_2$ and $F_1 \leftrightarrow F_2$.
- ▶ ...

Subformula

Propositional formulas:

- ▶ The formulas F_1, \dots, F_n are the immediate subformulas of the formulas $F_1 \wedge \dots \wedge F_n$ and $F_1 \vee \dots \vee F_n$.
- ▶ The formulas F is the immediate subformula of the formula $\neg F$.
- ▶ The formulas F_1, F_2 are the immediate subformulas of the formulas $F_1 \rightarrow F_2$ and $F_1 \leftrightarrow F_2$.
- ▶ ...

Quantified boolean formulas:

- ▶ The formula F_1 is the immediate subformula of the formulas $\forall p F_1$ and $\exists p F_1$.

Positions and Polarity

Let $F|_{\pi} = G$.

Propositional formulas:

- ▶ If G has the form $G_1 \wedge \dots \wedge G_n$ or $G_1 \vee \dots \vee G_n$, then for all $i \in \{1, \dots, n\}$ the position $\pi.i$ is a position in F and $pol(F, \pi.i) \stackrel{\text{def}}{=} pol(F, \pi)$.
- ▶ If G has the form $\neg G_1$, then $\pi.1$ is a position in F , $F|_{\pi.1} \stackrel{\text{def}}{=} G_1$ and $pol(F, \pi.1) \stackrel{\text{def}}{=} -pol(F, \pi)$.
- ▶ ...

Positions and Polarity

Let $F|_{\pi} = G$.


Propositional formulas:

- ▶ If G has the form $G_1 \wedge \dots \wedge G_n$ or $G_1 \vee \dots \vee G_n$, then for all $i \in \{1, \dots, n\}$ the position $\pi.i$ is a position in F and $pol(F, \pi.i) \stackrel{\text{def}}{=} pol(F, \pi)$.
- ▶ If G has the form $\neg G_1$, then $\pi.1$ is a position in F , $F|_{\pi.1} \stackrel{\text{def}}{=} G_1$ and $pol(F, \pi.1) \stackrel{\text{def}}{=} -pol(F, \pi)$.
- ▶ ...


Quantified boolean formulas:

- ▶ If G has the form $\forall p G_1$ or $\exists p G_1$, then $\pi.1$ is a position in F , $F|_{\pi.1} \stackrel{\text{def}}{=} G_1$ and $pol(F, \pi.1) \stackrel{\text{def}}{=} pol(F, \pi)$.


Example



Example



Example



Free and bound occurrences of variables

Let p be a boolean variable and $F|_{\pi} = p$.

- ▶ The occurrence of p at the position π in F is **bound** if π can be represented as a concatenation of two strings $\pi_1\pi_2$ such that $F|_{\pi_1}$ has the form $\forall pG$ or $\exists pG$ for some G .

Free and bound occurrences of variables

Let p be a boolean variable and $F|_{\pi} = p$.

- ▶ The occurrence of p at the position π in F is **bound** if π can be represented as a concatenation of two strings $\pi_1\pi_2$ such that $F|_{\pi_1}$ has the form $\forall pG$ or $\exists pG$ for some G .
In other words, a bound occurrence of p is an occurrence **in the scope of $\forall p$ or $\exists p$** .

Free and bound occurrences of variables

Let p be a boolean variable and $F|_{\pi} = p$.

- ▶ The occurrence of p at the position π in F is **bound** if π can be represented as a concatenation of two strings $\pi_1\pi_2$ such that $F|_{\pi_1}$ has the form $\forall pG$ or $\exists pG$ for some G .
In other words, a bound occurrence of p is an occurrence **in the scope of $\forall p$ or $\exists p$** .
- ▶ **Free occurrence**: not bound.

Free and bound occurrences of variables

Let p be a boolean variable and $F|_{\pi} = p$.

- ▶ The occurrence of p at the position π in F is **bound** if π can be represented as a concatenation of two strings $\pi_1\pi_2$ such that $F|_{\pi_1}$ has the form $\forall pG$ or $\exists pG$ for some G .
In other words, a bound occurrence of p is an occurrence **in the scope of $\forall p$ or $\exists p$** .
- ▶ **Free occurrence**: not bound.
- ▶ **Free (bound) variable** of a formula: a variable with at least one free (bound) occurrence.

Free and bound occurrences of variables

Let p be a boolean variable and $F|_{\pi} = p$.


- ▶ The occurrence of p at the position π in F is **bound** if π can be represented as a concatenation of two strings $\pi_1\pi_2$ such that $F|_{\pi_1}$ has the form $\forall pG$ or $\exists pG$ for some G .
In other words, a bound occurrence of p is an occurrence **in the scope of $\forall p$ or $\exists p$** .
- ▶ **Free occurrence**: not bound.
- ▶ **Free (bound) variable** of a formula: a variable with at least one free (bound) occurrence.
- ▶ **Warning** The same variable can be **both free and bound** in a formula.

Free and bound occurrences of variables

Let p be a boolean variable and $F|_{\pi} = p$.

- ▶ The occurrence of p at the position π in F is **bound** if π can be represented as a concatenation of two strings $\pi_1\pi_2$ such that $F|_{\pi_1}$ has the form $\forall pG$ or $\exists pG$ for some G .
In other words, a bound occurrence of p is an occurrence **in the scope of $\forall p$ or $\exists p$** .
- ▶ **Free occurrence**: not bound.
- ▶ **Free (bound) variable** of a formula: a variable with at least one free (bound) occurrence.
- ▶ **Warning** The same variable can be **both free and bound** in a formula.
- ▶ **Closed formula**: formula with no free variables.

Example: Free and Bound Variables



Only Free Variables Matter

The truth value of a formula depends only on the truth values of free variables of the formula:

Lemma

Let *for all free variables* p of a formula F we have $I_1(p) = I_2(p)$. Then $I_1 \models F$ if and only if $I_2 \models F$.

Only Free Variables Matter

The truth value of a formula depends only on the truth values of free variables of the formula:

Lemma

Let *for all free variables* p of a formula F we have $l_1(p) = l_2(p)$. Then $l_1 \models F$ if and only if $l_2 \models F$.

Theorem

Let F be a closed formula and l_1, l_2 be interpretations. Then $l_1 \models F$ if and only if $l_2 \models F$.

Truth, Validity and Satisfiability

Validity and satisfiability are defined as for propositional formulas.

Truth, Validity and Satisfiability

Validity and satisfiability are defined as for propositional formulas.

There is no difference between these notions for closed formulas:

Lemma

For every interpretation I and closed formula F the following propositions are equivalent: (i) $I \models F$; (ii) F is satisfiable; and (iii) F is valid.

Truth, Validity and Satisfiability

Validity and **satisfiability** are defined as for propositional formulas.

There is no difference between these notions for closed formulas:

Lemma

*For every interpretation I and **closed formula** F the following propositions are equivalent: (i) $I \models F$; (ii) F is satisfiable; and (iii) F is valid.*

Validity and satisfiability can be expressed through truth:

Lemma

Let F be a formula with free variables p_1, \dots, p_n .

- ▶ F is satisfiable if and only if the formula $\exists p_1 \dots \exists p_n F$ is satisfiable (true, valid).
- ▶ F is valid if and only if the formula $\forall p_1 \dots \forall p_n F$ is valid (true, satisfiable).

Substitutions for propositional formulas

Substitution: $(F)_p^G$: denotes the formula obtained from F by replacing all occurrences of the variable p by G .

Example:

$$\begin{aligned} ((p \vee s) \wedge (q \rightarrow p))_p^{(I \wedge s)} = \\ ((I \wedge s) \vee s) \wedge (q \rightarrow (I \wedge s)) \end{aligned}$$

Properties: If we apply **any substitution** to a **valid** formula then we also obtain a **valid** formula.

Substitutions for propositional formulas

Substitution: $(F)_p^G$: denotes the formula obtained from F by replacing all occurrences of the variable p by G .

Example:

$$\begin{aligned} & ((p \vee s) \wedge (q \rightarrow p))_p^{(I \wedge s)} = \\ & (((I \wedge s) \vee s) \wedge (q \rightarrow (I \wedge s))) \end{aligned}$$

Properties: If we apply any substitution to a valid formula then we also obtain a valid formula.

Substitutions for propositional formulas

Substitution: $(F)_p^G$: denotes the formula obtained from F by replacing all occurrences of the variable p by G .

Example:

$$\begin{aligned} & ((p \vee s) \wedge (q \rightarrow p))_p^{(I \wedge s)} = \\ & (((I \wedge s) \vee s) \wedge (q \rightarrow (I \wedge s))) \end{aligned}$$

Properties: If we apply **any substitution** to a **valid** formula then we also obtain a **valid** formula.

Substitution for quantified formulas

Some problems...

Substitution for quantified formulas

Some problems...

Consider $\exists q(\neg p \leftrightarrow q)$.

Substitution for quantified formulas

Some problems...

Consider $\exists q(\neg p \leftrightarrow q)$.

We cannot simply replace variables by formulas any more:

$\exists(r \rightarrow r)(\neg p \leftrightarrow r \rightarrow r)$???

Substitution for quantified formulas

Some problems...

Consider $\exists q(\neg p \leftrightarrow q)$.

We cannot simply replace variables by formulas any more:

$\exists(r \rightarrow r)(\neg p \leftrightarrow r \rightarrow r)$???

Free variables are **parameters**: we can only substitute for parameters.

But a variable can have both **free and bound** occurrences in a formula, e.g. $(\forall p p \rightarrow q) \wedge (q \vee (q \rightarrow p))$

Renaming bound variables

Notation: $\exists\forall$: any of \exists, \forall and $\forall\exists$: any of \forall, \exists .

Renaming bound variables

Notation: $\exists\forall$: any of \exists, \forall and $\forall\exists$: any of \forall, \exists .

Renaming bound variables in F :

Let $F[\exists\forall pG]$.

1. Take a **fresh** variable q (that is a variable **not occurring** in F);
2. Replace all free occurrences of p in G (note: not in F !) by q obtaining G' .
3. So we obtain the $F[\exists\forall qG']$ as the result.

Renaming bound variables

Notation: $\exists\forall$: any of \exists, \forall and $\forall\exists$: any of \wedge, \vee .

Renaming bound variables in F :

Let $F[\exists p G]$.

1. Take a **fresh** variable q (that is a variable **not occurring** in F);
2. Replace all free occurrences of p in G (note: not in F !) by q obtaining G' .
3. So we obtain the $F[\exists q G']$ as the result.

Lemma

$$F[\exists p G] \equiv F[\exists q G']$$

Renaming bound variables

Notation: $\exists\forall$: any of \exists, \forall and $\forall\exists$: any of \forall, \exists .

Renaming bound variables in F :

Let $F[\exists p G]$.

1. Take a **fresh** variable q (that is a variable **not occurring** in F);
2. Replace all free occurrences of p in G (note: not in F !) by q obtaining G' .
3. So we obtain the $F[\exists q G']$ as the result.

Lemma

$$F[\exists p G] \equiv F[\exists q G']$$

Example:

$$\exists q(\forall p((p \rightarrow q) \wedge p)) \vee p.$$

Then we can rename p into r obtaining

$$\exists q(\forall r((r \rightarrow q) \wedge r)) \vee p.$$

Rectified formulas

Rectified formula F :

- ▶ no variable appears both free and bound in F ;
- ▶ for every variable p , the formula F contains at most one occurrence of quantifiers $\exists p$.

Any formula can be transformed into a rectified formula by renaming bound variables.

We can use the usual notation $(F)_p^G$ for rectified formulas assuming that p occurs only free.

Rectified formulas

Rectified formula F :

- ▶ no variable appears both free and bound in F ;
- ▶ for every variable p , the formula F contains at most one occurrence of quantifiers $\exists p$.

Any formula can be transformed into a rectified formula by renaming bound variables.

We can use the usual notation $(F)_p^G$ for rectified formulas assuming that p occurs only free.

Rectified formulas

Rectified formula F :

- ▶ no variable appears both free and bound in F ;
- ▶ for every variable p , the formula F contains at most one occurrence of quantifiers $\exists p$.

Any formula can be transformed into a rectified formula by renaming bound variables.

We can use the usual notation $(F)_p^G$ for rectified formulas assuming that p occurs only free.

Rectification: Example

$$p \rightarrow \exists p(p \wedge \forall p(p \vee r \rightarrow \neg p)) \Rightarrow$$

$$p \rightarrow \exists p(p \wedge \forall p_1(p_1 \vee r \rightarrow \neg p_1)) \Rightarrow$$

$$p \rightarrow \exists p_2(p_2 \wedge \forall p_1(p_1 \vee r \rightarrow \neg p_1))$$

Rectification: Example

$$p \rightarrow \exists p(p \wedge \forall p(p \vee r \rightarrow \neg p)) \Rightarrow$$

$$p \rightarrow \exists p(p \wedge \forall p_1(p_1 \vee r \rightarrow \neg p_1)) \Rightarrow$$

$$p \rightarrow \exists p_2(p_2 \wedge \forall p_1(p_1 \vee r \rightarrow \neg p_1))$$

Rectification: Example

$$p \rightarrow \exists p(p \wedge \forall p(p \vee r \rightarrow \neg p)) \Rightarrow$$

$$p \rightarrow \exists p(p \wedge \forall p_1(p_1 \vee r \rightarrow \neg p_1)) \Rightarrow$$

$$p \rightarrow \exists p_2(p_2 \wedge \forall p_1(p_1 \vee r \rightarrow \neg p_1))$$

Another problem

$\exists q(\neg p \leftrightarrow q)$: there exists a truth value equivalent to $\neg p$. This formula is valid.

Substitute p by q .

$\exists q(\neg q \leftrightarrow q)$: there exists a truth value equivalent to its own negation. This formula is unsatisfiable.

Another problem

$\exists q(\neg p \leftrightarrow q)$: there exists a truth value equivalent to $\neg p$. This formula is valid.

Substitute p by q .

$\exists q(\neg q \leftrightarrow q)$: there exists a truth value equivalent to its own negation. This formula is unsatisfiable.

Another restriction

Suppose we want to substitute $(F)_p^G$.

Then we **require**: no free variable in G become **bound** in $(F)_p^G$.

In previous example $\exists q(\neg p \leftrightarrow q)$:

Substitute p by q . ($\exists q(\neg q \leftrightarrow q)$ does not satisfy above)

Uniform solution – renaming of bound variables

$\exists q(\neg p \leftrightarrow q) \equiv \exists r(\neg p \leftrightarrow r)$

Now we can substitute p by q obtaining $\exists r(\neg q \leftrightarrow r)$

From now on we always assume that:

- ▶ formulas are **rectified**.
- ▶ all substitutions **satisfy the requirement** above

Another restriction

Suppose we want to substitute $(F)_p^G$.

Then we **require**: no free variable in G become **bound** in $(F)_p^G$.

In previous example $\exists q(\neg p \leftrightarrow q)$:

Substitute p by q . ($\exists q(\neg q \leftrightarrow q)$ does not satisfy above)

Uniform solution – renaming of bound variables

$$\exists q(\neg p \leftrightarrow q) \equiv \exists r(\neg p \leftrightarrow r)$$

Now we can substitute p by q obtaining $\exists r(\neg q \leftrightarrow r)$

From now on we always assume that:

- ▶ formulas are **rectified**.
- ▶ all substitutions **satisfy the requirement** above

Another restriction

Suppose we want to substitute $(F)_p^G$.

Then we **require**: no free variable in G become **bound** in $(F)_p^G$.

In previous example $\exists q(\neg p \leftrightarrow q)$:

Substitute p by q . ($\exists q(\neg q \leftrightarrow q)$ does not satisfy above)

Uniform solution – renaming of bound variables

$\exists q(\neg p \leftrightarrow q) \equiv \exists r(\neg p \leftrightarrow r)$

Now we can substitute p by q obtaining $\exists r(\neg q \leftrightarrow r)$

From now on we always assume that:

- ▶ formulas are **rectified**.
- ▶ all substitutions **satisfy the requirement** above

Another restriction

Suppose we want to substitute $(F)_p^G$.

Then we **require**: no free variable in G become **bound** in $(F)_p^G$.

In previous example $\exists q(\neg p \leftrightarrow q)$:

Substitute p by q . ($\exists q(\neg q \leftrightarrow q)$ does not satisfy above)

Uniform solution – renaming of bound variables

$$\exists q(\neg p \leftrightarrow q) \equiv \exists r(\neg p \leftrightarrow r)$$

Now we can substitute p by q obtaining $\exists r(\neg q \leftrightarrow r)$

From now on we always assume that:

- ▶ formulas are **rectified**.
- ▶ all substitutions **satisfy the requirement** above

Equivalent replacement

Lemma

Let I be an interpretation and $I \models F_1 \leftrightarrow F_2$. Then $I \models G[F_1] \leftrightarrow G[F_2]$.

Theorem (Equivalent Replacement)

Let $F_1 \equiv F_2$. Then $G[F_1] \equiv G[F_2]$.

Prenex form

- ▶ **Quantifier-free formula:** no quantifiers (that is, propositional).
- ▶ **Prenex formula** has the form $\exists_1 p_1 \dots \exists_n p_n G$, where G is quantifier-free.
- ▶ **Outermost prefix of $\exists_1 p_1 \dots \exists_n p_n G$:** the longest subsequence $\exists_1 p_1 \dots \exists_k p_k$ of $\exists_1 p_1 \dots \exists_n p_n$ such that $\exists_1 = \dots = \exists_k$.
- ▶ A formula F is a **prenex form of a formula G** if F is prenex and $F \equiv G$.

Prenex form

- ▶ **Quantifier-free formula:** no quantifiers (that is, propositional).
- ▶ **Prenex formula** has the form $\exists_1 p_1 \dots \exists_n p_n G$, where G is quantifier-free.
- ▶ **Outermost prefix of $\exists_1 p_1 \dots \exists_n p_n G$:** the longest subsequence $\exists_1 p_1 \dots \exists_k p_k$ of $\exists_1 p_1 \dots \exists_n p_n$ such that $\exists_1 = \dots = \exists_k$.
- ▶ A formula F is a **prenex form of a formula G** if F is prenex and $F \equiv G$.

Prenex form

- ▶ **Quantifier-free formula:** no quantifiers (that is, propositional).
- ▶ **Prenex formula** has the form $\exists_1 p_1 \dots \exists_n p_n G$, where G is quantifier-free.
- ▶ **Outermost prefix of $\exists_1 p_1 \dots \exists_n p_n G$:** the longest subsequence $\exists_1 p_1 \dots \exists_k p_k$ of $\exists_1 p_1 \dots \exists_n p_n$ such that $\exists_1 = \dots = \exists_k$.
- ▶ A formula F is a **prenex form of a formula G** if F is prenex and $F \equiv G$.

Prenex form

- ▶ **Quantifier-free formula:** no quantifiers (that is, propositional).
- ▶ **Prenex formula** has the form $\exists_1 p_1 \dots \exists_n p_n G$, where G is quantifier-free.
- ▶ **Outermost prefix of $\exists_1 p_1 \dots \exists_n p_n G$:** the longest subsequence $\exists_1 p_1 \dots \exists_k p_k$ of $\exists_1 p_1 \dots \exists_n p_n$ such that $\exists_1 = \dots = \exists_k$.
- ▶ A formula F is a **prenex form of a formula G** if F is prenex and $F \equiv G$.

Prenexing rules

We assume that the formula **rectified** before application of

Prenexing rules:

$$\exists \forall p F_1 \bowtie \dots \bowtie F_n \Rightarrow \exists \forall p (F_1 \bowtie \dots \bowtie F_n)$$

$$F_1 \leftrightarrow F_2 \Rightarrow (F_1 \rightarrow F_2) \wedge (F_2 \rightarrow F_1)$$

$$\forall p F_1 \rightarrow F_2 \Rightarrow \exists p (F_1 \rightarrow F_2) \quad \exists p F_1 \rightarrow F_2 \Rightarrow \forall p (F_1 \rightarrow F_2)$$

$$F_1 \rightarrow \forall p F_2 \Rightarrow \forall p (F_1 \rightarrow F_2) \quad F_1 \rightarrow \exists p F_2 \Rightarrow \exists p (F_1 \rightarrow F_2)$$

$$\neg \forall p F \Rightarrow \exists p \neg F$$

$$\neg \exists p F \Rightarrow \forall p \neg F$$

Warning: Sound only when the formula is **rectified**!

Some useful equivalences: $\neg \forall p F \equiv \exists p \neg F$ and $\neg \exists p F \equiv \forall p \neg F$

Prenexing. Example I

$$\begin{aligned} & \exists q(q \rightarrow p) \rightarrow \neg \forall r(r \rightarrow p) \vee p \Rightarrow \\ & \forall q((q \rightarrow p) \rightarrow \neg \forall r(r \rightarrow p) \vee p) \Rightarrow \\ & \forall q((q \rightarrow p) \rightarrow \exists r \neg(r \rightarrow p) \vee p) \Rightarrow \\ & \forall q((q \rightarrow p) \rightarrow \exists r(\neg(r \rightarrow p) \vee p)) \Rightarrow \\ & \forall q \exists r((q \rightarrow p) \rightarrow \neg(r \rightarrow p) \vee p). \end{aligned}$$

Prenexing. Example II

$$\exists q(q \rightarrow p) \rightarrow \neg \forall r(r \rightarrow p) \vee p \Rightarrow$$

$$\exists q(q \rightarrow p) \rightarrow \exists r \neg(r \rightarrow p) \vee p \Rightarrow$$

$$\exists q(q \rightarrow p) \rightarrow \exists r(\neg(r \rightarrow p) \vee p) \Rightarrow$$

$$\exists r(\exists q(q \rightarrow p) \rightarrow \neg(r \rightarrow p) \vee p) \Rightarrow$$

$$\exists r \forall q((q \rightarrow p) \rightarrow \neg(r \rightarrow p) \vee p).$$

Summary

- ▶ quantified boolean formulas (QBF): $\exists x \forall y \exists z F$
- ▶ syntax, semantics
- ▶ evaluating QBF formula on an interpretations: and-or trees
- ▶ positions/polarity
- ▶ bound/free occurrences of variables
- ▶ for closed formulas: validity and satisfiability coincide;
for open formulas we can express satisfiability/validity using \exists/\forall quantifiers respectively.
- ▶ rectified formulas: i) no variable occurs free and bound, ii) every variable is quantified at most once.
- ▶ rectification: rename bound variables
- ▶ prenex normal form: all quantifiers are on the left-hand-side
- ▶ prenexing: rectify + apply prenexing rules