

Calculus III

UT Knoxville, Spring 2023, MATH 341

Stella Thistlethwaite, Alex Zhang

February 17, 2023

Contents

1	Introduction	2
2	Three-Dimensional Space	3
2.1	Points	3
2.2	Vectors	4
2.3	Gradient	7
2.4	Projecting Vectors	8
2.5	Cross Product	9
2.6	Torque	10
2.7	Cylinders and Quadratic Surfaces	11
3	Vector Functions	13
3.1	Limits and Continuity	13
	Index	15

Introduction

Much of our focus will be on Stoke's Theorem.

Three-Dimensional Space

In past math classes, we have been used to dealing in \mathbb{R}^2 where we work with two degrees of freedom: x and y . Now, we will be working in \mathbb{R}^3 with three degrees of freedom: x , y , and z .

2.1 Points

Definition 2.1.1 ► Point

A **point** in \mathbb{R}^n space is an n -tuple that specifies a location in that space.

$$p = (p_1, \dots, p_n) \in \mathbb{R}^n$$

Definition 2.1.2 ► Distance

Given two points $a, b \in \mathbb{R}^n$, the **distance** between the two points is defined as:

$$d(a, b) := \sqrt{(b_1 - a_1)^2 + \dots + (b_n - a_n)^2}$$

Example 2.1.1 ► Distance Between Points

Find the distance between $p_1 = (-1, -1, 4)$ and $p_2 = (-1, 4, -1)$.

$$\begin{aligned} d(p_1, p_2) &= \sqrt{(-1 - (-1))^2 + (4 - (-1))^2 + (-1 - 1)^2} \\ &= \sqrt{0^2 + 5^2 + (-5)^2} \\ &= \sqrt{50} \end{aligned}$$

Definition 2.1.3 ► Sphere

Given a point $c = (h, k, l) \in \mathbb{R}^3$, a **sphere** is the set of all points $(x, y, z) \in \mathbb{R}^3$ that are a distance r from the point $c = (h, k, l)$.

$$(x - h)^2 + (y - k)^2 + (z - l)^2 = r^2$$

Note that all the points of the sphere are equidistant to the center of the sphere. This means

the sphere is really a hollow shell.

Example 2.1.2 ► Circle

Show that the following quadratic equation represents a circle by rewriting it in standard form. Find the center $c = (h, k)$ and the radius r .

$$x^2 + y^2 + x = 0$$

To solve this, we will have to complete the square:

$$\begin{aligned} x^2 + x + y^2 &= 0 \\ \implies x^2 + x + \frac{1}{4} + y^2 &= \frac{1}{4} \\ \implies \left(x + \frac{1}{2}\right)^2 + y^2 &= \frac{1}{4} \end{aligned}$$

2.2 Vectors

Definition 2.2.1 ► Vector

A **vector** is a mathematical object that contains multiple objects of the same type.

$$\vec{v} = \langle v_1, \dots, v_n \rangle \in \mathbb{R}^n$$

As customary in most mathematics textbooks, we will always denote vectors using the little arrow thing. In the context of three-dimensional space, we will only be working with vectors with three components. In addition, we will think of vectors as having a magnitude and direction.

Definition 2.2.2 ► Scalar Multiplication

Given a vector \vec{v} and scalar k , we define **scalar multiplication** as:

$$k \cdot \vec{v} := \langle kv_1, \dots, kv_n \rangle$$

Note that scalar multiplication is associative, commutative, and distributive.

- $a(b\vec{v}) = b(a(\vec{v})) = (ab)\vec{v}$
- $(k_1 + k_2)\vec{v} = k_1\vec{v} + k_2\vec{v}$

- $k(\vec{v} + \vec{w}) = kv + k\vec{w}$

Definition 2.2.3 ► Norm

A vector's **norm** is its magnitude or length.

$$\|v\| := \sqrt{v_1^2 + \cdots + v_n^2}$$

Definition 2.2.4 ► Unit Vector

A **unit vector** is a vector whose magnitude is 1.

We will introduce shorthand notation for the three standard unit vectors:

- $\hat{i} := \langle 1, 0, 0 \rangle$
- $\hat{j} := \langle 0, 1, 0 \rangle$
- $\hat{k} := \langle 0, 0, 1 \rangle$

These three vectors form the **standard basis** for \mathbb{R}^3 . That is, we can express any vector in \mathbb{R}^3 as a linear combination of $\hat{i}, \hat{j}, \hat{k}$.

Technique 2.2.1 ► Finding a Unit Vector from a Given Vector

Given a vector $\vec{v} = \langle x, y, z \rangle \in \mathbb{R}^3$, we can find the **unit vector** \vec{u} with the same direction by:

$$\vec{u} = \frac{\vec{v}}{\|\vec{v}\|} = \left\langle \frac{x}{\|\vec{v}\|}, \frac{y}{\|\vec{v}\|}, \frac{z}{\|\vec{v}\|} \right\rangle$$

Definition 2.2.5 ► Dot Product

Given two vectors \vec{a} and \vec{b} whose cardinality are both n , we define the **dot product** of \vec{a} and \vec{b} as:

$$\vec{a} \cdot \vec{b} := a_1b_1 + \cdots + a_nb_n$$

Like scalar multiplication, dot product is also associative, commutative, and distributive.

Theorem 2.2.1 ► Angle Between Vectors

If \vec{a} and \vec{b} are vectors and θ is the angle between \vec{a} and \vec{b} , then:

$$\vec{a} \cdot \vec{b} = \|\vec{a}\| \|\vec{b}\| \cdot \cos(\theta)$$

Proof. TODO: finish proof

**Definition 2.2.6 ► Parallel, Perpendicular**

- Two vectors are **parallel** if the angle between the vectors is 0 deg.
- Two vectors are **perpendicular** if the angle between the vectors is 90 deg.

Definition 2.2.7 ► Orthogonal

\vec{a} and \vec{b} are **orthogonal** if $\vec{a} \cdot \vec{b} = 0$.

Given a vector $\vec{a} = \langle a_1, a_2, a_3 \rangle$, we have:

$$\frac{\vec{a}}{\|\vec{a}\|} = \langle \cos \alpha, \cos \beta, \cos \gamma \rangle$$

where:

- $\alpha = \cos^{-1} \left(\frac{a_1}{\|\vec{a}\|} \right)$ (angle between \vec{a} and the x -axis)
- $\beta = \cos^{-1} \left(\frac{a_2}{\|\vec{a}\|} \right)$ (angle between \vec{a} and the y -axis)
- $\gamma = \cos^{-1} \left(\frac{a_3}{\|\vec{a}\|} \right)$ (angle between \vec{a} and the z -axis)

Definition 2.2.8 ► Work

If F is a force moving a particle from a point P to a point Q , the **work** performed by the force is given by:

$$W = \vec{F} \cdot \vec{PQ}$$

Example 2.2.1 ► Finding Work

Find the work done by a force $\vec{F} = \langle 3, 4, 5 \rangle$ in moving an object from $p = (2, 1, 0)$ to $q = (4, 6, 2)$.

First, we find \vec{pq} as such:

$$\begin{aligned}\vec{pq} &= \langle 4 - 2, 6 - 1, 2 - 0 \rangle \\ &= \langle 2, 5, 2 \rangle\end{aligned}$$

Then, we can find work:

$$\begin{aligned}W &= \vec{F} \cdot \vec{PQ} \\ &= \langle 3, 4, 5 \rangle \cdot \langle 2, 5, 2 \rangle \\ &= 6 + 20 + 10 \\ &= 36\end{aligned}$$

2.3 Gradient

Definition 2.3.1 ► Gradient

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a function. The **gradient** of f is a function $\nabla f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by:

$$\nabla f(x_1, \dots, x_n) = \left\langle \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right\rangle$$

Example 2.3.1 ► Gradient

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a function defined by $f(T, L, \rho) = \frac{1}{2L} \sqrt{\frac{T}{\rho}}$

The gradient of $f(T, L, P)$ is denoted

$$\begin{aligned}\nabla f(T, L, \rho) &= \left\langle \frac{\partial f}{\partial T}, \frac{\partial f}{\partial L}, \frac{\partial f}{\partial \rho} \right\rangle \\ &= \left\langle \frac{1}{4L\sqrt{T\rho}}, -\frac{1}{2L^2} \sqrt{\frac{T}{\rho}}, -\frac{1}{4L} \sqrt{\frac{T}{\rho^3}} \right\rangle\end{aligned}$$

We can then calculate gradient as such:

$$\begin{aligned}\nabla f(2, 1, 1) &= \left\langle \frac{1}{4(1)\sqrt{(2)(1)}}, -\frac{1}{2(1)}\sqrt{\frac{2}{1}}, -\frac{1}{(4)(1)}\sqrt{\frac{2}{1}} \right\rangle \\ &= \left\langle \frac{1}{4\sqrt{2}}, -\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{4} \right\rangle\end{aligned}$$

Definition 2.3.2 ► Directional Derivative

The **directional derivative** of $f(x, y, z)$ in the direction of \vec{a} is defined as:

$$\nabla f(x, y, z) \cdot \frac{\vec{a}}{\|\vec{a}\|}$$

Example 2.3.2 ► Directional Derivative

If $f(x, y, z) = xy^2z^5$, find the directional derivative of $f(x, y, z)$ at the point $(1, 0, -2)$ in the direction of the unit vector $\vec{u} = \frac{\vec{a}}{\|\vec{a}\|}$, $\vec{a} = \langle 1, 2, -2 \rangle$.

For this, we calculate $\nabla f(1, 0, -1)$, then calculate the dot product of $\nabla f(1, 0, -1)$ with the unit vector $\vec{u} = \langle 1/3, 2/3, -2/3 \rangle$. Thus, the directional derivative of $f(x, y, z)$ at $(1, 0, -1)$ denoted by $Df(1, 0, -1)$ in the direction of \vec{u} is:

$$\begin{aligned}Df(1, 0, -1) &= \nabla f(1, 2, -2) \cdot \vec{u} \\ &= \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\rangle \cdot \vec{u} \\ &= \langle 0, 0, 0 \rangle \cdot \left\langle \frac{1}{3}, \frac{2}{3}, -\frac{2}{3} \right\rangle \\ &= 0\end{aligned}$$

2.4 Projecting Vectors

Projecting a vector onto another vector

Definition 2.4.1 ► Scalar Projection

Given \vec{a} and \vec{b} , the **scalar projection** of \vec{b} onto \vec{a} is the norm of the vector projection of \vec{b} onto \vec{a} .

$$\text{comp}_{\vec{a}} \vec{b} := \frac{\vec{b} \cdot \vec{a}}{\|\vec{a}\|}$$

Definition 2.4.2 ► Vector Projection

Given \vec{a} and \vec{b} that are non-zero vectors, the **vector projection** of \vec{b} onto the vector \vec{a} is defined by:

$$\text{proj}_{\vec{a}} \vec{b} := \text{comp}_{\vec{a}} \vec{b} \frac{\vec{a}}{\|\vec{a}\|}$$

2.5 Cross Product

Definition 2.5.1 ► Cross Product

Given two vectors $\vec{a}, \vec{b} \in \mathbb{R}^3$, the **cross product** of \vec{a} and \vec{b} is a vector that is orthogonal to both \vec{a} and \vec{b} .

$$\vec{a} \times \vec{b} := \vec{c} \quad \text{where} \quad \vec{a} \cdot \vec{c} = 0 \quad \text{and} \quad \vec{b} \cdot \vec{c} = 0$$

The cross product is exclusive to vectors in three dimensions.

Technique 2.5.1 ► Calculating Cross Product

Let $\vec{a} := \langle a_1, a_2, a_3 \rangle$ and $\vec{b} := \langle b_1, b_2, b_3 \rangle$ To find $\vec{a} \times \vec{b}$, we:

1. Create a matrix as such:

$$\begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix}$$

2. Find the determinant of the matrix by cofactor expansion on the first row.

$$\begin{aligned}\vec{a} \times \vec{b} &= \hat{i} \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} - \hat{j} \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} + \hat{k} \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \\ &= \hat{i}(a_2b_3 - b_2a_3) - \hat{j}(a_1b_3 - b_1a_3) + \hat{k}(a_1b_2 - b_1a_2) \\ &= \langle a_2b_3 - b_2a_3, -a_1b_3 + b_1a_3, a_1b_2 - b_1a_2 \rangle\end{aligned}$$

- $\vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$
- $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$
- $(\vec{a} + \vec{b}) \times \vec{c} = \vec{a} \times \vec{c} + \vec{b} \times \vec{c}$
- If $r \in \mathbb{R}$, then $(r\vec{a}) \times \vec{b} = \vec{a} \times (r\vec{b}) = r(\vec{a} \times \vec{b})$
- $\vec{a} \cdot (\vec{b} \times \vec{c}) = (\vec{a} \times \vec{b}) \cdot \vec{c}$
- $\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$

Theorem 2.5.1

If \vec{a} and \vec{b} are two non-zero vectors in \mathbb{R}^3 and θ is the angle between \vec{a} and \vec{b} , then:

$$\|\vec{a} \times \vec{b}\| = \|\vec{a}\|\|\vec{b}\| \sin \theta$$

2.6 Torque

Definition 2.6.1 ► Torque

If \vec{F} is a force applied to an object with position vector \vec{r} , then the torque \vec{T} produced by \vec{F} is given by:

$$\vec{T} := \vec{r} \times \vec{F}$$

2.7 Cylinders and Quadratic Surfaces

Definition 2.7.1 ► Planar Curve

A **planar curve** is any curve that lies on a single plane.

Definition 2.7.2 ► Cylinder

Given a planar curve c , the surface in \mathbb{R}^3 defined by all parallel lines crossing the curve c is called a **cylinder**.

Note that our broad definition of cylinder does not require the cylinder to be circular, nor does it require it to be straight. For example, we could have a planar curve defined by $x^2 + y^2 = 1$ and create a circular cylinder with radius 1. We could also have a planar curve defined by $y = x^2$ and create a **parabolic cylinder**.

Example 2.7.1

Consider the curve $x^2 + y^2 = z^2$. For every z_0 at $x = y = 0$, we have a point, say $p := (0, 0, 0)$.

Definition 2.7.3 ► Cone

Definition 2.7.4 ► Conic Surface

A **conic surface** is a surface that is attained by taking a cross-section of a cone.

There are four types of conic surfaces:

1. The cross-section parallel to the xy -plane is a **circle**.
2. The cross-section slightly angled from the xy -plane is a **ellipse**.
3. The cross-section parallel to a generating line is a **parabola**.
4. The cross-section parallel to the z axis is a **hyperbola**.

Definition 2.7.5 ► Quadratic Surface

A **quadratic surface** in \mathbb{R}^3 is the set of points whose coordinates satisfy a quadratic polynomial in the variables x, y, z .

For example, the standard equation for a sphere is a quadratic surface.

Definition 2.7.6 ► Ellipsoid

An **ellipsoid** is a quadratic surface in \mathbb{R}^3 defined by the equation:

$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} + \frac{(z-l)^2}{c^2} = 1$$

The cross-sections of an ellipsoid with each coordinate plane (xy -plane, xz -plane, yz -plane) is just an ellipse. In other words:

- If we set $z = l$, we get an ellipse in the xy -plane
- If we set $y = k$, we get an ellipse in the xz -plane
- If we set $x = h$, we get an ellipse in the yz -plane

We can also have **hyperboloids**:

- Type 1: $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$
- Type 2: $\frac{x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$

For type 2 hyperboloids, its cross-section with the xy and xz plane is a hyperbola. Note that there is no cross-section with the yz plane. This is because the equation $-\frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$ has no solution in the real numbers.

The standard form equation for a parabola is:

$$z - l = \pm c [(x - h)^2 + (y - k)^2]$$

TODO: more quadratic surfaces here

Vector Functions

Definition 3.0.1 ► Vector Function

A **vector function** $\vec{r}(t)$ is a function that maps each $t \in \mathbb{R}$ to a corresponding vector in \mathbb{R}^n . In other words:

$$\vec{r} : \mathbb{R} \rightarrow \mathbb{R}^n$$

For example, we can have $\vec{r}(t) = \langle 2 - 3t, 5 - 7t, t \rangle$. Then $\vec{r}(t)$ is a vector function that parameterizes a line that passes through $(2, 5, 0)$ and with vector direction $\vec{v} = \langle -3, -7, 1 \rangle$.

Example 3.0.1 ► Circle Vector Function

Let $\vec{r}_1(t) = \langle \cos t, \sin t \rangle$ where $0 \leq t < 2\pi$. This function's graph is a circle of radius 1. We can think of this as $\cos^2 t + \sin^2 t = 1$.

Similarly, consider $\vec{r}_2(t) = \langle 5 \cos t, 8 \sin t \rangle$. Thus, $x/5 = \cos t$ and $y/8 = \sin t$. Using the equation $\cos^2 t + \sin^2 t = 1$, we now have $(x/5)^2 + (y/8)^2 = 1$. It's an ellipse.

If a vector function is given by $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$, then the domain of $\vec{r}(t)$ is the intersection of the domains of f, g, h . This is denoted as $\text{Dom}(\vec{r}(t))$.

3.1 Limits and Continuity

If $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$ is a vector function, we say that:

$$\lim_{t \rightarrow a} \vec{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle$$

Recall L'Hopital's rule:

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$$

Only applies if the left-hand side is an indeterminate form (i.e. a fraction whose denominator is 0).

Definition 3.1.1 ▶ Continuity

We say a function $f : X \rightarrow Y$ is **continuous** at some value a if:

1. $a \in X$,
2. $\lim_{x \rightarrow a} f(x)$ exists, and
3. $f(a) = \lim_{x \rightarrow a} f(x)$

Similarly, a vector function is continuous if it satisfies the above conditions.

Recall that for a function $f : \mathbb{R} \rightarrow \mathbb{R}$, the derivative at some value x can be generalized by:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Similarly, the derivative of a vector function at some value x can be generalized as:

$$\vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h}$$

Thus, if $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$, we have:

$$\vec{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle$$

Theorem 3.1.1 ▶ Properties of the differentials of vector functions

Let $\vec{u}(t) = \langle f_1(t), g_1(t), h_1(t) \rangle$ and $\vec{v}(t) = \langle f_2(t), g_2(t), h_2(t) \rangle$ where $f_1, g_1, h_1, f_2, g_2, h_2$ are differentiable. Let c be a scalar. Then:

- $\frac{d}{dt} \vec{u}(t) = \langle f_1'(t), g_1'(t), h_1'(t) \rangle$
- $\frac{d}{dt} (\vec{u}(t) + \vec{v}(t)) = \frac{d}{dt} \vec{u}(t) + \frac{d}{dt} \vec{v}(t)$
- $\frac{d}{dt} c\vec{v}(t) = c \frac{d}{dt} \vec{v}(t)$
- $\frac{d}{dt} f(t)\vec{v}(t) = f'(t)\vec{v}(t) + f(t)\vec{v}'(t)$
- $\frac{d}{dt} \vec{u}(t) \cdot \vec{v}(t) = \vec{u}'(t) \cdot \vec{v}(t) + \vec{u}(t) \cdot \vec{v}'(t)$
- $\frac{d}{dt} \vec{u}(t) \times \vec{v}(t) = \vec{u}'(t) \times \vec{v}(t) + \vec{u}(t) \times \vec{v}'(t)$ (note that order here is sensitive)
- $\frac{d}{dt} \vec{u}(f(t)) = \vec{u}'(f(t))f'(t)$

If we consider the curve C formed by all the terminal points of the graph of a continuous vector function $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$, we call $\vec{r}(t)$ a **parameterization** of the curve C , and we call $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$ the **displacement vector**.

Example 3.1.1

Find the parametric equations of the tangent line to the helix with parameterization $\vec{r}(t) := \langle 2 \cos t, \sin t, t \rangle$ at the point $(0, 1, \pi/2)$

$$\vec{r}'\left(\frac{\pi}{2}\right) = \left\langle -2 \sin \frac{\pi}{2}, \cos \frac{\pi}{2}, 1 \right\rangle$$

Then the parameterized line is:

$$\left\langle 0, 1, \frac{\pi}{2} \right\rangle + t \langle -2, 0, 1 \rangle$$

Index

Definitions

2.1.1 Point	3
2.1.2 Distance	3
2.1.3 Sphere	3
2.2.1 Vector	4
2.2.2 Scalar Multiplication	4
2.2.3 Norm	5
2.2.4 Unit Vector	5
2.2.5 Dot Product	5
2.2.6 Parallel, Perpendicular	6
2.2.7 Orthogonal	6
2.2.8 Work	6
2.3.1 Gradient	7
2.3.2 Directional Derivative	8
2.4.1 Scalar Projection	9
2.4.2 Vector Projection	9
2.5.1 Cross Product	9
2.6.1 Torque	10
2.7.1 Planar Curve	11
2.7.2 Cylinder	11
2.7.3 Cone	11
2.7.4 Conic Surface	11
2.7.5 Quadratic Surface	11
2.7.6 Ellipsoid	12
3.0.1 Vector Function	13
3.1.1 Continuity	14

Examples

2.1.1 Distance Between Points	3
2.1.2 Circle	4

2.2.1 Finding Work	6
2.3.1 Gradient	7
2.3.2 Directional Derivative	8
2.7.1	11
3.0.1 Circle Vector Function	13
3.1.1	15

Techniques

2.2.1 Finding a Unit Vector from a Given Vector	5
2.5.1 Calculating Cross Product	9

Theorems

2.2.1 Angle Between Vectors	6
2.5.1	10
3.1.1 Properties of the differentials of vector functions	14