

# **Introduction to Analysis**

UT Knoxville, Spring 2023, MATH 341

Mike Frazier, Peter Humphries, Alex Zhang

April 5, 2023

# Contents

<b>Preface</b>	<b>2</b>
<b>1 Introduction</b>	<b>4</b>
<b>2 Logic and Proofs</b>	<b>6</b>
2.1 Basic Logic . . . . .	6
2.2 Proofs and Proof Techniques . . . . .	10
<b>3 Naive Set Theory</b>	<b>13</b>
3.1 Sets . . . . .	13
3.2 Functions . . . . .	16
3.3 Injectivity and Surjectivity . . . . .	19
<b>4 Number Systems</b>	<b>22</b>
4.1 Natural Numbers $\mathbb{N}$ and Induction . . . . .	22
4.2 Integers $\mathbb{Z}$ . . . . .	24
4.3 Rational Numbers $\mathbb{Q}$ . . . . .	25
4.4 Fields . . . . .	27
4.5 Ordered Fields . . . . .	29
4.6 Completeness . . . . .	30
<b>5 Suprema and Infima</b>	<b>34</b>
<b>6 Consequences of Completeness</b>	<b>37</b>
<b>7 Cardinality</b>	<b>41</b>
7.1 Additional Remarks . . . . .	47
<b>8 Sequences and Convergence</b>	<b>49</b>
8.1 Properties of Limits . . . . .	53
8.2 Subsequences . . . . .	57
8.3 Limit Superior and Inferior . . . . .	59
8.4 Cauchy Sequences . . . . .	62

<b>9</b>	<b>Open and Closed Sets</b>	<b>65</b>
9.1	Open Sets . . . . .	65
9.2	Closed Sets . . . . .	67
9.3	Closure . . . . .	69
<b>10</b>	<b>Compact Sets</b>	<b>72</b>
10.1	Sequential Compactness . . . . .	72
10.2	Open Cover Compactness . . . . .	74
	<b>Index</b>	<b>79</b>

# Preface

These are my notes for the **Introduction to Analysis** course at the University of Tennessee (MATH 341). It is compiled from several sources including lecture notes by Dr. Michael Frazier and Dr. Peter Humphries, as well as online resources such as the Mathematics Stack Exchange.

The first few weeks of the course are spent reviewing the material taught in **Introduction to Abstract Mathematics** (MATH 300): logic, set theory, number systems, and cardinality. They serve as a “primer” for the following material on real analysis.

# Introduction

Real analysis is the branch of mathematics concerned with the study of functions, limits, and continuity of real numbers. It encompasses a wide range of concepts and techniques, including differentiation and integration. Real analysis serves as a foundation for many areas of mathematics and science. Its principles are used extensively in various computational techniques and algorithms.

In this course, we will focus on the central idea of convergence, which is fundamental to many of these concepts and techniques. To motivate our exploration of convergence, we will examine classic iterative methods that rely on the concept of convergence to approximate solutions.

## Example 1.0.1 ► Newton's Method

Given  $c > 0$ , suppose we want to calculate  $\sqrt{c}$ . Start with some initial guess  $x_1 > 0$ .

$$\begin{aligned}\text{Let } x_2 &:= \frac{1}{2} \left( x_1 + \frac{c}{x_1} \right) \\ \text{Let } x_3 &:= \frac{1}{2} \left( x_2 + \frac{c}{x_2} \right) \\ &\vdots \\ \text{Let } x_{n+1} &:= \frac{1}{2} \left( x_n + \frac{c}{x_n} \right)\end{aligned}$$

We find that  $\lim_{n \rightarrow \infty} x_n = \sqrt{c}$ .

Does this method work for all  $c > 0$  and  $x_1 > 0$ ? Assuming  $\lim_{n \rightarrow \infty} x_n = x$  converges, then:

$$\begin{aligned}
 x_{n+1} &= \frac{1}{2} \left( x_n + \frac{c}{x_n} \right) \\
 \implies x &= \frac{1}{2} \left( x + \frac{c}{x} \right) \\
 \implies 2x &= x + \frac{c}{x} \\
 \implies x &= \frac{c}{x} \\
 \implies x^2 &= c \\
 \implies x &= \sqrt{c}
 \end{aligned}$$

The above calculation only makes sense if we know the sequence converges. Consider the sequence  $x_{n+1} = 6 - x_n$  where  $x_1 = 4$ . Then:

$$x_1 = 4, \quad x_2 = 2, \quad x_3 = 4, \quad x_4 = 2, \quad \dots$$

Since this sequence does not converge, there is no limit when  $n$  approaches infinity.

Let's look at a more complicated iterative method.

### Example 1.0.2 ► Picard's Method

Suppose we had to solve  $y' = f(x, y)$  where  $y(x_0) = y_0$  (i.e. find a function  $y$  that satisfies our two conditions). As it turns out, we can use an iterated method to solve this as well.

- Start with an initial guess  $y_1(x)$
- Define  $y_{n+1}(x) := y_0 + \int_{x_0}^x f(t, y_n) dt$ .

Provided that  $f$  and  $y_0$  are “well-behaving”, then the sequence of functions  $y_n(x)$  converges to the solution  $y(x)$ .

This idea that an infinite sequence of functions can converge suggests some notion of “distance” between functions. We can use a number of metrics for distance, some possibilities including:

- $\int_a^b |f(x) - g(x)| dx$  (total area between the two functions)
- $\sup \{x : x = |f(x) - g(x)|\}$  (max possible “vertical” distance between the two curves)

# Logic and Proofs

Formal logic is the foundation of mathematics. It enables us to construct logically consistent models by starting with a set of axioms and systematically deducing new statements from them. This process not only helps us to prove known results but also to uncover new mathematical concepts and theorems.

## 2.1 Basic Logic

### Definition 2.1.1 ► Statement

A **statement** is a claim that is either true or false.

$$p : \text{some claim}$$

We usually denote statements with a letter like  $p$ . For example, we can write “ $p : x > 2$ ”, which means  $p$  represents the statement “ $x$  is greater than 2”. Throughout this chapter, we will use  $p$  and  $q$  to represent arbitrary statements.

### Definition 2.1.2 ► Conjunction

Logical **conjunction** is an operation that takes two statements and produces a new statement that is true only when both input statements are true.

$$p \wedge q : p \text{ is true and } q \text{ is true}$$

### Definition 2.1.3 ► Disjunction

Logical **disjunction** is an operation that takes two statements and produces a new statement that is true when at least one of the input statements is true.

$$p \vee q : p \text{ is true or } q \text{ is true}$$

Conjunction and Disjunction follow our intuition of “and” and inclusive “or”, respectively. We can visualize the two logical connectives using *truth tables*.

#### Example 2.1.4 ▶ Truth Table of Conjunction

$p$	$q$	$p \implies q$
T	T	T
T	F	F
F	T	F
F	F	F

#### Example 2.1.5 ▶ Truth Table of Disjunction

$p$	$q$	$p \implies q$
T	T	T
T	F	T
F	T	T
F	F	F

#### Definition 2.1.6 ▶ Negation

The *negation* of a statement is a statement with opposite truth values.

$$\neg p$$

#### Definition 2.1.7 ▶ Implication

An *implication* “ $p$  implies  $q$ ” states “if  $p$  is true, then  $q$  is true”.

$$p \implies q$$

In the implication  $p \implies q$ , we call  $p$  the *hypothesis* and  $q$  the *conclusion*. If the hypothesis is false to begin with, then the implication is not really meaningful. Instead of assigning those kinds of implications no truth value, we simply consider them true by convention. These kinds of truths are called *vacuous truths*.



**Example 2.1.8 ▶ Truth Table of Implication**

$p$	$q$	$p \implies q$
T	T	T
T	F	F
F	T	T
F	F	T

**Example 2.1.9 ▶ Simple Statements**

Let  $p : x > 2$  and  $q : x^2 > 1$ . Consider the following statements:

- “For all real numbers  $x$ ,  $p \implies q$ ”

**True.** If  $x > 2$ , then  $x^2 > 1$ .<sup>a</sup>

- “For all real numbers  $x$ ,  $q \implies p$ ”

**False.** Consider  $x = 1.1$ . Then  $x^2 = 1.21 > 1$ , but  $x = 1.1 < 2$ .

<sup>a</sup>This is normally where we would rigorously prove such a statement, but we will omit this for now.

**Definition 2.1.10 ▶ Logical Equivalence**

$p$  and  $q$  are **logically equivalent** if  $p \implies q$  and  $q \implies p$ .

$$p \iff q$$

In other words,  $p \iff q$  means that  $p$  and  $q$  share the same truth value. Either  $p$  and  $q$  are **always both true**, or  $p$  and  $q$  are **always both false**. Logical equivalence says nothing about the truth of  $p$  and  $q$  themselves.

We can also say “ $p$  if and only if  $q$ ” or “ $p$  iff  $q$ ” to denote logical equivalence.

**Example 2.1.11 ▶ Truth Table of Logical Equivalence**

$p$	$q$	$p \iff q$
T	T	T
T	F	F
F	T	F
F	F	T

**Definition 2.1.12 ▶ Converse**

Given the implication  $p \implies q$ , its **converse** statement is  $q \implies p$ .

It's important to note that an implication and its converse have no intrinsic equivalence.

**Example 2.1.13 ▶ Truth Table of Converse**

$p$	$q$	$p \implies q$	$q \implies p$
T	T	T	T
T	F	F	T
F	T	T	F
F	F	T	T

**Definition 2.1.14 ▶ Contrapositive**

Given the implication  $p \implies q$ , its **contrapositive** statement is  $\neg q \implies \neg p$ .

Unlike the converse, an implication and its contrapositive are logically equivalent. To help our intuition, we can construct a truth table.

**Example 2.1.15 ▶ Truth Table of Contrapositive**

$p$	$q$	$\neg p$	$\neg q$	$p \implies q$	$\neg q \implies \neg p$
T	T	F	F	T	T
T	F	F	T	F	F
F	T	T	F	T	T
F	F	T	T	T	T

As we can see, no matter what the truth values of the hypothesis and conclusion are, an implication and its contrapositive always have the same truth values.

When constructing a truth table, we must include **all** intermediate statements, not just the final statement.

## 2.2 Proofs and Proof Techniques

While truth tables are a useful tool for evaluating simple statements, they quickly become impractical when dealing with more complex propositions. Moreover, they do not offer insights into the reasoning behind such statements. In contrast, proofs can provide us with a deeper understanding of logical relationships and help us reason about complex statements.

In particular, we often need to prove implications of the form  $p \implies q$ , where the truth of  $p$  guarantees the truth of  $q$ . To do so, we can use a variety of proof techniques:

1. **Direct Proof:** Assume  $p$  is true, then reason that  $q$  must be true as well.
2. **Proof by Contradiction:** Assume both  $p$  and  $\neg q$  are true, then logically derive some contradiction.
3. **Proof by Contrapositive:** Assume  $\neg q$  is true, then reason that  $\neg p$  must be true as well.

In mathematical proofs, there are two main types of reasoning: direct and indirect. A direct proof shows a clear path from the premises to the conclusion, providing valuable insights into the underlying mathematics. In contrast, indirect proofs rely on a contradictory hypothesis to establish the truth of the conclusion. While indirect proofs can be useful when a direct proof is not readily available, they may be less insightful since they do not provide much context surrounding the premises.

However, it is worth noting that an indirect proof may be easier to find than a direct proof in certain cases. While a direct proof requires identifying the correct path that leads to the conclusion, an indirect proof only needs to deduce any contradictory statement. Despite this advantage, indirect proofs should be used sparingly and only when a direct proof is not feasible.

### Technique 2.2.1 ► Proof by Contradiction

To prove  $p \implies q$  by contradiction, we carry out the following steps:

1. Assume  $p$  is true, and suppose for the sake of contradiction  $\neg q$  is true.
2. Logically derive a statement that contradicts something we know to be true.
3. Ultimately conclude that  $q$  must be true.

In terms of logic notation, proof by contradiction follows:

$$[(p \wedge (\neg q)) \implies \text{Contradiction}] \implies [p \implies q]$$

### Example 2.2.2 ► Truth Table of Proof by Contradiction

$p$	$q$	$p \implies q$	$\neg q$	$p \wedge (\neg q)$	$\neg [p \wedge (\neg q)]$
T	T	T	F	F	T
T	F	F	T	T	F
F	T	T	F	F	T
F	F	T	T	F	T

By the above truth table, we can safely assume the following logical equivalence:

$$(p \implies q) \iff \neg [p \wedge (\neg q)]$$

### Technique 2.2.3 ► Proof by Contrapositive

To prove  $p \implies q$  by contrapositive, we carry out the following steps:

1. Assume  $\neg q$  is true.
2. Directly prove that  $\neg p$  is true.

In terms of logic notation, proof by contrapositive follows:

$$(\neg q \implies \neg p) \iff (p \implies q)$$

We can actually prove this using proof by contradiction!

#### Example 2.2.4 ► Logical Equivalence of Contrapositive

Given statements  $p$  and  $q$ ,  $p \implies q$  and  $\neg q \implies \neg p$  are equivalent.

*Proof.* Assume  $p \implies q$ . To prove  $\neg q \implies \neg p$ , we can suppose for contradiction that  $\neg q$  and  $p$  are both true. But  $p \implies q$ , so  $q$  is true which contradicts  $\neg q$ . Hence, the assumption that  $p$  is true was incorrect. Thus,  $\neg q \implies \neg p$ .

Assume  $\neg q \implies \neg p$ . From above, we have  $\neg(\neg p) \implies \neg(\neg q)$ , so  $p \implies q$ . □

#### Example 2.2.5 ► Proving Simple Logic Statements

Let  $p$ ,  $q$ , and  $r$  be arbitrary statements. Prove that  $[p \implies (q \vee r)] \iff [(p \wedge \neg q) \implies r]$ .

*Proof.* Assume  $p \implies (q \vee r)$ . Suppose  $p \wedge \neg q$ . Then  $p$  is true, so  $q \vee r$  is true by assumption. Also,  $\neg q$  is true, so  $r$  must be true from  $q \vee r$ .

Assume  $(p \wedge \neg q) \implies r$ . Suppose  $p$  is true. There are two possibilities:

1. If  $q$  is true, then  $q \vee r$  is true.
2. If  $\neg q$  is true, then  $p \wedge \neg q$  is true. Thus,  $r$  is true by assumption. Hence,  $q \vee r$  is true.

□

# Naive Set Theory

Instead of forming a rigorous, axiomatic basis for sets, we will simply take an informal approach to sets guided by our intuition. Ultimately, our introduction to real analysis does not fiddle with the fine details of set theory, so it's safe to take a naive approach.

## 3.1 Sets

### Definition 3.1.1 ► Set

A **set** is a collection of distinct objects.

For example,  $\mathbb{N} := \{1, 2, 3 \dots\}$  is the set of all natural numbers, and  $\mathbb{Z} := \{\dots, 1, 2, 3, \dots\}$  is the set of all integers. It's conventional to use capital letters to denote sets and use lowercase letters to denote elements of sets. Throughout this chapter, we will use  $A$  and  $B$  to represent arbitrary sets.

### Definition 3.1.2 ► Membership, $\in$

We write  $a \in A$  to mean “ $a$  is in  $A$ ”.

### Definition 3.1.3 ► Subset, $\subseteq$

$A$  is a **subset** of  $B$  if everything in  $A$  is also in  $B$ .

$$A \subseteq B \iff \forall (x \in A)(x \in B)$$

### Definition 3.1.4 ► Set Equality, $=$

$A$  **equals**  $B$  if  $A$  is a subset of  $B$  and  $B$  is a subset of  $A$ .

$$A = B \iff (A \subseteq B \wedge B \subseteq A)$$

**Definition 3.1.5 ▶ Proper Subset,  $\subsetneq$** 

$A$  is a **proper subset** of  $B$  if  $A$  is a subset of  $B$  but  $B$  is not a subset of  $A$ .

$$A \subsetneq B \iff (A \subseteq B \wedge B \not\subseteq A)$$

In other words,  $A$  is a proper subset of  $B$  if everything in  $A$  is also in  $B$ , but  $B$  has something that  $A$  does not.

Among mathematical texts, the generic subset symbol  $\subset$  has no standardized definition. Some use it to represent subset or equal; others use it to represent proper subset. We will simply not use  $\subset$  to avoid any ambiguity.

**Definition 3.1.6 ▶ Empty Set ( $\emptyset$ )**

The **empty set** is the set that contains no elements.

$$\emptyset := \{\}$$

As convention, we assume that  $\emptyset$  is a subset of every set, including itself.

**Technique 3.1.7 ▶ Proving a Subset Relation**

To prove that  $A \subseteq B$ :

1. Let  $x$  be an arbitrary element of  $A$ .
2. Show that  $x \in B$ .

To prove that  $A \not\subseteq B$ , choose a specific  $x \in A$  and show  $x \notin B$ .

**Example 3.1.8 ▶ Proving Simple Subset Relation**

Suppose that  $A \subseteq B$  and  $B \subseteq C$ . Prove that  $A \subseteq C$ .

*Proof.* Let  $x \in A$  be arbitrary. Since  $A \subseteq B$ , then  $x \in B$ . Similarly, since  $B \subseteq C$ , then  $x \in C$ . Therefore,  $A \subseteq C$ . □

**Definition 3.1.9 ▶ Union**

The **union** of two sets is the set of all things that are in one or the other set.

$$A \cup B := \{x : x \in A \vee x \in B\}$$

### Definition 3.1.10 ► Intersection

The **intersection** of two sets is the set of all things that are in both sets.

$$A \cap B := \{x : x \in A \wedge x \in B\}$$

More generally, we can apply union and intersection to an arbitrary number of sets, finite or infinite. We use a notation similar to summation using  $\sum$ . Let  $\Lambda$  be an indexing set, and for each  $\lambda \in \Lambda$ , let  $A_\lambda$  be a set.

$$\bigcup_{\lambda \in \Lambda} A_\lambda = \{x : x \in A_\lambda \text{ for some } \lambda \in \Lambda\}$$

$$\bigcap_{\lambda \in \Lambda} A_\lambda = \{x : x \in A_\lambda \text{ for all } \lambda \in \Lambda\}$$

### Example 3.1.11 ► Indexed Sets

For  $n \in \mathbb{N}$ , let  $A_n = \left[\frac{1}{n}, 1\right] = \{x \in \mathbb{R} : \frac{1}{n} \leq x \leq 1\}$ . Prove that:

- (a)  $\bigcup_{n=1}^{\infty} A_n = (0, 1]$
- (b)  $\bigcap_{n=1}^{\infty} A_n = \{1\}$

*Proof of (a).* Suppose  $x \in \bigcup_{n=1}^{\infty} A_n$ . Then there exists  $n \in \mathbb{N}$  such that  $x \in A_n = \left[\frac{1}{n}, 1\right]$ . That is,  $0 < \frac{1}{n} \leq x \leq 1$ . Therefore,  $x \in (0, 1]$ .

Suppose  $x \in (0, 1]$ . Then  $x > 0$ , so there exists  $n_0 \in \mathbb{N}$  such that  $\frac{1}{n_0} < x$ . Then  $\frac{1}{n_0} \leq x \leq 1$ , so  $x \in A_{n_0}$ . Therefore,  $x \in \bigcup_{n=1}^{\infty} A_n$ .  $\square$

*Proof of (b).* Suppose  $x \in \bigcap_{n=1}^{\infty} A_n$ . Then  $x \in A_1 = \{1\}$ .

Suppose  $x \in \{1\}$ . Then  $x = 1 \in \left[\frac{1}{n}, 1\right]$  for all  $n \in \mathbb{N}$ . Therefore,  $x \in \bigcap_{n=1}^{\infty} A_n$ .  $\square$

### Definition 3.1.12 ► Set Minus

The **set difference** of two sets is the set of all things that are in first set but not the second set.



$$A \setminus B = \{x \in A : x \notin B\}$$

### Definition 3.1.13 ► Complement

Let  $X$  be a set called the **universal set**. The **complement** of  $A$  in  $X$  is defined as  $X \setminus A$ .

$$A^c = X \setminus A = \{x \in X : x \notin A\}$$

### Theorem 3.1.14 ► De Morgan's Laws for Sets

Suppose  $X$  is a set, and for any subset  $S$  of  $X$ , let  $S^c = X \setminus S$ . Suppose that  $A_\lambda \subseteq X$  for every  $\lambda$  belonging to some index set  $\Lambda$ . Prove that:

- (a)  $\left(\bigcup_{\lambda \in \Lambda} A_\lambda\right)^c = \bigcap_{\lambda \in \Lambda} A_\lambda^c$ ;
- (b)  $\left(\bigcap_{\lambda \in \Lambda} A_\lambda\right)^c = \bigcup_{\lambda \in \Lambda} A_\lambda^c$ .

*Proof of (a).* First, let  $a \in \left(\bigcup_{\lambda \in \Lambda} A_\lambda\right)^c$ . Then,  $a \in X \setminus \left(\bigcup_{\lambda \in \Lambda} A_\lambda\right)$ , so  $a \in X$  but  $a \notin \left(\bigcup_{\lambda \in \Lambda} A_\lambda\right)$ . Thus,  $a \notin A_\lambda$  for any  $\lambda \in \Lambda$ , so  $a \in X \setminus A_\lambda$  for all  $\lambda \in \Lambda$ . In other words,  $a \in \bigcap_{\lambda \in \Lambda} A_\lambda^c$ .

Next, let  $a \in \bigcap_{\lambda \in \Lambda} A_\lambda^c$ . Then  $a \in A_\lambda^c$  for all  $\lambda \in \Lambda$ , so  $a \in X$  but  $a \notin A_\lambda$  for all  $\lambda \in \Lambda$ . That is,  $a \notin \left(\bigcup_{\lambda \in \Lambda} A_\lambda\right)$ . In other words,  $a \in \left(\bigcup_{\lambda \in \Lambda} A_\lambda\right)^c$ .  $\square$

*Proof of (b).* First, let  $a \in \left(\bigcap_{\lambda \in \Lambda} A_\lambda\right)^c$ . Then,  $a \in X \setminus \bigcap_{\lambda \in \Lambda} A_\lambda$ , so  $a \in X$  but  $a \notin \bigcap_{\lambda \in \Lambda} A_\lambda$ . That is,  $a \notin A_\lambda$  for some  $\lambda \in \Lambda$ . Thus,  $a \in X \setminus A_\lambda$  for some  $\lambda \in \Lambda$ . Therefore,  $a \in \bigcup_{\lambda \in \Lambda} A_\lambda^c$ .

Next, let  $a \in \bigcup_{\lambda \in \Lambda} A_\lambda^c$ . Then  $a \in A_\lambda^c$  for some  $\lambda \in \Lambda$ , so  $a \in X$  but  $a \notin A_\lambda$  for some  $\lambda \in \Lambda$ . That is,  $a \notin \left(\bigcap_{\lambda \in \Lambda} A_\lambda\right)$ . Therefore,  $a \in \left(\bigcap_{\lambda \in \Lambda} A_\lambda\right)^c$ .  $\square$

## 3.2 Functions

We generally think of functions as a “map” or “rule” that assigns numbers to other numbers. For example,  $f(x) = 2x$  maps  $1 \mapsto 2$ ,  $2 \mapsto 4$ , etc. More formally, we define functions in terms of sets.

**Definition 3.2.1 ▶ Cartesian Product**

Let  $X$  and  $Y$  be sets. The **Cartesian product** of  $X$  and  $Y$  is the set of all ordered pairs  $(x, y)$  where  $x \in X$  and  $y \in Y$ .

$$X \times Y := \{(x, y) : x \in X \wedge y \in Y\}$$

**Definition 3.2.2 ▶ Relation**

Let  $X$  and  $Y$  be sets. A **relation** between  $X$  and  $Y$  is a subset of the Cartesian product  $X \times Y$ .

**Definition 3.2.3 ▶ Function**

Let  $X$  and  $Y$  be sets. A **function** from  $X$  to  $Y$  is a relation from  $X$  to  $Y$  such that for every  $x \in X$ , there exists a unique  $y \in Y$  where  $(x, y) \in f$ .

More formally, a **function**  $f : X \rightarrow Y$  is a subset of  $X \times Y$  satisfying:

1.  $\forall(x \in X) [\exists(y \in Y)((x, y) \in f)]$
2.  $(x, y_1), (x, y_2) \in f \implies y_1 = y_2$

Given  $f : X \rightarrow Y$ , we call  $X$  the **domain** of  $f$  and  $Y$  the **codomain** of  $f$ . Given  $x \in X$ , we write  $f(x)$  to denote the unique element of  $Y$  such that  $(x, y) \in f$ .

$$f(x) = y \iff (x, y) \in f$$

**Definition 3.2.4 ▶ Function Image**

Let  $f : X \rightarrow Y$  be a function and  $A \subseteq X$ . The **image** of  $A$  under  $f$  is the set containing all possible function outputs from all inputs in  $A$ .

$$f[A] := \{f(a) : a \in A\}$$

Given  $f : X \rightarrow Y$ , we call  $f[X]$  the **range** of  $f$ .

**Example 3.2.5 ▶ Function Images**

Suppose  $f : X \rightarrow Y$  is a function, and  $A_\lambda \subseteq X$  for each  $\lambda \in \Lambda$ . Then:

$$(a) \ f\left[\bigcup_{\lambda \in \Lambda} A_\lambda\right] = \bigcup_{\lambda \in \Lambda} f[A_\lambda]$$

$$(b) f\left[\bigcap_{\lambda \in \Lambda} A_\lambda\right] \subseteq \bigcap_{\lambda \in \Lambda} f[A_\lambda]$$

In this example, we will only prove the “forward” direction. That is, we want to show that  $f\left[\bigcup_{\lambda \in \Lambda} A_\lambda\right] \subseteq \bigcup_{\lambda \in \Lambda} f[A_\lambda]$ .

*Proof of (a).* Let  $y \in f\left[\bigcup_{\lambda \in \Lambda} A_\lambda\right]$ . By definition of Function Image, there exists  $x \in \bigcup_{\lambda \in \Lambda} A_\lambda$  such that  $y = f(x)$ . Thus, there exists  $\lambda_0 \in \Lambda$  such that  $x \in \lambda_0$ . That is,  $y \in f[A_{\lambda_0}]$ . Therefore,  $y \in \bigcup_{\lambda \in \Lambda} f[A_\lambda]$ .  $\square$

### Definition 3.2.6 ▶ Function Inverse Image

Let  $f : X \rightarrow Y$  be a function and  $B \subseteq Y$ . The **inverse image** of  $B$  under  $f$  is the set containing all possible function inputs whose output is in  $B$ .

$$f^{-1}[B] := \{x \in X : f(x) \in B\}$$

Note the following logical equivalence:

$$x \in f^{-1}[B] \iff f(x) \in B$$

### Example 3.2.7 ▶ Function Inverse Images

Suppose  $f : X \rightarrow Y$  is a function, and  $B_\lambda \subseteq Y$  for each  $\lambda \in \Lambda$ . Then:

$$f^{-1}\left[\bigcup_{\lambda \in \Lambda} B_\lambda\right] = \bigcup_{\lambda \in \Lambda} f^{-1}[B_\lambda]$$

Again, we will only prove the “forward direction”.

*Proof.* Let  $x \in f^{-1}\left[\bigcup_{\lambda \in \Lambda} B_\lambda\right]$ . Then,  $f(x) \in \bigcup_{\lambda \in \Lambda} B_\lambda$ . That is,  $f(x) \in B_{\lambda_0}$  for some  $\lambda_0 \in \Lambda$ . Thus,  $x \in f^{-1}[B_{\lambda_0}]$ , so  $x \in \bigcup_{\lambda \in \Lambda} f^{-1}[B_\lambda]$ .  $\square$

### 3.3 Injectivity and Surjectivity

#### Definition 3.3.1 ► Injective, One-to-one

A function  $f : X \rightarrow Y$  is **injective** or **one-to-one** if no two inputs in  $X$  have the same output in  $Y$ .

$$\forall (x_1, x_2 \in X) [x_1 \neq x_2 \implies f(x_1) \neq f(x_2)]$$

We can also think of injectivity as, “if two inputs have the same output, then the two inputs must be the same”. It’s really just the contrapositive of our initial definition, which we know must be logically equivalent.

$$\forall (x_1, x_2 \in X) [f(x_1) = f(x_2) \implies x_1 = x_2]$$

For example, the function  $f(x) = x^2$  is not injective, because  $f(-1) = 1$  and  $f(1) = 1$ . We have two distinct inputs that map to the same output.

#### Technique 3.3.2 ► Proving a Function is Injective

To prove a function  $f : X \rightarrow Y$  is injective:

1. Let  $x_1, x_2 \in X$  where  $f(x_1) = f(x_2)$ .
2. Reason that  $x_1 = x_2$ .

#### Example 3.3.3 ► Proving Injectivity

$f(x) = -3x - 7$  is injective.

*Proof.* Suppose  $f(x_1) = f(x_2)$ . Then  $-3x_1 - 7 = -3x_2 - 7$ , so  $-3x_1 = -3x_2$ . Thus,  $x_1 = x_2$ , so  $f$  is injective. □

#### Example 3.3.4 ► Disproving Injectivity

Prove that  $f(x) = x^2$  is not injective.

*Proof.*  $f(-1) = 1$  and  $f(1) = 1$ , but  $-1 \neq 1$ . Thus,  $f$  is not injective. □

**Definition 3.3.5 ▶ Surjective, Onto**

A function  $f : X \rightarrow Y$  is **surjective** or **onto** if everything in  $Y$  has a corresponding input in  $X$ .

$$\forall(y \in Y)[\exists(x \in X)(f(x) = y)]$$

Note that  $f : X \rightarrow f[X]$  is **always** surjective.

**Technique 3.3.6 ▶ Proving a Function is Surjective**

To prove a function  $f : X \rightarrow Y$  is surjective:

1. Let  $y \in Y$  be arbitrary.
2. “Undo” the function  $f$  to obtain  $x \in X$  where  $f(x) = y$ .

**Example 3.3.7 ▶ Proving Surjectivity**

Prove that  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = -3x + 7$  is surjective.

*Proof.* Let  $y \in Y$  be arbitrary. Let  $x := \frac{y-7}{-3}$ . Then  $x \in \mathbb{R}$ , and:

$$\begin{aligned} f(x) &= -3\left(\frac{y-7}{-3}\right) + 7 \\ &= (y-7) + 7 \\ &= y \end{aligned}$$

Therefore,  $f$  is surjective. □

**Definition 3.3.8 ▶ Bijective**

A function  $f : X \rightarrow Y$  is **bijective** if it is both injective and surjective.

**Definition 3.3.9 ▶ Function Composition**

Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be functions. The **composition** of  $f$  and  $g$  is a function  $g \circ f : X \rightarrow Z$  defined by:

$$(g \circ f)(x) := g(f(x))$$

**Theorem 3.3.10 ▶ Composition Preserves Injectivity and Surjectivity**

Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are functions.

- (a) If  $f$  and  $g$  are injective, then  $g \circ f$  is injective.
- (b) If  $f$  and  $g$  are surjective, then  $g \circ f$  is surjective.
- (c) If  $f$  and  $g$  are bijective, then  $g \circ f$  is bijective.

*Proof of (a).* Let  $x_1, x_2 \in X$ . Suppose that  $(g \circ f)(x_1) = (g \circ f)(x_2)$ . Then,  $g(f(x_1)) = g(f(x_2))$ . Because  $g$  is injective, we have  $f(x_1) = f(x_2)$ . Because  $f$  is injective, we have  $x_1 = x_2$ . Therefore,  $g \circ f$  is injective.  $\square$

*Proof of (b).* Let  $z \in Z$ . Because  $g$  is surjective, there exists an element  $y \in Y$  such that  $g(y) = z$ . Because  $f$  is surjective, there exists an element  $x \in X$  such that  $f(x) = y$ . Thus,  $(g \circ f)(x) = g(f(x)) = g(y) = z$ . Therefore,  $g \circ f$  is surjective.  $\square$

*Proof of (c).* We know that from (a) and (b) composition preserves injectivity and surjectivity. Thus, composition must also preserve bijectivity.  $\square$

**Definition 3.3.11 ▶ Inverse Function**

Let  $f : X \rightarrow Y$  be a bijection. The **inverse function** of  $f$  is a function  $f^{-1} : Y \rightarrow X$  defined by:

$$f^{-1} := \{(y, x) \in Y \times X : (x, y) \in f\}$$

The notation for inverse functions conflicts with the notation for inverse images. A key distinction to make it that only bijections can have an inverse function, but we can apply the inverse image to any function. Thus, given a bijection  $f : X \rightarrow Y$ , we know  $f^{-1}(f(x)) = x$  for all  $x \in X$ , and  $f(f^{-1}(y)) = y$  for all  $y \in Y$ .

**Example 3.3.12**

Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  be functions such that  $(g \circ f)(x) = x$  for all  $x \in X$ , and  $(f \circ g)(y) = y$  for all  $y \in Y$ .  $f^{-1} = g$ .

*Proof.* todo: finish proof  $\square$

# Number Systems

Our goal is to create an axiomatic basis for the real numbers  $\mathbb{R}$ . We need to establish axioms for  $\mathbb{R}$  and then derive all further properties from the axioms. We would like these axioms to be as minimal and agreeable as possible; however, finding axioms that characterize  $\mathbb{R}$  is not easy. Instead, we'll start from the natural numbers  $\mathbb{N}$  and expand from there.

## 4.1 Natural Numbers $\mathbb{N}$ and Induction

How do we define the natural numbers? Listing every natural number is definitely not an option. We could try to define the natural numbers as  $\mathbb{N} := \{1, 2, \dots\}$ . However, the “...” is ambiguous. Instead, we can define  $\mathbb{N}$  in terms of its properties.

### Definition 4.1.1 ► Peano Axioms for $\mathbb{N}$

The **Peano axioms** are five axioms that can be used to define the natural numbers  $\mathbb{N}$ .

1.  $1 \in \mathbb{N}$
2. Every  $n \in \mathbb{N}$  has a successor called  $n + 1$ .
3. 1 is **not** the successor of any  $n \in \mathbb{N}$ .
4. If  $n, m \in \mathbb{N}$  have the same successor, then  $n = m$ .
5. If  $1 \in S$  and every  $n \in S$  has a successor, then  $\mathbb{N} \subseteq S$ .

Note that there is not one “prescribed” way to do define the natural numbers. This is just the most popular approach.

From the fifth Peano axiom, we can derive a new proof technique for proving statements about consecutive natural numbers.

### Theorem 4.1.2 ► Principle of Induction (by the Peano Axioms)

Let  $P(n)$  be a statement for each  $n \in \mathbb{N}$ . Suppose that:

1.  $P(1)$  is true, and
2. if  $P(n)$  is true, then  $P(n + 1)$  is true.

Then  $P(n)$  is true for all  $n \in \mathbb{N}$ .

*Proof.* Let  $S := \{n \in \mathbb{N} : P(n)\}$ . Then  $1 \in S$  because  $P(1)$  is true. Note that if  $n \in S$ , then  $P(n)$  is true. Hence,  $P(n+1)$  is true by assumption, so  $n+1 \in S$ . By the fifth Peano axiom, we have  $\mathbb{N} \subseteq S$ . Since  $S$  was defined as a subset of  $\mathbb{N}$ , we have  $\mathbb{N} = S$ . Therefore,  $P(n)$  is true for all  $n \in \mathbb{N}$ .  $\square$

A proof by induction has a “domino effect”. Imagine a domino for each natural number 1, 2, 3, and so on, arranged in an infinite row. Knocking the 1st domino will knock them all down.

$$\underbrace{P(1)}_{\text{by 1.}} \implies \underbrace{P(2)}_{\text{by 2.}} \implies \underbrace{P(3)}_{\text{by 2.}} \implies \dots$$

### Technique 4.1.3 ▶ Proof by Induction

To prove a statement  $P(n)$  for all  $n \in \mathbb{N}$ , we need to prove two statements:

1. **Base Case:** Prove  $P(1)$ .
2. **Induction Step:** Assume  $P(n)$  is true from some  $n \in \mathbb{N}$ , then prove  $P(n) \implies P(n+1)$ .

It is crucial that we actually use our assumption that  $P(n)$  is true in the induction step. Otherwise, our proof is most likely wrong.

### Example 4.1.4 ▶ Simple Proof by Induction

Prove that  $1 + 2 + \dots + n = \frac{n(n+1)}{2}$  for all  $n \in \mathbb{N}$ .

*Proof.* Let  $P(n)$  be the statement  $1 + 2 + \dots + n = \frac{n(n+1)}{2}$ .

**Base Case:** When  $n = 1$ , LHS = 1 and RHS =  $\frac{1(1+1)}{2} = 1$ , so  $P(1)$  is true.

**Induction Step:** Assume that  $P(n)$  is true for some  $n \in \mathbb{N}$ . Then:

$$\begin{aligned} 1 + 2 + \dots + n + (n+1) &= \frac{n(n+1)}{2} + (n+1) \\ &= (n+1) \left( \frac{n}{2} + 1 \right) \\ &= \frac{(n+1)(n+2)}{2} \end{aligned}$$

That is,  $P(n+1)$  is true. By the Principle of Induction (by the Peano Axioms),  $P(n)$  is



true for all  $n \in \mathbb{N}$ .



## 4.2 Integers $\mathbb{Z}$

From the natural numbers, we can easily construct the integers. First, we assume the existence of an operation, addition (+) and multiplication ( $\cdot$ ). On  $\mathbb{N}$ , we assume addition and multiplication satisfy the following properties for all  $a, b, c \in \mathbb{N}$ :

- **Commutativity**  $a + b = b + a$   $a \cdot b = b \cdot a$
- **Associativity**  $(a + b) + c = a + (b + c)$   $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- **Distributivity**  $a \cdot (b + c) = a \cdot b + a \cdot c$
- **Identity**  $1 \cdot n = n$

We can expand this number system by including:

1. an **additive identity** ( $n + 0 = n$  for all  $n \in \mathbb{N}$ )
2. **additive inverses** (for all  $n \in \mathbb{N}$ , add  $-n$  so  $-n + n = 0$ )

From this, we can construct the set of integers.

### Definition 4.2.1 ► Integers $\mathbb{Z}$

The set of **integers** is defined as:

$$\mathbb{Z} := \mathbb{N} \cup \{0\} \cup \{-n : n \in \mathbb{N}\}$$

### Definition 4.2.2 ► Even, Odd, Parity

Let  $a \in \mathbb{Z}$ .

- $a$  is **even** if there exists  $k \in \mathbb{Z}$  where  $a = 2k$ .
- $a$  is **odd** if there exists  $k \in \mathbb{Z}$  where  $a = 2k + 1$ .
- **Parity** describes whether an integer is even or odd.

### Theorem 4.2.3 ► Parity Exclusivity

Every integer is either even or odd, never both.

TODO: prove this

Sort  
out this  
wonky  
format-  
ting

**Example 4.2.4 ▶ Parity of Square**

For  $n \in \mathbb{Z}$ , if  $n^2$  is even, then  $n$  is even.

*Proof.* We proceed by contraposition. Suppose  $n$  is not even. Then  $n$  is odd, and thus can be expressed as  $n = 2k + 1$  for some  $k \in \mathbb{Z}$ . Then:

$$\begin{aligned} n^2 &= (2k + 1)(2k + 1) \\ &= 4k^2 + 4k + 1 \end{aligned}$$

Since the integers are closed under addition and multiplication, then  $4k^2 + 4k \in \mathbb{Z}$ . Thus,  $n^2$  is odd. □

**4.3 Rational Numbers  $\mathbb{Q}$** 

We can further expand this number system by the following:

1. Include **multiplicative inverses** (for all  $n \in \mathbb{Z} \setminus \{0\}$ , define  $1/n$  such that  $n \cdot 1/n = 1$ )
2. Define  $m \cdot 1/n := m/n$  when  $n \neq 0$ .

From this, we can construct the set of rational numbers.

**Definition 4.3.1 ▶ Rational Numbers  $\mathbb{Q}$** 

The set of **rational numbers** is defined as:

$$\mathbb{Q} := \left\{ \frac{m}{n} : m, n \in \mathbb{Z} \wedge n \neq 0 \right\}$$

To ensure multiplication works as intended, we also define  $\frac{m}{n} \cdot \frac{k}{l} := \frac{m \cdot k}{n \cdot l}$ .

We say  $\frac{m_1}{n_1} = \frac{m_2}{n_2}$  if and only if  $m_1 n_2 = m_2 n_1$  where  $n_1, n_2 \neq 0$ . In other words,  $\frac{m_1}{n_1} \sim \frac{m_2}{n_2} \iff m_1 n_2 = m_2 n_1$ . Thus,  $\mathbb{Q}$  is the set of equivalence classes for this relation.

If  $n = kp$  and  $m = kq$ , where  $k, p, q \in \mathbb{Z}$ ,  $k \neq 0$ ,  $q \neq 0$ , then:

$$\frac{n}{m} = \frac{kp}{kq} = \frac{k}{q}, \quad \text{because } kpq = kqp$$

If  $n$  and  $m$  have no common factor (except  $\pm 1$ ), then we say that  $n/m \in \mathbb{Q}$  is in the “lowest terms” or “reduced terms”. The set  $(\mathbb{Q}, +, \cdot)$  forms a field. However, we cannot write  $x = n/m$

where  $x^2 = 2$ .

### Theorem 4.3.2 ▶ $\sqrt{2}$ is not a Rational Number

$$\sqrt{2} \notin \mathbb{Q}$$

*Proof.* Suppose for contradiction  $\sqrt{2}$  is a rational number. Then, there exist  $n, m \in \mathbb{Z}$  such that  $(n/m)^2 = 2$ . If  $n = kp$  and  $m = kq$ , then we can “cancel” the common factor  $k$  to write  $n/m = p/q$ . That is, we can assume that  $n$  and  $m$  have no (non-trivial) common factors. Now,  $n^2/m^2 = 2$ , so by multiplying both sides by  $m^2$ , we get  $n^2 = 2m^2$ . Thus,  $n^2$  is an even number, so  $n$  is also even (Example 4.2.4). Then, we can write  $n = 2k$  where  $k \in \mathbb{Z}$ . Then:

$$\implies (2k)^2 = 2m^2$$

$$\implies 4k^2 = 2m^2$$

$$\implies 2k^2 = m^2$$

Then  $m^2$  is even, so  $m$  is even. Thus,  $m$  and  $n$  are both even, so they are multiples of 2. This contradicts the fact that we defined  $n/m$  in the lowest terms.  $\square$

Does there exist  $r \in \mathbb{Q}$  such that  $r^2 = 3$ ?

### Definition 4.3.3 ▶ Divides

For  $a, b \in \mathbb{Z}$ , we say  $a$  **divides**  $b$  if  $b$  is a multiple of  $a$ .

$$a \mid b \iff \exists(c \in \mathbb{Z})(b = ac)$$

### Theorem 4.3.4 ▶ Division Algorithm

Suppose  $a, b \in \mathbb{Z}$ . Then  $a = kb + r$  where  $k \in \mathbb{Z}$  and  $r \in \mathbb{Z}$  where  $0 \leq r < a$ .

### Example 4.3.5

If  $p \in \mathbb{N}$  and  $3 \mid p^2$ , then  $3 \mid p$ .

*Proof.* By the division algorithm,  $p = 3k + j$  where  $k \in \mathbb{Z}$  and  $j \in \mathbb{Z}$  where  $0 \leq j < 3$ . Then,  $p^2 = (3k + j)^2 = 9k^2 + 6kj + j^2$ . Suppose that  $3 \mid p^2$ . Then,  $p^2 = 3l = 9k^2 + 6kj + j^2$ .

Fix this whole section up. Very confusing.

Need proof here

Thus:

$$j^2 = 3l - 9k^2 - 6kj = 3(l - 3k^2 - 2kj)$$

We have  $3 \mid j^2$ . Hence,  $j \neq 1, j \neq 2$ , leaving only  $j = 0$ . Therefore,  $p = 3k + 0$ , so  $3 \mid p$ .  $\square$

#### Example 4.3.6 $\triangleright \sqrt{3}$ is not a Rational Number

*Proof.* Suppose for contradiction  $\sqrt{3}$  is a rational number. Then, there exist  $n, m \in \mathbb{Z}$  such that  $(n/m)^2$ . If  $n$  and  $m$  share a common factor, then we can “cancel” the common factor to where  $n/m = {}^kp/kq = p/q$ . Thus, we may assume that  $n$  and  $m$  have no nontrivial common factor.

$$\begin{aligned} \left(\frac{n}{m}\right)^2 &= 3 \\ \implies \frac{n^2}{m^2} &= 3 \\ \implies n^2 &= 3m^2 \end{aligned}$$

Thus,  $3 \mid n^2$ , so  $3 \mid n$  by the previous lemma. Writing  $n = 3k$  for some  $k \in \mathbb{Z}$ , we have:

$$\begin{aligned} (3k)^2 &= 3m^2 \\ \implies 9k^2 &= 3m^2 \\ \implies 3k^2 &= m^2 \end{aligned}$$

That is,  $3 \mid m^2$  so  $3 \mid m$ . Thus, 3 divides both  $n$  and  $m$ . This contradicts the fact that we defined  $n/m$  in the lowest terms.  $\square$

## 4.4 Fields

### Definition 4.4.1 $\triangleright$ Field

A **field** is a set  $F$  with two defined operations, addition and multiplication, satisfying the following for all  $a, b, c \in F$ :

Axiom	Addition	Multiplication
<b>Associativity</b>	$(a + b) + c = a + (b + c)$	$(ab)c = a(bc)$
<b>Commutativity</b>	$a + b = b + a$	$ab = ba$
<b>Distributivity</b>	$a(b + c) = ab + ac$	$(a + b)c = ac + bc$
<b>Identities</b>	$\exists(0 \in \mathbb{F})(a + 0 = a)$	$\exists(1 \in \mathbb{F})(1 \neq 0 \wedge 1a = a)$
<b>Inverses</b>	$\exists(-a \in \mathbb{F})(a + (-a) = 0)$	$(a \neq 0) \iff \exists(a^{-1} \in \mathbb{F})(aa^{-1} = 1)$

All the “standard facts” of arithmetic and algebra in  $\mathbb{R}$  follows from these axioms.

$\mathbb{Q}$ ,  $\mathbb{R}$ , and  $\mathbb{C}$  are infinite fields, but  $\mathbb{Z}_p$  (arithmetic modulo  $p$ ) is a finite field if  $p$  is prime.

More generally,  $F_q$  where  $q = p^k$  is a finite field.

#### Theorem 4.4.2 ► Facts about Fields

Let  $F$  be a field. For all  $a, b, c \in F$ :

- (a) if  $a + c = b + c$ , then  $a = b$
- (b)  $a \cdot 0 = 0$
- (c)  $(-a) \cdot b = -(a \cdot b)$
- (d)  $(-a) \cdot (-b) = a \cdot b$
- (e) if  $a \cdot c = b \cdot c$  and  $c \neq 0$ , then  $a = b$
- (f) if  $a \cdot b = 0$ , then  $a = 0$  or  $b = 0$
- (g)  $-(-a) = a$
- (h)  $-0 = 0$

*Proof of (g).*

$$\begin{aligned}
 -(-a) &= -(-a) + 0 \\
 &= -(-a) + (a + (-a)) \\
 &= -(-a) + (-a + a) \\
 &= (-(-a) + (-a)) + a \\
 &= ((-a) + -(-a)) + a \\
 &= 0 + a \\
 &= a + 0 \\
 &= a
 \end{aligned}$$



## 4.5 Ordered Fields

### Definition 4.5.1 ► Ordered Field

An **ordered field** is a field with a relation  $<$  such that for all  $a, b, c \in F$ :

Axiom	Description
<b>Trichotomy</b>	Only one is true: $a < b$ , $a = b$ , or $b < a$
<b>Transitivity</b>	if $a < b$ and $b < c$ then $a < c$
<b>Additive Property</b>	if $b < c$ , then $a + b < a + c$
<b>Multiplicative Property</b>	if $b < c$ and $0 < a$ , then $a \cdot b < a \cdot c$

We then define  $>$  as the inverse relation of  $<$ .

### Theorem 4.5.2 ► Facts about Ordered Fields

- if  $a < b$  then  $-b < -a$
- if  $a < b$  and  $c < 0$ , then  $cb < ca$
- if  $a \neq 0$ , then  $a^2 = a \cdot a > 0$
- $0 < 1$
- if  $0 < a < b$  then  $0 < 1/b < 1/a$

Although  $\mathbb{C}$  is a field, it is not an ordered field. We can certainly define some kind of “order” on  $\mathbb{C}$ , but there is no way to make it satisfy the four axioms of an ordered field. For example,  $i^2 = -1 < 0$ , contradicting the fact that any nonzero number’s square is greater than 0 in an ordered field.

$\mathbb{R}$  and  $\mathbb{Q}$  are ordered fields.

### Definition 4.5.3 ► Absolute Value

Let  $F$  be an ordered field. For  $a \in F$ , we define the **absolute value** of  $a$  as:

$$|a| := \begin{cases} a, & a \geq 0 \\ -a, & a < 0 \end{cases}$$

We can think of  $|a - b|$  as the distance between  $a$  and  $b$ . More generally,  $|a - b| = d(a, b)$  is the metric we will be using throughout real analysis.

#### Theorem 4.5.4 ► Properties of Absolute Value

- $|a| \geq 0$ ,  $a \leq |a|$ , and  $-a \leq |a|$
- $|ab| = |a||b|$

#### Theorem 4.5.5 ► Triangle Inequality

Let  $F$  be an ordered field. For any  $a, b \in F$ ,  $|a + b| \leq |a| + |b|$ .

*Proof.* There are two cases to consider. If  $a + b \geq 0$ , then:

$$\begin{aligned} |a + b| &= a + b \\ &\leq |a| + b \\ &\leq |a| + |b| \end{aligned}$$

If  $a + b < 0$ , then:

$$\begin{aligned} |a + b| &= -(a + b) \\ &= -a - b \\ &\leq |a| - b \\ &\leq |a| + |b| \end{aligned}$$

□

## 4.6 Completeness

#### Definition 4.6.1 ► Bounded Above, Bounded Below, Bounded

Let  $F$  be an ordered field, and let  $A \subseteq F$ .

- $A$  is **bounded above** if there exists  $b \in F$  such that  $a \leq b$  for all  $a \in A$ . In this context,  $b$  is an **upper bound** for  $A$ .
- $A$  is **bounded below** if there exists  $c \in F$  such that  $c \leq a$  for all  $a \in A$ . In this context,  $c$  is a **lower bound** for  $A$ .
- $A$  is **bounded** if  $A$  is bounded above and bounded below.

**Example 4.6.2 ▶ Upper and Lower Bounds**

Consider the set  $(0, 1) := \{x \in \mathbb{R} : 0 < x < 1\}$ .

- $(0, 1)$  is bounded above by 1 and any number greater than 1.
- $(0, 1)$  is bounded below by 0 and any negative number.

Consider the set  $[3, \infty) := \{x \in \mathbb{R} : 3 \leq x\}$ .

- $[3, \infty)$  is not bounded above.
- $[3, \infty)$  is bounded below by 3 and any number less than 3.

**Definition 4.6.3 ▶ Maximum, Minimum**

Let  $F$  be an ordered field, and let  $A \subseteq F$ .

- If there exists  $M \in A$  such that  $M$  is an upper bound for  $A$ , then  $M$  is the **maximum** of  $A$ , denoted  $M = \max A$
- If there exists  $m \in A$  such that  $m$  is a lower bound for  $A$ , then  $m$  is the **minimum** of  $A$ , denoted  $m = \min A$ .

Note that from the above example,  $(0, 1)$  has neither a maximum nor a minimum. However, 3 is the minimum of  $[3, \infty)$ .

**Definition 4.6.4 ▶ Supremum**

Let  $F$  be an ordered field, and let  $A \subseteq F$ .  $s \in F$  is a **supremum** of  $A$  if:

1.  $s$  is an upper bound for  $A$ , and
2. if  $t$  is an upper bound for  $A$ , then  $s \leq t$ .

In other words, the supremum is the least upper bound for  $A$ . If  $A$  has a supremum, then that supremum is unique. \_\_\_\_\_

Prove  
this

**Theorem 4.6.5 ▶ Maximum is the Supremum**

Let  $F$  be an ordered field, and let  $A \subseteq F$ . If  $A$  has a maximum  $M$ , then  $M = \sup A$ .

*Proof.* Since  $M = \max A$ , we know  $M$  is an upper bound for  $A$ . Let  $t$  be an upper bound for  $A$ . Since  $M \in A$ , then  $t \geq M$ . Thus,  $M$  is less than or equal to any upper bound  $t$ , so  $M = \sup A$ . □



**Example 4.6.6 ▶ Supremum of  $(0, 1)$** 

Prove that  $\sup(0, 1) = 1$ .

*Proof.* First, note that 1 is an upper bound for  $(0, 1)$ . Next, suppose that  $t \in \mathbb{Q}$  is an upper bound for  $(0, 1)$ . Since  $0 < 1/2 < 1$ , then  $0 < 1/2 \leq t$ . By transitivity,  $t > 0$ . Suppose for contradiction  $t < 1$ . Because  $0 < t < 1$ , we have  $1 < 1 + t < 2$ . Dividing across by 2, we have  $1/2 < 1 + t/2 < 1$ . That is,  $1 + t/2 \in (0, 1)$ . But  $t < 1$ , so  $2t < 1 + t$ . Thus,  $t < 1 + t/2$ . This contradicts our assumption that  $t$  is an upper bound for  $(0, 1)$ . Therefore,  $t \geq 1$ , so  $\sup(0, 1) = 1$ .  $\square$

**Definition 4.6.7 ▶ Completeness**

An ordered field  $F$  is **complete** if every nonempty subset of  $F$  that is bounded above has a supremum in  $F$ .

**Theorem 4.6.8 ▶  $\mathbb{Q}$  is not complete**

*Proof sketch.* Let  $A := \{x \in \mathbb{Q} : x^2 < 2\}$ . In other words,  $A = (-\sqrt{2}, \sqrt{2}) \subseteq \mathbb{Q}$ . Then  $A$  is nonempty and bounded above. Suppose for contradiction that  $\mathbb{Q}$  is complete. Then  $A$  has a supremum, say  $s = \sup(A)$ . Consider the following cases:

1. If  $s^2 < 2$ , let  $n \in \mathbb{N}$  such that  $(s + 1/n)^2 < 2$ . Then  $s + 1/n \in A$ , contradicting  $s$  being an upper bound for  $A$ .
2. If  $s^2 > 2$ , let  $n \in \mathbb{N}$  such that  $(s - 1/n)^2 > 2$ . Then  $s - 1/n$  is an upper bound smaller than  $s$ , contradicting  $s$  being the least upper bound (supremum).
3. If  $s^2 = 2$ , then  $s \notin \mathbb{Q}$  (Theorem 4.3.2).

Thus,  $A \subseteq \mathbb{Q}$  does not have a supremum. Therefore,  $\mathbb{Q}$  is not complete.  $\square$

**Definition 4.6.9 ▶ Real Numbers  $\mathbb{R}$** 

The **real numbers** are a set  $\mathbb{R}$  with two operations,  $+$  and  $\cdot$ , and order relation  $<$  such that:

1.  $(\mathbb{R}, +, \cdot)$  is a field,
2.  $(\mathbb{R}, +, \cdot, <)$  is an ordered field, and
3.  $(\mathbb{R}, +, \cdot, <)$  is complete.

Alternatively,  $\mathbb{R}$  can be constructed explicitly using “Dedekind cuts”. Either way,  $\mathbb{R}$  is the **only** unique complete ordered field up to isomorphism. That is, if there is some other imposter complete ordered field  $\mathbb{R}'$ , we can map every element of  $\mathbb{R}$  to  $\mathbb{R}'$  such that we preserve all the

operations and relations between things in  $\mathbb{R}$ . More formally, there exists an isomorphism  $T : \mathbb{R} \rightarrow \mathbb{R}'$  where  $T$  is bijective, and:

- $T(x + y) = T(x) + T(y)$
- $T(xy) = T(x)T(y)$
- $x < y \iff T(x) < T(y)$

Additionally,  $\mathbb{N} \subseteq \mathbb{R}$  where  $\mathbb{N}$  satisfies the Peano axioms.

#### Theorem 4.6.10 $\sqrt{2}$ is a Real Number

*Proof sketch.* Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

- Show  $A \neq \emptyset$  and  $A$  is bounded above
- Completeness says  $s := \sup A$  exists
- Show  $s^2 = 2 \implies s = \sqrt{2} \in \mathbb{R}$ .

More generally, if  $n, m \in \mathbb{N}$ , then  $\sqrt[n]{m} \in \mathbb{R}$ .



# Suprema and Infima

## Definition 5.0.1 ► Infimum

Let  $F$  be an ordered field, and let  $A \subseteq F$ .  $s$  is the *infimum* of  $A$  if:

1.  $s$  is a lower bound for  $A$ , and
2.  $s$  is greater than every other lower bound for  $A$ .

We can prove that the existence of infima is already implied by completeness.

## Theorem 5.0.2 ► Existence of Infima in $\mathbb{R}$

Let  $A \subseteq \mathbb{R}$  be nonempty and bounded below. Then  $A$  has an infimum in  $\mathbb{R}$ .

*Proof.* Let  $A \subseteq \mathbb{R}$  be nonempty and bounded below. Let  $B$  be the set of all lower bounds for  $A$ . In other words,  $B := \{b \in \mathbb{R} : \forall(a \in A)(b < a)\}$ . Since  $A$  is bounded below, then  $B$  is nonempty. Note also that  $B$  is bounded above by element of  $A$ . By completeness,  $s := \sup B$  exists. Now, we need to show that  $\sup B = \inf A$ .

1. Every  $a \in A$  is an upper bound for  $B$ , and  $\sup B$  is the least upper bound for  $B$ . Then,  $\sup B \leq a$ . That is,  $\sup B$  is a lower bound for  $A$ .
2. Let  $t$  be a lower bound for  $A$ . Then, by definition of  $B$ , it follows that  $t \in B$ . Then  $t \leq \sup B$  as required.

Therefore,  $\sup B = \inf A$  in  $\mathbb{R}$ . □

## Theorem 5.0.3 ► Well-Ordering Principle

Every non-empty subset of  $\mathbb{N}$  has a minimum.

*Proof.* We will use induction. For convenience, let  $P(n)$  represent the following statement: “If  $A \subseteq \mathbb{N}$  and  $A \cap \{1, 2, \dots, n\} \neq \emptyset$ , then  $A$  has a minimum.”

**Base Case:** First, we will prove  $P(1)$ . If  $A \subseteq \mathbb{N}$  and  $A \cap \{1\} \neq \emptyset$ , then  $1 \in A$ , so  $A$  has a minimum.

**Induction Step:** Assume that  $P(n)$  holds for some  $n \in \mathbb{N}$ . Suppose  $A \subseteq \mathbb{N}$  and  $A \cap \{1, 2, \dots, n+1\} \neq \emptyset$ .

1. If  $A \cap \{1, 2, \dots, n\} \neq \emptyset$ , then  $A$  has a minimum by  $P(n)$ .
2. If  $A \cap \{1, 2, \dots, n\} = \emptyset$ , then  $n + 1 \in A$ , so  $\min A = n + 1$ .

By induction,  $P(n)$  holds for all  $n \in \mathbb{N}$ . If  $A \subseteq \mathbb{N}$  and  $A \neq \emptyset$ , then there exists  $m \in A$  such that  $m \in \mathbb{N}$ . By  $P(m)$  (which is true by induction), the set  $A$  has a minimum.  $\square$

#### Theorem 5.0.4 ▶ Pushing Supremum

Let  $A$  be a nonempty subset of  $\mathbb{R}$ , and let  $b, c$  be real numbers.

- (a) If  $a \leq b$  for all  $a \in A$ , then  $\sup A \leq b$ .
- (b) If  $c \leq a$  for all  $a \in A$ , then  $c \leq \inf A$ .

**Intuition:** Consider the interval  $A := (0, 1)$ . Because  $a \leq 1$  for all  $a \in (0, 1)$ , we have  $\sup A \leq 1$ . Because  $0 \leq a$  for all  $a \in (0, 1)$ , we have  $0 \leq \inf A$ .

*Proof of (a).* Since  $a \leq b$  for all  $a \in A$ , then  $b$  is an upper bound for  $A$ . By completeness,  $A$  has a supremum, and  $s := \sup A$  is the least upper bound for  $A$ . Thus,  $s \leq b$ .  $\square$

*Proof of (b).*  $\square$

#### Example 5.0.5

Suppose  $A, B \subseteq \mathbb{R}$ ,  $A \neq \emptyset$ ,  $A \subseteq B$ , and  $B$  is bounded above. Prove that  $A$  is bounded above and  $\sup A \leq \sup B$ .

*Proof.* Since  $A \subseteq B$  and  $A \neq \emptyset$ , then  $B \neq \emptyset$ . Also,  $B$  is bounded above, so  $B$  has a supremum (by completeness). Let  $a \in A$  be arbitrary. Then  $a \in B$ , so  $a \leq \sup B$ . Thus,  $A$  is bounded above, so  $A$  has a supremum (by completeness). By Pushing Supremum,  $\sup A \leq \sup B$ .  $\square$

#### Theorem 5.0.6 ▶ Approximation Property of Suprema and Infima

Suppose  $A$  is a nonempty subset of  $\mathbb{R}$ , and  $s, r \in \mathbb{R}$ . Then:

- (a)  $s = \sup A$  if and only if (i)  $s$  is an upper bound for  $A$ , and (ii) for all  $\epsilon > 0$ , there exists  $a \in A$  such that  $s - \epsilon < a$ .
- (b)  $r = \inf A$  if and only if (i)  $r$  is a lower bound for  $A$ , and (ii) for all  $\epsilon > 0$ , there exists  $a \in A$  such that  $a < r + \epsilon$ .

**Intuition:** If we nudge the supremum ever so slightly to the left, then we must have moved past something in  $A$ .

*Proof of (a).* Let  $s := \sup A$ . Then (i) holds by definition of suprema. To prove (ii), let  $\epsilon > 0$ . Since  $s - \epsilon < s$ , then  $s - \epsilon$  is not an upper bound for  $A$ . Therefore, there exists  $a \in A$  such that  $s - \epsilon < a$ .

Conversely, suppose that (i) and (ii) hold. We need to show  $s = \sup A$ . From (i), we know that  $s$  is an upper bound for  $A$ . Now, we need to show that  $s$  is the least upper bound. Let  $t$  be an upper bound for  $A$ . Suppose for contradiction that  $t < s$ . Let  $\epsilon := s - t > 0$ . Then  $t = s - \epsilon$ . By (ii), there exists  $a \in A$  such that  $a > s - \epsilon = t$ . This contradicts  $t$  being an upper bound for  $A$ . Thus, there is no upper bound less than  $s$ . Therefore,  $s = \sup A$ .  $\square$

# Consequences of Completeness

## Theorem 6.0.1 ► $\mathbb{N}$ is not Bounded Above

*Proof.* Suppose for contradiction  $\mathbb{N}$  is bounded above. Since  $\mathbb{N}$  is not empty, then  $\mathbb{N}$  has a supremum in  $\mathbb{R}$ . Let  $s := \sup \mathbb{N} \in \mathbb{R}$ . Then  $n \leq s$  for all  $n \in \mathbb{N}$ . By the Peano axioms,  $n$  has a successor  $n + 1 \in \mathbb{N}$ , so  $n + 1 \leq s$  for all  $n \in \mathbb{N}$ . Therefore,  $n \leq s - 1$  for all  $n \in \mathbb{N}$ . This contradicts  $s$  being the least upper bound for  $\mathbb{N}$ .  $\square$

## Theorem 6.0.2 ► Archimedean Principle

Suppose  $x, y \in \mathbb{R}$  where  $x > 0$ . Then, there exists  $n \in \mathbb{N}$  such that  $nx > y$ .

**Intuition:** This is basically an extension of the fact that  $\mathbb{N}$  is not bounded above.

*Proof.* Since  $y/x$  is not an upper bound for  $\mathbb{N}$ , then there exists  $n \in \mathbb{N}$  such that  $n > y/x$ . Since  $x > 0$ , then  $nx > y$ .  $\square$

## Theorem 6.0.3 ► Density of $\mathbb{Q}$ in $\mathbb{R}$

Suppose  $x, y \in \mathbb{R}$  where  $x < y$ . Then there exists  $r \in \mathbb{Q}$  such that  $x < r < y$ .

**Intuition:** Given any two different real numbers, there's some rational number between them.

*Proof.* We will consider three cases:

1. If  $x \geq 0$ , then  $0 \leq x < y$ . Since  $y - x > 0$ , then by the Archimedean Principle, there exists  $n \in \mathbb{N}$  such that  $n(y - x) > 1$ . We want to show there is a natural number between  $nx$  and  $ny$ . Let  $A := \{k \in \mathbb{N} : k > nx\}$ . Since  $\mathbb{N}$  isn't bounded above, then  $A$  is not empty. By the Well-Ordering Principle,  $A$  has a minimum. Let  $m := \min A$ . Then  $m > nx$ , and  $m - 1 \leq nx$ . Thus,  $m \leq nx + 1$ , so:

$$nx < m \leq nx + 1 < ny$$

Dividing across by  $n$  yields  $x < m/n < y$ . Note that  $m, n \in \mathbb{N} \subseteq \mathbb{Z}$ , so  $m/n \in \mathbb{Q}$ .

2. If  $x < 0$  and  $y > 0$ , then  $x < 0 < y$  where  $0 \in \mathbb{Q}$ .
3. If  $x < 0$  and  $y \leq 0$ , then  $x < y \leq 0$ . Multiplying across by  $-1$ , we have  $-x > -y \geq 0$ . By the first case, there must exist  $t \in \mathbb{Q}$  where  $-y < t < -x$ . Multiply across by  $-1$  again to attain  $y > -t > x$  where  $-t \in \mathbb{Q}$ .

This completes the proof. □

### Theorem 6.0.4 ▶ $\sqrt{2}$ is a Real Number

There exists  $s \in \mathbb{R}$  such that  $s^2 = 2$ .

*Proof.* Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ . Since  $0^2 < 2$ , then  $0 \in A$ , so  $A$  is not empty. Also,  $A$  is bounded above, for example by 2. By completeness,  $A$  must have a supremum in  $\mathbb{R}$ . Let  $s := \sup A$ . We will use trichotomy to show that  $s^2 = 2$ .

1. If  $s^2 > 2$ , then...

**Scratchwork:** We need to show that this is not possible, i.e. show there is some  $s - 1/n$  that is less than  $s$  but is still an upper bound for  $A$ . We want  $(s - 1/n)^2 > 2$ . Then,  $s^2 - 2s/n + 1/n^2 > 2$ . We can chop off the  $1/n^2$ , reducing the inequality to  $s^2 - 2s/n > 2$ . Thus, we need to choose  $n > \frac{2s}{s^2-2}$ .

... let  $n \in \mathbb{N}$  such that  $n > \frac{2s}{s^2-2}$ . Then:

$$\begin{aligned}
 & n > \frac{2s}{s^2-2} \\
 \implies & s^2 - \frac{2s}{n} > 2 \\
 \implies & s^2 - \frac{2s}{n} + \frac{1}{n^2} > 2 \\
 \implies & \left(s - \frac{1}{n}\right)^2 > 2
 \end{aligned}$$

Thus,  $s - 1/n$  is an upper bound for  $A$  that is less than  $s$ . This contradicts  $s$  being the supremum for  $A$ .

2. If  $s^2 < 2$ , then...

**Scratchwork:** Again, we need to show that this is not possible. We know that in this case,  $s \in A$ , so we need to find another thing in  $A$  that is bigger than  $s$ . In other words, we want some  $(s + 1/n)^2 < 2$ . Then,  $s^2 + 2s/n + 1/n^2 < 2$ . Choose  $n > 1/2s$  and  $n > \frac{4s}{2-s^2}$ .

$$\left(s + \frac{1}{n}\right)^2 = s^2 + \frac{2s}{n} + \frac{1}{n^2}$$

... let  $n \in \mathbb{N}$  such that  $n > \max\left\{\frac{1}{2s}, \frac{4s}{2-s^2}\right\}$ . Then  $\frac{1}{n} < 2s$  and  $s^2 + \frac{4s}{n} < 2$ . So:

$$\begin{aligned} \left(s + \frac{1}{n}\right)^2 &= s^2 + \frac{2s}{n} + \frac{1}{n^2} \\ &< s^2 + \frac{2s}{n} + \frac{2s}{n} \\ &= s^2 + \frac{4s}{n} < 2 \end{aligned}$$

That is,  $s + \frac{1}{n} \in A$ . This contradicts  $s$  being an upper bound for  $A$ .  
By trichotomy,  $s^2 = 2$ . □

### Theorem 6.0.5 ▶ Nested Interval Property

Suppose that for each  $n \in \mathbb{N}$ ,  $a_n, b_n \in \mathbb{R}$  with  $a_n \leq b_n$ , and  $a_n \leq a_{n+1} \leq b_{n+1} \leq b_n$  for all  $n \in \mathbb{N}$ . Then  $\bigcap_{n=1}^{\infty} [a_n, b_n] \neq \emptyset$ .

**Intuition:** We can move the two borders of an open interval closer and closer to each other, and it won't be empty.

*Proof.* Note that  $a_n \leq a_{n+1} \leq a_{n+2} \leq \dots$  and  $\dots \leq b_{n+2} \leq b_{n+1} \leq b_n$ . If  $k \leq n$ , then  $a_k \leq a_n \leq b_n$ .

- If  $k \leq n$ , then  $a_k \leq a_n \leq b_n$ .
- If  $k \geq n$ , then  $a_k \leq b_k \leq b_n$ .

That is,  $a_k \leq b_n$  for all  $k, n \in \mathbb{N}$ . Let  $A := \{a_k : k \in \mathbb{N}\}$ . Then  $A$  is bounded above, for example by  $b_1$ . Also,  $A$  is not empty. By completeness,  $A$  has a supremum. Let  $s := \sup A$ . Note that since  $s$  is an upper bound for  $A$ , then  $a_n \leq \sup A$  for all  $n \in \mathbb{N}$ . Also note that  $\sup A$  is the least upper bound for  $A$ , so  $\sup A \leq b_n$  for all  $n \in \mathbb{N}$ . Thus,  $a_n \leq \sup A \leq b_n$  for all  $n \in \mathbb{N}$ , so  $\sup A \in [a_n, b_n]$  for all  $n \in \mathbb{N}$ . Thus,  $\sup A \in \bigcap_{n=1}^{\infty} [a_n, b_n]$ , so it is not empty. □



The nested interval property is actually false for open intervals!

$$\forall (x \in (0, 1)) \exists (n \in \mathbb{N}) (1/n < x \implies x \notin (0, 1/n))$$

# Cardinality

## Definition 7.0.1 ► Cardinality

**Cardinality** is a measure of the amount of elements in a set, denoted  $|A|$ . We say two sets have the same cardinality if there exists a bijection between them.

For finite sets, we can think of cardinality as the number of elements in that set. For infinite sets, cardinality can sometimes go against our intuition. For any sets  $A, B, C$ :

1.  $|A| = |A|$
2. if  $|A| = |B|$ , then  $|B| = |A|$
3. if  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

Hence, equality of cardinalities is an equivalence relation.

## Example 7.0.2 ► Cardinality of $\mathbb{N}$ and $2\mathbb{N}$

Let  $2\mathbb{N} := \{2n : n \in \mathbb{N}\}$  (i.e. the set of even natural numbers). Then  $|\mathbb{N}| = |2\mathbb{N}|$ .

*Proof.* To show that these two sets have the same cardinality, we need to find some bijection between the sets. Let  $f : \mathbb{N} \rightarrow 2\mathbb{N}$  be a function defined by  $f(n) = 2n$ . Note that  $f$  is well-defined (i.e. is actually a function) because  $f(n) \in 2\mathbb{N}$  for all  $n \in \mathbb{N}$ . To prove that  $f$  is a bijection, we need to prove it is both injective and surjective.

1. Let  $n_1, n_2 \in \mathbb{N}$  such that  $f(n_1) = f(n_2)$ . Then  $2n_1 = 2n_2$ , so  $n_1 = n_2$ . Thus,  $f$  is injective.
2. Let  $m \in 2\mathbb{N}$ . Then  $m = 2k$  for some  $k \in \mathbb{N}$ , so  $m = 2k = f(k)$  for some  $k \in \mathbb{N}$ . Thus,  $f$  is surjective.

Therefore,  $f$  is a bijection, so  $|\mathbb{N}| = |2\mathbb{N}|$ . □

## Example 7.0.3 ► Cardinality of Intervals

Let  $a, b \in \mathbb{R}$  where  $a < b$ . Then  $|(0, 1)| = |(a, b)|$ .

*Proof.* We need to find a bijection from  $(0, 1)$  to  $(a, b)$ . We need to “scale” the interval  $(0, 1)$  to the width of  $(a, b)$ , then translate it to match  $(a, b)$ . Define  $f : (0, 1) \rightarrow (a, b)$  by  $f(x) = a + (b - a)x$ . (We need to check  $f$  is well-defined). Let  $x \in (0, 1)$ . Then  $0 < x < 1$ , so multiplying by  $(b - a)$  which is positive gives  $0 < (b - a)x < b - a$ . Adding  $a$ , we get  $a < a + (b - a)x < b$ . Now we need to show  $f$  is a bijection:

1. Let  $x_1, x_2 \in (0, 1)$  such that  $f(x_1) = f(x_2)$ . Then  $a + (b - a)x_1 = a + (b - a)x_2$ . Subtracting  $a$  from both sides, we get  $(b - a)x_1 = (b - a)x_2$ . Since  $(b - a) \neq 0$ , we can divide both side by  $(b - a)$  to get  $x_1 = x_2$ .
2. Let  $y \in (a, b)$ .

**Scratchwork:** We want to find some  $x \in (0, 1)$  where  $y = f(x) = a + (b - a)x$ . Using some algebra to solve for  $x$ , we have  $x = \frac{y - a}{b - a}$

Let  $x = \frac{y - a}{b - a}$ . First, we show  $x \in (0, 1)$ :

$$\begin{aligned} & a < y < b \\ \implies & 0 < y - a < b - a \\ \implies & 0 < \frac{y - a}{b - a} < 1 \end{aligned}$$

Thus,  $x \in (0, 1)$ . Also:

$$f(x) = a + (b - a)\left(\frac{y - a}{b - a}\right) = a + (y - a) = y$$

Thus,  $f$  is surjective.

Therefore,  $f$  is a bijective, so  $|(0, 1)| = |(a, b)|$ . □

#### Definition 7.0.4 ► Power Set

Let  $A$  be a set. The **power set** of  $A$  is the set of all subsets of  $A$ .

$$\mathcal{P}(A) = \{B : B \subseteq A\}$$

For example, the power set of  $\{1, 2, 3\}$  is  $\{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$ . For any finite set with  $n$  elements in it, its power set has  $2^n$  elements in it.

**Example 7.0.5 ▶ Cardinality of  $\mathbb{N}$  and  $\mathcal{P}(\mathbb{N})$** 

$$|\mathbb{N}| \neq |\mathcal{P}(\mathbb{N})|$$

*Proof.* We will show that any function  $f : \mathbb{N} \rightarrow \mathcal{P}(\mathbb{N})$  cannot be surjective, and thus not bijective. Let  $f : \mathbb{N} \rightarrow \mathcal{P}(\mathbb{N})$  be any function defined by  $f(n) = A_n$ . Note  $A_n \subseteq \mathbb{N}$ , so  $A_n \in \mathcal{P}(\mathbb{N})$ . Now we will define a set that isn't in  $f[\mathbb{N}]$ . For each  $n \in \mathbb{N}$ , if  $n \in A_n$ , then  $n \notin A$ , and if  $n \notin A_n$ , then  $n \in A$ . More formally,  $A := \{n \in \mathbb{N} : n \notin A_n\}$ . For all  $k \in \mathbb{N}$ , note that:

- if  $k \in A_k$ , then  $k \notin A$ , so  $A \neq A_k$ , and
- if  $k \notin A_k$ , then  $k \in A_k$ , so  $A \neq A_k$ .

Hence,  $A \subseteq \mathbb{N}$ , but  $f(k) \neq A$  for any  $k \in \mathbb{N}$ . Thus,  $f$  is not surjective. □

**Definition 7.0.6 ▶ Finite, Countably Infinite, Countable, Uncountable**

Let  $A$  be a set. We say  $A$  is:

- **finite** if  $A \neq \emptyset$  or  $|A| = |\{1, 2, \dots, n\}|$  for some  $n \in \mathbb{N}$ .
- **countably infinite** if  $|A| = |\mathbb{N}|$ .
- **countable** if  $A$  is finite or countably infinite
- **uncountable** if  $A$  is not countable

**Theorem 7.0.7 ▶  $\mathcal{P}(\mathbb{N})$  is uncountable.**

*Proof.* We know from Example 7.0.5 that  $\mathcal{P}(\mathbb{N})$  is not countably infinite. We need to show that  $\mathcal{P}(\mathbb{N})$  is not finite. Since  $\{1\} \in \mathcal{P}(\mathbb{N})$ , then it cannot be empty. Suppose for contradiction  $|\{1, 2, \dots, n\}| = |\mathcal{P}(\mathbb{N})|$  for some  $n \in \mathbb{N}$ , then there exists a bijection  $f : \{1, 2, \dots, n\} \rightarrow \mathcal{P}(\mathbb{N})$ . Define  $g : \mathbb{N} \rightarrow \{1, 2, \dots, n\}$  by:

$$g(k) = \begin{cases} k, & 1 \leq k \leq n \\ 1, & k > n \end{cases}$$

Then  $g$  is surjective, so  $f \circ g : \mathbb{N} \rightarrow \mathcal{P}(\mathbb{N})$  is surjective. This contradicts the fact that no such function exists (by Example 7.0.5). □

Generally, there is never a bijection from a set to its power set.

**Intuition:** A set is countable if its elements can be “listed” or “counted”. That is, for finite sets:

$$X = \{x_1, x_2, \dots, x_n\} = \{x_k\}_{k=1}^n$$

For infinitely countable sets:

$$X = \{x_1, x_2, \dots\} = \{x_k\}_{k=1}^{\infty}$$

If  $X$  is finite, then there exists a bijection  $f : \{1, 2, \dots, n\} \rightarrow X$ . Thus,  $X = \{f(1), f(2), \dots, f(n)\}$ . If  $X$  is countably infinite, then there exists a bijection  $f : \mathbb{N} \rightarrow X$ . Thus,  $X = \{f(1), f(2), \dots\}$ .

### Theorem 7.0.8 ► Subsets of Countable Sets are Countable

The subset of a countable set is still countable. (i.e. a countable set cannot contain an uncountable subset).

*Proof.* Let  $X$  be a countable set, and let  $A \subseteq X$ . We will consider two cases. First, if  $A$  is finite, then  $A$  is countable, and we are done. Otherwise,  $A$  is infinite, and hence  $X$  is infinite. Then  $X$  is countably infinite, so  $X = \{x_1, x_2, \dots\} = \{x_k\}_{k=1}^{\infty}$ .

**Idea:** Our set  $A$  might look something like  $\{x_3, x_4, x_6, \dots\}$ . We need to align these indices to 1, 2, 3, and so on. We'll let  $k_1 = \min\{3, 4, 6, \dots\}$ , let  $k_2 = \min\{4, 6, \dots\}$ , and so on.

Let  $k_1 := \min\{k \in \mathbb{N} : x_k \in A\}$ . Let  $a_1 := x_{k_1}$ . For all  $j \in \mathbb{N}$  such that  $j > 1$ , we define  $k_j := \min\{k \in \mathbb{N} : (x_k \in A) \wedge (k > k_{j-1})\}$ . Let  $a_j := x_{k_j}$ . Then  $1 \leq k_1 < k_2 < k_3 < \dots$ , so  $k_j$  approaches infinity. Let  $g : \mathbb{N} \rightarrow A$  be a function defined by  $g(j) = a_j$ . We need to show that  $g$  is both injective and surjective, and thus a bijection.

- Suppose that  $g(j_1) = g(j_2)$  for some  $j_1, j_2 \in \mathbb{N}$ . Then  $a_{j_1} = a_{j_2}$ , so  $x_{k_{j_1}} = x_{k_{j_2}}$ . Then  $k_{j_1} = k_{j_2}$ , so  $j_1 = j_2$ . Thus,  $g$  is injective.
- Let  $a \in A$ . Since  $A \subseteq X$ , then  $a \in X$ . Thus,  $a = x_l$  for some  $l \in \mathbb{N}$ . Let  $m := \min\{j \in \mathbb{N} : k_j \geq l\}$ . Since  $m \in \{j \in \mathbb{N} : k_j \geq l\}$ , then  $k_m \geq l$ . Also,  $m - 1 \notin \{j \in \mathbb{N} : k_j \geq l\}$ , so  $k_{m-1} < l$ . Now,  $k_m = \min\{k \in \mathbb{N} : (x_k \in A) \wedge (k > k_{m-1})\}$ . But  $x_l \in A$ , and  $l > k_{m-1}$ , so  $l \in \{k \in \mathbb{N} : (x_k \in A) \wedge (k > k_{m-1})\}$ . Thus,  $k_m \leq l$ , because  $k_m$  is the minimum of the set containing  $l$ . By trichotomy,  $k_m = l$ . Therefore:

$$g(m) = a_m = x_{k_m} = x_l = a$$

So  $g$  is surjective.

Since  $g$  is a bijection, then  $|\mathbb{N}| = |A|$ , so  $|A|$  is countable.  $\square$

### Theorem 7.0.9 ► Injectivity and Cardinality

A set  $A$  is countable if and only if there exists an injective function  $f : A \rightarrow \mathbb{N}$ .

*Proof.* First, suppose  $A$  is a countable set. We consider two cases:

- If  $A$  is countably infinite, then there exists a bijection  $f : A \rightarrow \mathbb{N}$ .
- If  $A$  is finite, then there exists a bijection  $f : A \rightarrow \{1, 2, \dots, n\}$  for some  $n \in \mathbb{N}$ . Let  $g : \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  be a function defined by  $g(x) = x$  (i.e. an inclusion mapping). Then  $f$  and  $g$  are both injective, so  $g \circ f : A \rightarrow \mathbb{N}$  is injective.

Conversely, suppose  $f : A \rightarrow \mathbb{N}$  is an injection. Then  $f[A] \subseteq \mathbb{N}$ , so  $f[A]$  is countable by Theorem 7.0.8. Define  $g : A \rightarrow f[A]$  by  $g(a) = f(a)$ . Then  $g$  is injective because  $f$  is injective, and  $g$  is surjective because  $g[A] = f[A]$ . Thus,  $g$  is a bijection, so  $|A| = |f[A]|$ . Therefore,  $A$  is countable.  $\square$

### Theorem 7.0.10 ► $|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$

$\mathbb{N} \times \mathbb{N}$  is countable.

*Proof.* Let  $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  be a function defined by  $f(n, m) = 2^n 3^m$ . We now show that  $f$  is bijective. To prove  $f$  is injective, suppose  $f(n_1, m_1) = f(n_2, m_2)$ . Then  $2^{n_1} 3^{m_1} = 2^{n_2} 3^{m_2}$ .

- If  $n_1 > n_2$ , then  $2^{n_1 - n_2} = 3^{m_2 - m_1}$ . Since  $n_1 > n_2$ , we have  $n_1 - n_2 > 0$ , so  $2^{n_1 - n_2} \in \mathbb{N}$ . Then also  $3^{m_2 - m_1} \in \mathbb{N}$ . But  $2^{n_1 - n_2}$  is even, and  $3^{m_2 - m_1}$  is odd. This contradicts the fact that  $2^{n_1 - n_2} = 3^{m_2 - m_1}$ .
- If  $n_2 > n_1$ , then  $3^{m_1 - m_2} = 2^{n_2 - n_1}$ . By a similar argument,  $2^{n_2 - n_1}$  is even and  $3^{m_1 - m_2}$  is odd, producing the same contradiction.
- If  $n_1 = n_2$ , then  $2^{n_1} = 2^{n_2}$ , so cancelling gives  $3^{m_1} = 3^{m_2}$ . Thus,  $m_1 = m_2$ .

Hence,  $(n_1, m_1) = (n_2, m_2)$ , so  $f$  is injective. By Theorem 7.0.9,  $\mathbb{N} \times \mathbb{N}$  is countable. Also,  $\mathbb{N} \times \mathbb{N}$  is infinite, so  $|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$ .  $\square$

### Theorem 7.0.11 ► Collection of Countable Sets

Suppose that for all  $k \in \mathbb{N}$ ,  $A_k$  is a countable set. Then  $\cup_k A_k \subseteq \mathbb{N}$  is countable. (i.e. a countable union of countable sets is countable)

*Proof.* Let  $k \in \mathbb{N}$ .  $A_k$  is countable, so  $A_k$  can be listed as such:

$$\begin{aligned} A_1 &= \{a_{11}, a_{12}, a_{13}, a_{14}, \dots\} = \{a_j\}_{j \in \mathbb{N}} \\ A_2 &= \{a_{21}, a_{22}, a_{23}, a_{24}, \dots\} \\ &\vdots \\ A_k &= \{a_{k1}, a_{k2}, a_{k3}, a_{k4}, \dots\} \end{aligned}$$

We want to define some function  $f : \bigcup_{k \in \mathbb{N}} A_k \rightarrow \mathbb{N} \times \mathbb{N}$  where  $f(a_{kj}) = (k, j) \in \mathbb{N} \times \mathbb{N}$ . However, we need to consider the possibility that the sets  $A_k$  are not disjoint. If  $a_{12} = a_{34}$ , then  $a_{(12)} = (1, 2)$  and  $f(a_{34}) = (3, 4)$ .

Given  $a \in \bigcup_{k \in \mathbb{N}} A_k$ , let  $k(a) := \min\{k \in \mathbb{N} : a \in A_k\}$ . If  $a \in A_{k(a)}$ , then there is a unique  $j(a) \in \mathbb{N}$  such that  $a = a_{k(a)j(a)}$ . Now define  $f : \bigcup_{k \in \mathbb{N}} A_k \rightarrow \mathbb{N} \times \mathbb{N}$  by  $f(a) = (k(a), j(a))$ . We must show that  $f$  is injective. Let  $x, y \in \bigcup_{k \in \mathbb{N}} A_k$  such that  $f(x) = f(y)$ . That is,  $(k(x), j(x)) = (k(y), j(y))$ . Then  $x = a_{k(x)j(x)} = a_{k(y)j(y)} = y$ . By Theorem 7.0.10, there exists some injection  $g : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ . Hence,  $g \circ f : \bigcup_{k \in \mathbb{N}} A_k \rightarrow \mathbb{N}$  is injective. By Theorem 7.0.9,  $\bigcup_{k \in \mathbb{N}} A_k$  is countable.  $\square$

This theorem shows that, in order to prove a countable union of countable sets is countable, we just need to show that each set in the union is countable. We'll use this in our proof that the set of rational numbers is a countable set.

### Theorem 7.0.12 $\blacktriangleright$ $\mathbb{Q}$ is Countable

*Proof.* Let  $\mathbb{Q}^+ := \{r \in \mathbb{Q} : r > 0\}$ , and let  $\mathbb{Q}^- := \{r \in \mathbb{Q} : r < 0\}$ . First, we'll prove that  $\mathbb{Q}^+$  is countable. Let  $f : \mathbb{Q}^+ \rightarrow \mathbb{N} \times \mathbb{N}$  be a function defined as  $f(r) = (p, q)$  such that  $r = p/q$  where  $p, q \in \mathbb{N}$  and  $p$  shares no common factors with  $q$ . To show  $f$  is injective, let  $r_1, r_2 \in \mathbb{Q}^+$  where  $f(r_1) = f(r_2)$ . Then  $r_1 = p_1/q_1, r_2 = p_2/q_2$  where  $p_1, q_1 \in \mathbb{N}$  with no common factors, and  $p_2, q_2 \in \mathbb{N}$  with no common factors. Thus,  $(p_1, q_1) = (p_2, q_2)$ , so  $p_1 = p_2$  and  $q_1 = q_2$ . Thus,  $r_1 = p_1/q_1 = p_2/q_2 = r_2$ , so  $f$  is injective. Since there exists an injection  $g : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ , then  $g \circ f : \mathbb{Q}^+ \rightarrow \mathbb{N}$  is injective. Thus,  $\mathbb{Q}^+$  is countable.

Next, we'll prove that  $\mathbb{Q}^-$  is countable. Let  $h : \mathbb{Q}^- \rightarrow \mathbb{Q}^+$  by  $h(r) = -r$ . We show  $h$  is injective. If  $h(r_1) = h(r_2)$  where  $r_1, r_2 \in \mathbb{Q}^-$ , then  $-r_1 = -r_2$ , so  $r_1 = r_2$ . Thus,  $h$  is injective. From above, there exist an injection  $\phi : \mathbb{Q}^+ \rightarrow \mathbb{N}$ . Hence,  $h \circ \phi : \mathbb{Q}^{-1} \rightarrow \mathbb{N}$  is injective.

Finally,  $\{0\}$  is countable because it is finite. Since  $\mathbb{Q} = \mathbb{Q}^+ \cup \{0\} \cup \mathbb{Q}^-$  is a countable union of countable sets, then  $\mathbb{Q}$  is countable.  $\square$

This means we can “list” the rational numbers (disregarding order) as  $\mathbb{Q} = \{r_1, r_2, \dots\} = \{r_n\}_{n \in \mathbb{N}}$ .

### Theorem 7.0.13 ▶ $\mathbb{R}$ is Uncountable

*Proof.* Suppose for contradiction  $\mathbb{R}$  is countable. Then  $\mathbb{R}$  can be “listed” as  $\mathbb{R} = \{x_1, x_2, \dots\} = \{x_n\}_{n \in \mathbb{N}}$ . We will define a sequence of non-empty closed intervals  $\{I_k\}_{k \in \mathbb{N}}$  such that  $I_{k+1} \subseteq I_k$  and  $x_k \notin I_k$  for all  $k \in \mathbb{N}$ . Let  $I_0 := [0, 1]$ . Divide  $I_0$  into three equal closed intervals. Then, at least one of these three intervals does not contain  $x_1$ . Choose such an interval and call it  $I_1$ . Divide  $I_1$  into three equal closed intervals. Then, at least one of these three intervals does not contain  $x_2$ . Choose such an interval and call it  $I_2$ . Given  $I_k$  for some  $k \in \mathbb{N}$ , divide  $I_k$  into three equal closed intervals, then choose the interval that does not contain  $x_{k+1}$  and call it  $I_{k+1}$ . By induction, we have  $\{I_k\}_{k=1}^\infty$  where each  $I_k$  is a (nonempty) closed interval, and  $I_{k+1} \subseteq I_k$  for each  $k \in \mathbb{N}$ . By the Nested Interval Property,  $\bigcap_{k \in \mathbb{N}} I_k$  is not empty, so there exists  $x \in \mathbb{R}$  such that  $x \in \bigcap_{k \in \mathbb{N}} I_k$ . Since  $x \in \mathbb{R}$ , we have  $x = x_n$  for some  $n \in \mathbb{N}$  (by our supposition that  $\mathbb{R}$  is countable). However, we constructed  $I_n$  such that  $x_n \notin I_n$ , so  $x_n \notin \bigcap_{k \in \mathbb{N}} I_k$ . This contradiction renders our initial supposition false. Therefore,  $\mathbb{R}$  is uncountable.  $\square$

### Theorem 7.0.14 ▶ Irrational Numbers are Uncountable

*Proof.* Suppose for contradiction  $\mathbb{R} \setminus \mathbb{Q}$  is countable. Then  $\mathbb{R} = (\mathbb{R} \setminus \mathbb{Q}) \cup \mathbb{Q}$ , so  $\mathbb{R}$  is a countable union of countable sets, making  $\mathbb{R}$  countable. This contradicts the fact that  $\mathbb{R}$  is uncountable, so  $\mathbb{R} \setminus \mathbb{Q}$  is uncountable.  $\square$

## 7.1 Additional Remarks

### Definition 7.1.1 ▶ Algebraic Number, Transcendental Number

If  $\alpha$  is a root of the polynomial  $a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 = 0$  where each  $a_i \in \mathbb{Z}$ , then  $\alpha$  is called an **algebraic number**. If  $\alpha$  is not algebraic, then we call it a **transcendental number**.

The set of all algebraic numbers is countable, so “most” real numbers are transcendental.

- Even though  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , there are “more” irrational numbers than rational numbers



- The set  $\{\sqrt[n]{m} : n, m \in \mathbb{N}\}$  is countable, so “most” real numbers are not radicals.
- The set of algebraic numbers is countable, so “most” real numbers are transcendental.
- $|\mathcal{P}(\mathbb{N})| = |\mathbb{R}|$ , but the set  $\{A \subseteq \mathbb{N} : A \text{ is finite}\}$  is countable.
- If there exists an injection  $f : A \rightarrow B$ , then we say  $|A| \leq |B|$ . If there exists an injection from  $A$  to  $B$ , but there does not exist an injection from  $B$  to  $A$ , then we say  $|A| < |B|$ .

### Theorem 7.1.2 ▶ Schroeder-Bernstein

If there exists an injection  $f : A \rightarrow B$  and an injection  $g : B \rightarrow A$ , then there exists a bijection  $h : A \rightarrow B$ .

### Theorem 7.1.3 ▶ Continuum Hypothesis

There is no cardinality between  $|\mathbb{N}|$  and  $|\mathbb{R}|$ .

In Zermelo-Fraenkel with Choice (ZFC) set theory, the continuum hypothesis cannot be proven to be true nor false. In 1938, Godel proved that the continuum hypothesis is consistent with ZFC. In 1963, Cohen proved that the negation of the continuum hypothesis is also consistent with ZFC.

Just because you can write a description of a set does not mean that the set exists nor makes sense.

- For example, let  $A := \{\bigcup\{B : B \text{ is a set}\}\}$ . Then  $\mathcal{P}(A) \subseteq A$ , so  $|\mathcal{P}(A)| \leq |A| < |\mathcal{P}(A)|$ .
- Another example: let  $B := \{\text{all sets}\}$ . Let  $C := \{A : A \notin A\}$ . Is  $C \in C$ ?

# Sequences and Convergence

## Definition 8.0.1 ► Sequence

A **sequence** is an ordered list of real numbers.

$$s = (s_1, s_2, s_3, s_4, \dots)$$

Formally, a **sequence** is a function  $s : \mathbb{N} \rightarrow \mathbb{R}$ . We write  $s_n$  to denote  $s(n)$ .

We can define a sequence using an expression, like  $s_n := n^2$ . Then  $s = (1, 4, 9, 16, \dots)$ . Also, we can informally define a sequence in terms of its elements, like  $s = (3, 1, 4, 1, 5, 9, \dots)$ . We could just have a random sequence like  $s := (12.3, e^2, 1 - \pi, 10000, \dots)$ .

Let's consider how we can formalize the definitions of limits and convergence. Consider the sequence  $s_n := 1/n$ , then  $(s_n) = (1, 1/2, 1/3, 1/4, \dots)$ . We have an intuitive idea that, as  $n$  gets bigger, then  $1/n$  gets closer to 0. We can say that this sequence “converges” to 0.

Now consider the sequence  $s := (1, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, \dots)$ . Does this sequence converge? This really depends on our definition of convergence. We might define this as, “ $s_n$  gets close to  $l$  as  $n$  gets large”. It certainly matches our intuition, but what exactly does “close to  $l$ ” mean? Maybe we could say, “ $|s_n - l|$  gets small as  $n$  gets large”. More precisely, this might be “for all  $\epsilon > 0$ ,  $|s_n - l| < \epsilon$  when  $n$  is large”. That “ $n$  is large” is still imprecise. Fixing that part, we get the formal definition for convergence:

## Definition 8.0.2 ► Convergence

Let  $s := (s_n)_{n \in \mathbb{N}}$  be a sequence of real numbers, and let  $l \in \mathbb{R}$ . We say  $s_n$  **converges** to  $l$  if, for all  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $|s_n - l| < \epsilon$  for all  $n > N$ .

$$\forall(\epsilon > 0) \exists(N \in \mathbb{N}) \forall(n > N) (|s_n - l| < \epsilon)$$

Like in the approximation property, we use  $\epsilon$  to denote some arbitrarily tiny value that's really really close to 0, but not actually 0. We can also write  $\lim_{n \rightarrow \infty} s_n = l$  or  $s_n \rightarrow l$  to mean  $s_n$  converges to  $l$ .

**Technique 8.0.3 ▶ Proving Convergence**

To prove that a sequence  $s$  converges to  $l$ , we carry out the following steps:

1. As some scratch work, solve the inequality  $|s_n - l| < \epsilon$  for  $n$ .
2. In the formal proof, let  $\epsilon > 0$ , and let  $N$  be greater than the solved thing. Let  $n > N$ , then work towards  $|s_n - l| < \epsilon$ .

Make  
this ex-  
planation  
better

**Example 8.0.4 ▶  $1/n$  converges to 0**

Prove that  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ .

**Intuition:** Since we're proving something for all  $\epsilon > 0$ , let's start by choosing some arbitrary  $\epsilon > 0$ . Next, we need to choose some  $N \in \mathbb{N}$  where  $|s_n - l| < \epsilon$  for all  $n > N$ . Thus:

$$\begin{aligned} |s_n - l| &< \epsilon \\ \left| \frac{1}{n} - 0 \right| &< \epsilon \\ \frac{1}{n} &< \epsilon \\ n &> \frac{1}{\epsilon} \end{aligned}$$

So we choose  $N > \frac{1}{\epsilon}$ .

*Proof.* Let  $\epsilon > 0$ . Let  $N \in \mathbb{N}$  where  $N > 1/\epsilon$ . If  $n > N > 1/\epsilon$ , then  $1/n < \epsilon$ . Thus:

$$|s_n - l| = |1/n - 0| = 1/n < \epsilon$$

Therefore,  $s$  converges to 0. □

**Example 8.0.5**

Prove that  $\lim_{n \rightarrow \infty} \frac{2n+3}{3n+7} = \frac{2}{3}$ .

**Intuition:** This time, we want to choose some  $N \in \mathbb{N}$  such that  $|s_n - l| < \epsilon$ . Thus:

$$\begin{aligned} \left| \frac{2n+3}{3n+7} - \frac{2}{3} \right| &< \epsilon \\ \left| \frac{6n+9-6n-14}{9n+21} \right| &< \epsilon \\ \frac{5}{9n+21} &< \epsilon \\ \frac{5}{\epsilon} &< 9n+21 \\ \frac{1}{9} \left( \frac{5}{\epsilon} - 21 \right) &< n \end{aligned}$$

Thus, we want to choose  $N > \frac{1}{9}(\frac{5}{\epsilon} - 21)$ .

*Proof.* Let  $\epsilon > 0$ . Let  $N \in \mathbb{N}$  such that  $N > \frac{1}{9}(\frac{5}{\epsilon} - 21)$ . If  $n > N > \frac{1}{9}(\frac{5}{\epsilon} - 21)$ , then:

$$\begin{aligned} 9n &> \frac{5}{\epsilon} - 21 \\ 9n &> \frac{5}{\epsilon} - 21 \\ 9n + 21 &> \frac{5}{\epsilon} \\ \frac{5}{9n+21} &< \epsilon \end{aligned}$$

Thus:

$$\begin{aligned} |s_n - l| &= \left| \frac{2n+3}{3n+7} - \frac{2}{3} \right| \\ &= \left| \frac{6n+9-6n-14}{9n+21} \right| \\ &= \frac{5}{9n+21} \\ &< \epsilon \end{aligned}$$

□

The above proof chooses a sort of “optimal” or “best possible”  $N$ . We could have thrown away the 21 in the denominator, and the inequality we’re aiming for will still be the same.

*Alternate proof.* Let  $\epsilon > 0$ . Let  $N \in \mathbb{N}$  such that  $N > \frac{5}{9\epsilon}$ . If  $n > N > \frac{5}{9\epsilon}$ , then  $\frac{5}{9n} < \epsilon$ , so  $\frac{5}{9n+21} < \frac{5}{9n} < \epsilon$ . Then:

$$|s_n - l| = \left| \frac{2n+3}{3n+7} - \frac{2}{3} \right| = \frac{5}{9n+21} < \epsilon$$



### Example 8.0.6

Prove that  $\lim_{n \rightarrow \infty} \frac{2n+3}{3n-7} = \frac{2}{3}$ .

**Intuition:** Here, we have to be careful about throwing away terms.

$$\begin{aligned} |s_n - l| &< \epsilon \\ \left| \frac{2n+3}{3n-7} - \frac{2}{3} \right| &< \epsilon \\ \left| \frac{6n+9-6n+14}{9n-21} \right| &< \epsilon \\ \frac{23}{|9n-21|} &< \epsilon \end{aligned}$$

We want  $9n - 21 > 0$ , so we must have  $n \geq 3$ . We can apply this restriction on  $n$  to get rid of the absolute value:

$$\begin{aligned} \frac{23}{9n-21} &< \epsilon \\ \frac{23}{\epsilon} &< 9n-21 \\ \frac{1}{9} \left( \frac{23}{\epsilon} + 21 \right) &< n \end{aligned}$$

Thus, we want to choose some  $N > \frac{1}{9} \left( \frac{23}{\epsilon} + 21 \right)$  and  $N \geq 3$ .

*Proof.* Let  $\epsilon > 0$ . Let  $N \in \mathbb{N}$  such that  $N > \frac{1}{9} \left( \frac{23}{\epsilon} + 21 \right)$ . Then  $N > \frac{21}{9}$ , and since  $N$  is a natural number, then  $N \geq 3$ . Let  $n \in \mathbb{N}$  where  $n > N$ . Then:

$$\begin{aligned} 9n &> \frac{23}{\epsilon} + 21 \\ 9n - 21 &> \frac{23}{\epsilon} \\ \epsilon &> \frac{23}{9n-21} \end{aligned}$$

Thus:

$$|s_n - l| = \left| \frac{2n+3}{3n-7} - \frac{2}{3} \right| = \left| \frac{23}{9n-21} \right| = \frac{23}{9n-21} < \epsilon$$



**Definition 8.0.7 ▶ Divergence**

A sequence **diverges** if it does not converge.

$$\exists(\epsilon > 0)\forall(N \in \mathbb{N})\exists(n > N)(|s_n - l| \geq \epsilon)$$

**Example 8.0.8 ▶ Diverging Sequence**

Prove that  $s = (1, 0, 1, 0, 0, 1, 0, 0, 0, \dots)$  does not converge to 0.

*Proof.* Let  $\epsilon = 1/2$ . Then for all  $N \in \mathbb{N}$ , there exists  $n > N$  such that  $s_n = 1$ . Then:

$$|s_n - 0| = |1 - 0| > \epsilon$$

Therefore,  $s$  does not converge. □

## 8.1 Properties of Limits

A sequence can only converge to one value, not more. That is, if a sequence has a limit, then that limit is unique.

**Lemma 8.1.1 ▶ Approximating Zero**

Let  $x \in \mathbb{R}$ . If  $x < \epsilon$  for all  $\epsilon > 0$ , then  $x \leq 0$ .

*Proof.* We proceed by contraposition. Suppose  $x > 0$ . Let  $\epsilon := x/2 > 0$ . Then  $x \geq \epsilon = x/2$ . □

**Theorem 8.1.2 ▶ Uniqueness of Limits**

Let  $s_n$  be a sequence of real numbers. If  $s_n$  converges to  $l_1$  and converges to  $l_2$ , then  $l_1 = l_2$ .

*Proof.* Let  $\epsilon > 0$ . Since  $s_n$  converges to  $l_1$ , then there exists  $N_1 \in \mathbb{N}$  such that  $|s_n - l_1| < \epsilon/2$  for all  $n > N_1$ . Similarly, since  $s_n$  converges to  $l_2$ , then there exists  $N_2 \in \mathbb{N}$  such that  $|s_n - l_2| < \epsilon/2$  for all  $n > N_2$ .

Let  $n \in \mathbb{N}$  where  $n > N_1$  and  $n > N_2$ . Then:

$$|l_1 - l_2| = |l_1 - s_n + s_n - l_2| \leq \underbrace{|l_1 - s_n| + |s_n - l_2|}_{\text{Triangle Inequality}} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

Hence,  $|l_1 - l_2| < \epsilon$  for all  $\epsilon > 0$ . Thus, by Lemma 8.1.1,  $|l_1 - l_2| \leq 0$ . However, we know that  $|l_1 - l_2| \geq 0$  since it's an absolute value. Thus, we have  $|l_1 - l_2| = 0$ , so  $l_1 = l_2$ .  $\square$

Definitions of bounds for sequences, show that convergent implies boundedness

Split the theorem below into four separate theorems?

### Theorem 8.1.3 ► Properties of Limits

Let  $(s_n)$  and  $(t_n)$  be convergent sequences of real numbers, and let  $s, t \in \mathbb{R}$  such that  $s_n$  converges to  $s$  and  $t_n$  converges to  $t$ . Then:

1. For any  $c \in \mathbb{R}$ ,  $cs_n$  converges to  $cs$ ,
2.  $s_n + t_n$  converges to  $s + t$ ,
3.  $s_nt_n$  converges to  $st$ , and
4. if  $t_n \neq 0$ , then for all  $n$  and  $t \neq 0$ ,  $\frac{s_n}{t_n}$  converges to  $\frac{s}{t}$ .

*Proof of 1.* Let  $\epsilon > 0$ . Since  $(s_n)$  converges to  $s$ , then there exists  $N \in \mathbb{N}$  such that  $|s_n - s| < \frac{\epsilon}{1+|c|}$  for all  $n > N$ . Then, for all  $n > N$ , we have:

$$|cs_n - cs| = |c(s_n - s)| = |c||s_n - s| < |c|\frac{\epsilon}{1+|c|} = \frac{|c|}{1+|c|}\epsilon < \epsilon$$

$\square$

*Proof of 2.* Let  $\epsilon > 0$ . Since  $(s_n)$  converges to  $s$ , then there exists  $N_1 \in \mathbb{N}$  such that  $|s_n - s| < \epsilon/2$  for all  $n > N_1$ . Similarly, since  $t_n$  converges to  $t$ , then there exists  $N_2 \in \mathbb{N}$  such that  $|t_n - t| < \epsilon/2$ . Let  $N \in \mathbb{N}$  where  $N \geq N_1$  and  $N \geq N_2$ . Then:

$$|(s_n + t_n) - (s + t)| = |s_n - s + t_n - t| \leq |s_n - s| + |t_n - t| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

That is,  $s_n + t_n$  converges to  $s + t$ .  $\square$

*Proof of 3.* Let  $\epsilon > 0$ . Since  $s_n$  converges to  $s$ , then there exists  $N_1 \in \mathbb{N}$  such that  $|s_n - s| < \epsilon/2(|t| + 1)$  for all  $n > N_1$ . Also,  $(s_n)$  converges, so  $(s_n)$  is bounded. That is, there exists  $M \in \mathbb{R}$  such that  $|s_n| \leq M$  for all  $n \in \mathbb{N}$ . Since  $t_n$  converges to  $t$ , there exists  $N_2 \in \mathbb{N}$  such that

$|t_n - t| < \frac{\epsilon}{2(M+1)}$  for all  $n > N$ . Let  $N \in \mathbb{N}$  such that  $N \geq N_1$  and  $N \geq N_2$ . If  $n > N$ , then:

$$\begin{aligned}
 |s_n t_n - st| &= |s_n t_n - s_n t + s_n t - st| \\
 &= |s_n(t_n - t) + (s_n - s)t| \\
 &\leq |s_n(t_n - t)| + |(s_n - s)t| \\
 &= |s_n||t_n - t| + |s_n - s||t| \\
 &< M \frac{\epsilon}{2(1+M)} + \frac{\epsilon}{2(1+|t|)}|t| \\
 &= \frac{M}{1+M} \frac{\epsilon}{2} + \frac{\epsilon}{2} \frac{|t|}{1+|t|} \\
 &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\
 &= \epsilon
 \end{aligned}$$

□

*Proof of 4.* We will prove that  $\frac{1}{t_n}$  converges to  $\frac{1}{t}$ . Let  $\epsilon > 0$ . Since  $t_n$  converges to  $t$  and  $t \neq 0$ , then there exists  $N_1 \in \mathbb{N}$  such that  $|t_n - t| < \frac{\epsilon t^2}{2}$ . By ??, there exists  $N_2 \in \mathbb{N}$  such that  $|t_n| > \frac{|t|}{2}$  for all  $n > N_2$ . Let  $N \in \mathbb{N}$  such that  $N > N_1$  and  $N > N_2$ . Let  $n \in \mathbb{N}$  be arbitrary. Then:

$$\begin{aligned}
 \left| \frac{1}{t_n} - \frac{1}{t} \right| &= \left| \frac{t - t_n}{t_n t} \right| \\
 &= \frac{1}{|t_n|} \frac{1}{|t|} |t - t_n| \\
 &< \frac{2}{|t|} \frac{1}{|t|} \frac{\epsilon t^2}{2} \\
 &= \epsilon
 \end{aligned}$$

By 3, if  $s_n$  converges to  $s$ , then  $\frac{s_n}{t_n} = s_n \left( \frac{1}{t_n} \right)$  converges to  $s \left( \frac{1}{t} \right) = \frac{s}{t}$ .

□

#### Lemma 8.1.4 ▶ Limit of a Constant Sequence

If  $s_n$  is a constant sequence  $(l, l, l, \dots)$ , then  $s_n$  converges to  $l$ .

*Proof.* Let  $\epsilon > 0$ . For all  $n \in \mathbb{N}$ ,  $|s_n - l| = 0 < \epsilon$ .

□

lemma  
14.6 for  
bounding  
in proof  
of 4.

Explain  
new no-  
tation

better



**Example 8.1.5 ▶ Using the Properties**

Prove  $\lim_{n \rightarrow \infty} \frac{5n^3 - 8n^2 + 15}{7n^3 + 19n + 4} = \frac{5}{7}$ .

*Proof.*

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{5n^3 - 8n^2 + 15}{7n^3 + 19n + 4} &= \lim_{n \rightarrow \infty} \frac{5 - 8/n + 15/n^3}{7 + 19/n^2 + 4/n^3} \\ &= \frac{\lim_{n \rightarrow \infty} 5 - 8/n + 15/n^3}{\lim_{n \rightarrow \infty} 7 + 19/n^2 + 4/n^3} \\ &= \frac{\lim_{n \rightarrow \infty} 5 - \lim_{n \rightarrow \infty} 8/n + \lim_{n \rightarrow \infty} 15/n^3}{\lim_{n \rightarrow \infty} 7 + \lim_{n \rightarrow \infty} 19/n^2 + \lim_{n \rightarrow \infty} 4/n^3} \end{aligned}$$

Now we can work with each limit independently. Note that  $\lim_{n \rightarrow \infty} \frac{1}{n^2} = \left(\lim_{n \rightarrow \infty} \frac{1}{n}\right)\left(\lim_{n \rightarrow \infty} \frac{1}{n}\right)$ , so:

$$\lim_{n \rightarrow \infty} \frac{5n^3 - 8n^2 + 15}{7n^3 + 19n + 4} = \frac{5}{7}$$

□

**Definition 8.1.6 ▶ Increasing, Decreasing, Monotonic**

A sequence  $(s_n)$  is:

- **increasing** if  $s_n \leq s_{n+1}$  for all  $n \in \mathbb{N}$ .
- **strictly increasing** if  $s_n < s_{n+1}$  for all  $n \in \mathbb{N}$ .
- **decreasing** if  $s_n \geq s_{n+1}$  for all  $n \in \mathbb{N}$ .
- **strictly decreasing** if  $s_n > s_{n+1}$  for all  $n \in \mathbb{N}$ .

If  $(s_n)$  satisfies any of these properties, then we say  $(s_n)$  is **monotonic**.

For example,  $(s_n) = \left(\frac{1}{n}\right) = \left(1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\right)$  is strictly decreasing and thus monotonic.

**Theorem 8.1.7 ▶ Monotone Sequence Theorem**

Let  $(s_n)$  be a sequence of real numbers.

1. If  $(s_n)$  is increasing and bounded above, then  $(s_n)$  converges to  $\sup\{s_n : n \in \mathbb{N}\}$ .
2. If  $(s_n)$  is decreasing and bounded below, then  $(s_n)$  converges to  $\inf\{s_n : n \in \mathbb{N}\}$ .

**Idea:** Assuming  $s$  is our limit, we want to find  $N \in \mathbb{N}$  such that  $|s - s_n| < \epsilon$ , or  $s - \epsilon < s_n$  for all  $n > N$ . Then  $s - \epsilon < s_n \leq s$  for all  $n > N$ .

*Proof of 1.* Let  $\epsilon > 0$ . Because  $\{s_n : n \in \mathbb{N}\}$  is non-empty and bounded above, then it has a supremum. Let  $s := \sup\{s_n : n \in \mathbb{N}\}$ . Thus, there exists  $N \in \mathbb{N}$  such that  $s_N > s - \epsilon$  (by the approximation property). Since  $(s_n)$  is increasing, we have:

$$\forall (n > N) (s - \epsilon < s_N \leq s_n \leq s)$$

Hence,  $\epsilon < s_n - s \leq 0$ , so  $|s_n - s| < \epsilon$ . □

*Proof of 2.* Suppose  $(s_n)$  is decreasing and bounded below. Then  $s_{n+1} \leq s_n$  for all  $n \in \mathbb{N}$ . Moreover, there exists  $m \in \mathbb{R}$  such that  $s_n \geq m$  for all  $n \in \mathbb{N}$ . That is,  $-s_{n+1} \geq -s_n$  for all  $n \in \mathbb{N}$ , and  $-s_n \leq -m$  for all  $n \in \mathbb{N}$ . Therefore,  $(-s_n)$  is increasing and bounded above. By the first part, we know  $(-s_n)$  converges to  $\sup\{-s_n : n \in \mathbb{N}\} = -\inf\{s_n : n \in \mathbb{N}\}$ . Hence,  $(s_n)$  converges to  $\inf\{s_n : n \in \mathbb{N}\}$ . □

## 8.2 Subsequences

So far, we've only looked at well-behaving sequences that converge. What about sequences that don't converge? Can we still find some nice properties that describe their behavior?

$$(s_n) = (0, 1, 0, 1, 0, 1, \dots)$$

Consider a sequence  $(t_n)$  where  $t_n = s_{2n}$ . That is:

$$(t_n) = (s_2, s_4, s_6, \dots) = (1, 1, 1, \dots)$$

Inside this diverging sequence, we can find a convergent **subsequence**! Intuitively, we can make a subsequence by “throwing away” terms but keeping the same order. We can formally define a subsequence as follows:

### Definition 8.2.1 ▶ Subsequence

Given a sequence  $(s_n)$ , a **subsequence** is any sequence of the form  $(t_k)_{k \in \mathbb{N}}$  where  $t_k = s_{n_k}$  for all  $k \in \mathbb{N}$ ,  $n_k \in \mathbb{N}$  for all  $k \in \mathbb{N}$ , and  $n_k < n_{k+1}$  for all  $k \in \mathbb{N}$ .

For example, if we had  $s = (s_1, s_2, s_3, s_4, s_5, s_6, s_7, \dots, s_{213}, s_{214}, s_{215}, \dots)$ , we can have a subse-

quence like:

$$(t_n) = (s_3, s_5, s_{213}, \dots)$$

Here, we would have  $n_1 = 3$ ,  $n_2 = 5$ ,  $n_3 = 213$ , and so on.

### Example 8.2.2 ▶ Subsequences

Let  $(s_n) := (1, 1/2, 1/3, 1/4, 1/5, 1/6, \dots)$

1.  $(t_n) := (1, 1/4, 1/9, 1/16, 1/25, \dots)$  is a subsequence of  $(s_n)$  where  $t_n = 1/n^2$ , or  $t_n = s_{n^2}$ .
2.  $(t_n) := (1/5, 1/25, 1/125, \dots)$  is also a subsequence of  $(s_n)$  with  $t_n = \frac{1}{5^n}$  or  $t_n = s_{5^n}$ .
3.  $(t_n) := (1/7, 1/2, 1/12, 1/6)$  is **not** a subsequence of  $(s_n)$  because the indices in  $s_n$  are not strictly increasing. We have  $n_1 = 7$ , but  $n_2 = 2$ .

**In general:**

$$(s_n) = (s_1, s_2, s_3, \dots)$$

$$(t_n) = (s_{n_k}) = (s_{n_1}, s_{n_2}, s_{n_3})$$

### Lemma 8.2.3 ▶ Indices of Subsequences

If  $(s_{n_k})_{k \in \mathbb{N}}$  is a subsequence of  $(s_n)_{n \in \mathbb{N}}$ , then  $n_k \geq k$  for all  $k \in \mathbb{N}$ .

We will use induction.

**Base Case:** Since  $n_1 \in \mathbb{N}$ , then  $n_1 \geq 1$ .

**Induction Step:** Suppose  $n_k \geq k$  for some  $k \in \mathbb{N}$ . Since  $n_{k+1} > n_k$ , we have  $n_{k+1} \geq n_k + 1 \geq k + 1$ .

Hence,  $n_k \geq k$  for all  $k \in \mathbb{N}$ .

### Theorem 8.2.4 ▶ Limits of Subsequences

Suppose  $(s_n)$  is a sequence of real numbers, and  $s_n$  converges to  $s$  for some  $s \in \mathbb{R}$ . If  $(s_{n_k})$  is a subsequence of  $(s_n)$ , then  $s_{n_k}$  converges to  $s$ .

*Proof.* Let  $\epsilon > 0$ . Since  $s_n$  converges to  $s$ , then there exists  $N \in \mathbb{N}$  such that  $|s_n - s| < \epsilon$  for all  $n > N$ . Suppose  $k > N$ . By lemma 8.2.3,  $n_k \geq k > N$ , so  $|s_{n_k} - s| < \epsilon$ .  $\square$

## 8.3 Limit Superior and Inferior

Suppose  $(s_n)$  is a bounded sequence. Then there exists  $M \in \mathbb{R}$  such that  $-M \leq s_n \leq M$  for all  $n \in \mathbb{N}$ . Let:

$$\begin{aligned} t_1 &:= \sup\{s_1, s_2, s_3, \dots\} = \sup\{s_k : k \geq 1\} \\ t_2 &:= \sup\{s_2, s_3, s_4, \dots\} = \sup\{s_k : k \geq 2\} \\ t_3 &:= \sup\{s_3, s_4, s_5, \dots\} = \sup\{s_k : k \geq 3\} \\ &\vdots \\ t_n &:= \sup\{s_n, s_{n+1}, s_{n+2}, \dots\} = \sup\{s_k : k \geq n\} \\ t_{n+1} &:= \sup\{s_{n+1}, s_{n+2}, s_{n+3}, \dots\} = \sup\{s_k : k \geq n+1\} \end{aligned}$$

Then:

$$-M \leq s_n \leq t_n$$

and:

$$t_{n+1} \leq t_n$$

so  $(t_n)$  is bounded below and decreasing. Hence,  $(t_n)$  converges by the Monotone Sequence Theorem.

### Definition 8.3.1 ► Limit Superior, Limit Inferior

Let  $(s_n)$  be a bounded sequence of real numbers. The **limit superior** is defined as:

$$\limsup s_n := \lim_{n \rightarrow \infty} \sup\{s_k : k \geq n\}$$

Similarly, the **limit inferior** is defined as:

$$\liminf s_n := \lim_{n \rightarrow \infty} \inf\{s_k : k \geq n\}$$

### Example 8.3.2

$$\text{Define } s_n := \begin{cases} 3 + \frac{1}{n}, & n \text{ is even} \\ 1 - \frac{1}{n}, & n \text{ is odd} \end{cases}$$

$$(s_n) = (0, 3 + 1/2, 2/3, 3 + 1/4, 4/5, 3 + 1/6)$$

Let's try to calculate the limit superior of  $s_n$ . Define  $(t_n)$  as follows:

$$\begin{aligned} t_1 &:= \sup\{s_1, s_2, s_3, \dots\} = 3 + \frac{1}{2} \\ t_2 &:= \sup\{s_2, s_3, s_4, \dots\} = 3 + \frac{1}{2} \\ t_3 &:= \sup\{s_3, s_4, s_5, \dots\} = 3 + 1/4 \\ t_4 &:= \sup\{s_4, s_5, s_6, \dots\} = 3 + 1/4 \\ t_5 &:= \sup\{s_5, s_6, s_7, \dots\} = 3 + 1/6 \\ &\vdots \end{aligned}$$

We can see that  $\limsup s_n = \lim_{n \rightarrow \infty} \sup\{s_k : k \geq n\} = 3$ . We might refer to 3 as the “largest limit point”.

Now let's try to calculate the limit inferior of  $s_n$ . Define  $(r_n)$  as follows:

$$\begin{aligned} r_1 &:= \inf\{s_1, s_2, s_3, \dots\} = 0 \\ r_2 &:= \inf\{s_2, s_3, s_4, \dots\} = \frac{2}{3} \\ r_3 &:= \inf\{s_3, s_4, s_5, \dots\} = \frac{2}{3} \\ r_4 &:= \inf\{s_4, s_5, s_6, \dots\} = \frac{4}{5} \\ &\vdots \end{aligned}$$

We can see that  $\liminf s_n = \lim_{n \rightarrow \infty} \inf\{s_k : k \geq n\} = 1$ . We might refer to 1 as the “smallest limit point”.

### Theorem 8.3.3

Suppose  $(s_n)$  is a bounded sequence of real numbers, and suppose that  $(s_{n_k})$  is a convergent subsequence of  $(s_n)$ . Then  $\liminf s_n \leq \lim_{k \rightarrow \infty} s_{n_k} \leq \limsup s_n$ .

*Proof.* Let  $r_n := \inf\{s_k : k \geq n\}$  and  $t_n := \sup\{s_k : k \geq n\}$ . Then  $r_n \leq s_n \leq t_n$  for all  $n \in \mathbb{N}$ . In particular,  $r_{n_k} \leq s_{n_k} \leq t_{n_k}$  for all  $k \in \mathbb{N}$ . By (todo: theroem),  $\lim_{k \rightarrow \infty} r_{n_k} = \lim_{n \rightarrow \infty} r_n$ . Note that  $\lim_{n \rightarrow \infty} r_n = \liminf s_n$ , and  $\lim_{k \rightarrow \infty} t_{n_k} = \lim_{n \rightarrow \infty} t_n = \limsup s_n$ . By the

(todo: problem set squeeze theorem), we have:

$$\liminf s_n = \lim_{k \rightarrow \infty} r_{n_k} \leq \lim_{k \rightarrow \infty} s_{n_k} \leq \lim_{k \rightarrow \infty} t_{n_k} = \limsup s_n$$

□

### Theorem 8.3.4 ► Bolzano-Weierstrass Theorem

Suppose  $(s_n)$  is a bounded sequence of real numbers. The  $(s_n)$  has a subsequence that converges to  $\limsup s_n$ , and  $(s_n)$  has a subsequence that converges to  $\liminf s_n$ .

#### Intuition:

- Let  $t_k := \sup\{s_k, s_{k+1}, s_{k+2}, \dots\}$ , so  $\limsup s_n = \lim_{k \rightarrow \infty} t_k$ .
- For each  $k \in \mathbb{N}$  we can find some  $n_k \geq k$  such that  $t_k - 1/k < s_{n_k}$ .
- Thus,  $-1/k < s_{n_k} - t_k \leq 0$ , so  $|s_{n_k} - t_k| < 1/k$ .
- By (todo: problem set),  $s_{n_k} - t_k \rightarrow 0$ , so  $s_{n_k} = s_{n_k} - t_k + t_k \rightarrow \limsup s_n$ .
- But: we need  $n_1 < n_2 < \dots < n_k < n_{k+1} < \dots$ . So we need to choose  $n_k$  inductively!

*Proof for limsup.* We will choose a subsequence of  $(s_n)$  that converges to  $\limsup s_n$ . For each  $k \in \mathbb{N}$ , let  $t_k := \sup\{s_k, s_{k+1}, s_{k+2}, \dots\}$ . For convenience, let  $P(n)$  be the statement “there exists  $n_k \in \mathbb{N}$  such that  $n_k > n_{k-1}$  and  $|s_{n_k} - t_{1+n_{k-1}}| < \frac{1}{k}$ .” We define  $n_0 := 0$ .

**Base Case:** Let  $t_1 := \sup\{s_1, s_2, \dots\}$ . By the approximation property (todo ref), there exists  $n_1 \in \mathbb{N}$  such that  $t_1 - 1 < s_{n_1} \leq t_1$ . Subtracting across by  $t_1$ , we have  $-1 < s_{n_1} - t_1 \leq 0$ . Thus,  $|s_{n_1} - t_1| < 1$ .

**Induction Step:** Now we aim to prove  $P(k-1) \implies P(k)$ . There exists  $n_k \in \mathbb{N}$  such that  $n_k > n_{k-1}$ , and:

$$\begin{aligned} & t_{1+n_{k-1}} - \frac{1}{k} < s_{n_k} \leq t_{1+n_{k-1}} \\ \implies & -\frac{1}{k} < s_{n_k} - t_{1+n_{k-1}} \leq 0 \\ \implies & |s_{n_k} - t_{1+n_{k-1}}| < \frac{1}{k} \end{aligned}$$

That is,  $\lim_{k \rightarrow \infty} (s_{n_k} - t_{1+n_{k-1}}) = 0$ . Since  $n_k > n_{k-1}$  for all  $k \in \mathbb{N}$ ,  $(s_{n_k})$  is a subsequence

of  $(s_n)$ . But  $(t_{1+n_{k-1}})$  is a subsequence of  $(t_k)$ , so:

$$\lim_{k \rightarrow \infty} t_{1+n_{k-1}} = \lim_{k \rightarrow \infty} t_k = \limsup s_n$$

Thus:

$$s_{n_k} = s_{n_k} - t_{1+n_{k-1}} + t_{1+n_{k-1}}$$

so:

$$\lim_{k \rightarrow \infty} s_{n_k} = \lim_{k \rightarrow \infty} (s_{n_k} - t_{1+n_{k-1}}) + \lim_{k \rightarrow \infty} t_{1+n_{k-1}} = 0 + \limsup s_n$$

Therefore,  $(s_{n_k})$  is a subsequence of  $(s_n)$  that converges to  $\limsup s_n$ . □

### Theorem 8.3.5 ▶ Convergence iff $\limsup = \liminf$

Let  $(s_n)$  be a bounded sequence of real numbers. Then  $(s_n)$  converges if and only if  $\liminf s_n = \limsup s_n$

*Proof.* First, suppose  $s_n$  converges to some  $s \in \mathbb{R}$ . By the Bolzano-Weierstrass Theorem, there exists a subsequence  $(s_{n_k})$  of  $(s_n)$  such that  $\lim_{k \rightarrow \infty} s_{n_k} = \limsup s_n$ . But  $s_n$  converges to  $s$ , so  $s_{n_k}$  also converges to  $s$ . That is,  $s = \lim_{k \rightarrow \infty} s_{n_k} = \limsup s_n$ . By the same reasoning, we have  $s = \liminf s_n$ . Hence,  $\liminf s_n = \lim s_n = \limsup s_n$ .

Conversely, suppose  $\liminf s_n = \limsup s_n$ . Let  $r_n := \inf\{s_k : k \geq n\}$  and  $t_n := \sup\{s_k : k \geq n\}$ . Then  $r_n \leq s_n \leq t_n$  for all  $n \in \mathbb{N}$ . Then  $\lim r_n = \liminf s_n = \limsup s_n = \lim t_n$ . Therefore, by the Squeeze Theorem (todo: ref),  $s_n$  converges to  $\liminf s_n$ . □

## 8.4 Cauchy Sequences

To show that a sequence  $(s_n)$  converges using the definition of limit, we need to know what limit is beforehand. Consider the following limit:

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{k^3}$$

This sequence of partial sums converges, but its limit is unknown. Certainly we can get a decimal approximation for this value, but there is no known “closed” form of this value.

**Definition 8.4.1 ▶ Cauchy Sequence**

We say a sequence is **Cauchy** if, for all  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $|s_n - s_m| < \epsilon$  for all  $n > N$  and  $m > N$ .

$$\forall(\epsilon > 0) \exists(N \in \mathbb{N}) \forall(n > N, m > N) (|s_n - s_m| < \epsilon)$$

In other words, a sequence is Cauchy if **all** terms in the tail can be made arbitrarily close to each other. Or, for any arbitrarily small distance, there exists some “tail” of the sequence that exists entirely within that distance. This definition circumvents any mention of a specific “limit”. But we can prove that any Cauchy sequence of real numbers is convergent, and vice versa.

**Lemma 8.4.2 ▶ Convergent Sequences are Cauchy**

If a sequence of real numbers converges, then that sequence is Cauchy.

*Proof.* Let  $(s_n)$  be a convergent sequence of real numbers. Let  $\epsilon > 0$ . Since  $(s_n)$  converges to some  $s \in \mathbb{R}$ , there exists  $N \in \mathbb{N}$  such that  $|s_n - s| < \epsilon/2$  for all  $n > N$ . If  $n > N$  and  $m > N$ , then:

$$|s_n - s_m| = |s_n - s + s - s_m| \leq |s_n - s| + |s - s_m| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

□

**Lemma 8.4.3**

Suppose  $(s_n)$  is a Cauchy sequence, and that  $(s_{n_k})$  is a convergent subsequence of  $(s_n)$  where  $s_{n_k}$  converges to some  $s \in \mathbb{R}$ . Then  $(s_n)$  converges, and  $\lim s_n = s$ .

*Proof.* Let  $\epsilon > 0$ . Since  $(s_n)$  is Cauchy, then there exists some  $N \in \mathbb{N}$  such that  $|s_n - s_m| < \epsilon/2$  for all  $n > N$  and  $m > N$ . Since  $(s_{n_k})$  converges to  $s$ , there exists  $N_1 \in \mathbb{N}$  such that  $|s_{n_k} - s| < \frac{\epsilon}{2}$  for all  $k > N_1$ . Let  $k \in \mathbb{N}$  where  $k > N$  and  $k > N_1$ . Since  $n_k \geq k$ , then  $n_k > N$  and  $n_k > N_1$ . For all  $n > k$ , we have:

$$|s_n - s| = |s_n - s_{n_k} + s_{n_k} - s| \leq \underbrace{|s_n - s_{n_k}|}_{n_1, n_k > N} + \underbrace{|s_{n_k} - s|}_{n_k > N_1} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

□



**Lemma 8.4.4 ▶ Cauchy Sequences are Bounded**

If a sequence of real numbers is Cauchy, then that sequence is bounded.

*Proof.* Let  $(s_n)$  be a Cauchy sequence of real numbers. Then there exists  $N \in \mathbb{N}$  such that  $|s_n - s_m| < 1$  for all  $n > N$  and  $m > N$ . Let  $m := N + 1$ . Then, for all  $n > N$ , we have:

$$|s_n| = |s_n - s_m + s_m| \leq |s_n - s_m| + |s_m| < 1 + |s_m| = 1 + |s_{N+1}|$$

Thus, for all  $n \in \mathbb{N}$ , we have:

$$|s_n| \leq \max\{|s_1|, |s_2|, \dots, |s_N|, 1 + |s_{N+1}|\}$$

Therefore,  $(s_n)$  is bounded. □

**Theorem 8.4.5 ▶ Cauchy Criterion**

A sequence of real numbers converges if and only if it is Cauchy.

*Proof.* By Lemma 8.4.2, we know that convergence implies Cauchy.

If  $(s_n)$  is Cauchy, then by Lemma 8.4.4,  $(s_n)$  is bounded. By the Bolzano-Weierstrass Theorem,  $(s_n)$  has a convergent subsequence. By Lemma 8.4.3,  $(s_n)$  converges. □

Our definition of completeness in  $\mathbb{R}$  predicates on a notion of order between the elements. Specifically, we said  $\mathbb{R}$  is complete because every subset of  $\mathbb{R}$  that is bounded above has a supremum. What does it mean to say  $\mathbb{R}^2$  is complete?

**Definition 8.4.6 ▶ Completeness (in terms of Cauchy Sequences)**

A (metric) space is **complete** if every Cauchy sequence converges to a point in the space.

The intuition is the same: there are no points “missing” from the space.

# Open and Closed Sets

We will describe some concepts that generalize open/closed intervals. This chapter also serves as a very light introduction to topology—specifically, the topology of the real number line.

## 9.1 Open Sets

### Definition 9.1.1 ► Open Set

Intuitively, set is **open** if it does not contain any of its “boundary points”, such as minimum or maximum.

More formally, we say  $A \subseteq \mathbb{R}$  is **open** if, for all  $x \in A$ , there exists  $r > 0$  such that  $(x - r, x + r) \subseteq A$ .

$$\forall(x \in A)\exists(r > 0)((x - r, x + r) \subseteq A)$$

### Example 9.1.2 ► $[0, 1)$ is not open

The interval  $[0, 1)$  is not open.

*Proof.*  $0 \in [0, 1)$ , but  $(0 - r, 0 + r) \not\subseteq [0, 1)$  for any  $r > 0$ . □

### Definition 9.1.3 ► Open Ball

We call the interval  $(x - r, x + r)$  the **open ball** of radius  $r$  centered at  $x$ , notated as  $B(x, r)$  or  $B_r(x)$ .

$$B(x, r) = B_r(x) = (x - r, x + r)$$

This new notation lets us write ideas more succinctly. For example,  $\mathbb{R}$  is open. Given any  $x \in \mathbb{R}$ , then any  $r > 0$  will give us  $B(x, r) \subseteq \mathbb{R}$ . Also,  $\emptyset$  is vacuously open.

**Lemma 9.1.4 ▶ Open Intervals are Open Sets**

Let  $a, b \in \mathbb{R}$  where  $a < b$ . Then  $(a, b)$  is an open set.

*Proof.* Let  $c := \frac{a+b}{2}$ , and let  $R := \frac{b-a}{2}$ . Then  $(a, b) = B(c, R)$ . Let  $x \in B(c, R)$ . Then  $|x - c| < R$ . Let  $r := R - |x - c| > 0$ . We now prove  $B(x, r) \subseteq B(c, R)$ . Let  $y \in B(x, r)$ . Then  $|x - y| < r$ , so:

$$|y - c| = |y - x + x - c| \leq |y - x| + |x - c| < r + |x - c| = R - |x - c| + |x - c| = R$$

Hence,  $y \in B(c, R) = (a, b)$ . Therefore,  $(a, b)$  is an open set.  $\square$

As we prove below, an arbitrary union of open sets is itself an open set.

**Theorem 9.1.5 ▶ Union of Open Sets is Open**

Suppose  $\Lambda$  is a set, and for each  $\lambda \in \Lambda$ ,  $O_\lambda$  is an open subset of  $\mathbb{R}$ . Then  $\bigcup_{\lambda \in \Lambda} O_\lambda$  is an open set.

*Proof.* Let  $x \in \bigcup_{\lambda \in \Lambda} O_\lambda$ . Then there exists some  $\lambda_0 \in \Lambda$  such that  $x \in O_{\lambda_0}$ . Since  $O_{\lambda_0}$  is open, there exists  $r > 0$  such that:

$$(x - r, x + r) \subseteq O_{\lambda_0} \subseteq \bigcup_{\lambda \in \Lambda} O_\lambda$$

$\square$

The intersection of open sets is more troublesome. Countable intersections of open sets may not be open. For example, let  $A_n := \left(-\frac{1}{n}, \frac{1}{n}\right)$  for each  $n \in \mathbb{N}$ . Then  $\bigcap_{n \in \mathbb{N}} A_n = \{0\}$  is not open!

**Theorem 9.1.6 ▶ Finite Intersection of Open Sets is Open**

Let  $n \in \mathbb{N}$ , and let  $O_1, O_2, \dots, O_n$  be open subsets of  $\mathbb{R}$ . Then  $\bigcap_{k=1}^n O_k$  is open.

*Proof.* Let  $x \in \bigcap_{k=1}^n O_k$ . Then  $x \in O_k$  for  $k = 1, 2, \dots, n$ . Then, for each  $k \in \{1, 2, \dots, n\}$ , there must be some radius  $r_k > 0$  such that  $B(x, r_k) \subseteq O_k$ . Since there are only finitely many open sets, we can take the minimum radius. Let  $r := \min\{r_1, r_2, \dots, r_n\}$ . Then,  $r \leq r_k$  for each  $k \in \{1, 2, \dots, n\}$ . Hence:

$$B(x, r) \subseteq B(x, r_k) \subseteq O_k \quad \text{for all } k \in \{1, 2, \dots, n\}$$

Therefore,  $B(x, r) \subseteq \bigcap_{k=1}^n O_k$ , so it is open. □

Note how the above theorem only works by taking the minimum radius of all the open sets. We can only take this minimum radius because there are only a finite number of open sets.

## 9.2 Closed Sets

### Definition 9.2.1 ► Closed Set

Intuitively, a set is **closed** if it contains all of its “boundary points”.

More formally, a set  $E \subseteq \mathbb{R}$  is **closed** if every convergent sequence  $(s_n)$  where  $s_n \in E$  for all  $n \in \mathbb{N}$  satisfies  $\lim_{n \rightarrow \infty} s_n \in E$ .

### Example 9.2.2 ► $(0, 1]$ is not closed

The interval  $[0, 1)$  is not closed.

*Proof.* Consider the sequence  $(s_n) := 1/n$ . Then  $(s_n)$  converges to 0, but  $0 \notin (0, 1]$ . □

Note that this interval  $(0, 1]$  is neither open nor closed! It is wrong to think of open/closed as strictly one or the other (i.e. openness and closedness are not mutually exclusive). Moreover, a set can be both open and closed (or **clopen**), going against the intuition of open and closed sets. There are only two clopen sets in the real numbers:  $\mathbb{R}$  itself, and  $\emptyset$ .

### Lemma 9.2.3 ► Closed Intervals are Closed Sets

Let  $a, b \in \mathbb{R}$  with  $a < b$ . Then  $[a, b]$  is a closed set.

*Proof.* Let  $(s_n)$  be an arbitrary convergent sequence of real numbers where  $a \leq s_n \leq b$  for all  $n \in \mathbb{N}$ . Since  $(s_n)$  is convergent, then  $\lim_{n \rightarrow \infty} s_n$  exists. By the properties of limits, we have:

$$\lim_{n \rightarrow \infty} a \leq \lim_{n \rightarrow \infty} s_n \leq \lim_{n \rightarrow \infty} b$$

Hence,  $\lim_{n \rightarrow \infty} s_n \in [a, b]$ . Therefore,  $[a, b]$  is a closed set. □

**Theorem 9.2.4 ▶ Intersection of Closed Sets is Closed**

Let  $\Lambda$  be a set, and let  $E_\lambda \subseteq \mathbb{R}$  be closed for all  $\lambda \in \Lambda$ . Then  $\bigcap_{\lambda \in \Lambda} E_\lambda$  is a closed set.

*Proof.* Let  $(s_n)$  be an arbitrary convergent sequence of real numbers entirely contained within  $\bigcap_{\lambda \in \Lambda} E_\lambda$ . Since  $(s_n)$  is convergent, then  $\lim_{n \rightarrow \infty} s_n$  exists. Let  $l$  denote that limit. Let  $\lambda \in \Lambda$  be arbitrary. Then  $s_n \in E_\lambda$  for all  $n \in \mathbb{N}$ . Since  $E_\lambda$  is closed, then any convergent sequence contained in  $E_\lambda$  has its limit in  $E_\lambda$ . Thus,  $\lim_{n \rightarrow \infty} s_n \in E_\lambda$ . Since  $\lambda \in \Lambda$  is arbitrary, then  $\lim_{n \rightarrow \infty} s_n \in E_\lambda$  for all  $\lambda \in \Lambda$ . Therefore,  $s \in \bigcap_{\lambda \in \Lambda} E_\lambda$ , so this set is closed.  $\square$

Similar to the intersection of open sets, the union of closed sets is guaranteed to be closed if it is a finite union. For example, the union  $\left(\bigcup_{n \in \mathbb{N}} [1/n, 1]\right) = (0, 1]$  is not closed!

**Theorem 9.2.5 ▶ Finite Union of Closed Sets is Closed**

Let  $n \in \mathbb{N}$ , and let  $E_1, E_2, \dots, E_n$  be closed subsets of  $\mathbb{R}$ . Then  $\bigcup_{k=1}^n E_k$  is a closed set.

A direct proof of this theorem can be found in the textbook.

The direct proof here is rather wordy and awkward. We will first establish a concrete relationship between open and closed sets, then leverage that to prove this theorem “indirectly”.

**Theorem 9.2.6 ▶ Complement of an Open Set is Closed**

Let  $O \subseteq \mathbb{R}$  be open. Then  $\mathbb{R} \setminus O$  is closed.

*Proof.* Let  $(x_n)$  be an arbitrary convergent sequence entirely contained within  $\mathbb{R} \setminus O$ . Let  $l_x := \lim_{n \rightarrow \infty} x_n$ . Suppose for contradiction that  $l_x \notin \mathbb{R} \setminus O$ . Then  $l_x \in O$ . Since  $O$  is open, there exists some radius  $r > 0$  such that  $B(l_x, r) \subseteq O$ . Since  $(x_n)$  converges to  $l_x$ , then there exists  $N \in \mathbb{N}$  such that  $|x_n - l_x| < r$  for all  $n > N$ . That is,  $x_n \in B(l_x, r) \subseteq O$  for all  $n > N$ . This contradicts  $x_n \in \mathbb{R} \setminus O$ . Thus,  $l_x \in \mathbb{R} \setminus O$ , so  $\mathbb{R} \setminus O$  is closed.  $\square$

**Theorem 9.2.7 ▶ Complement of a Closed Set is Open**

Let  $C \subseteq \mathbb{R}$  be closed. Then  $\mathbb{R} \setminus C$  is open.

*Proof.* Let  $x \in \mathbb{R} \setminus C$ . We must prove the following statement:

$$\exists (n \in \mathbb{N}) (B(x, 1/n) \subseteq \mathbb{R} \setminus C)$$

Suppose for contradiction the negation of the previous statement holds. That is:

$$\forall (n \in \mathbb{N}) (B(x, 1/n) \not\subseteq \mathbb{R} \setminus C)$$

In other words, for all  $n \in \mathbb{N}$ , there exists  $x_n \in B(x, 1/n)$  such that  $x_n \in C$ . Hence, the sequence  $(x_n)$  satisfies  $x_n \in C$  for all  $n \in \mathbb{N}$  and  $|x_n - x| < 1/n$ . Thus,  $(x_n)$  converges to  $x$ . However,  $C$  is closed, and  $(x_n)$  is a sequence in  $C$ , so  $x \in C$ . This contradicts  $x \in \mathbb{R} \setminus C$ . Therefore, our original statement holds, so  $\mathbb{R} \setminus C$  is open.  $\square$

Combining the two above theorems, we can infer a pretty useful relationship between open and closed sets.

$$A \text{ is open} \iff \mathbb{R} \setminus A \text{ is closed}$$

$$B \text{ is closed} \iff \mathbb{R} \setminus B \text{ is open}$$

We can apply this relationship to directly prove that the finite union of closed sets is closed (Theorem 9.2.5).

*Proof of Theorem 9.2.5.* By De Morgan's Laws, we have:

$$\mathbb{R} \setminus \left( \bigcup_{k=1}^n E_k \right) = \bigcap_{k=1}^n (\mathbb{R} \setminus E_k)$$

Since each  $\mathbb{R} \setminus E_k$  is open, the finite intersection  $\bigcap_{k=1}^n (\mathbb{R} \setminus E_k)$  is also open. Hence,  $\mathbb{R} \setminus \left( \bigcup_{k=1}^n E_k \right)$  is open, so  $\bigcup_{k=1}^n E_k$  is closed.  $\square$

## 9.3 Closure

### Definition 9.3.1 ► Closure of a Set

Intuitively, the **closure** of a set is the set containing its elements and boundary points.

Formally, for  $A \subseteq \mathbb{R}$ , the **closure** of  $A$  is the set:

$$\overline{A} := \{x \in \mathbb{R} : \exists (\text{sequence } (x_n)) \forall (n \in \mathbb{N}) (x_n \in A) \text{ and } x_n \rightarrow x\}$$

For example, let's consider the closure the open interval  $A := (0, 1)$ .

- We can take the constant sequence of  $1/2$  contained in  $(0, 1)$  which converges to  $1/2$ , so  $1/2 \in \bar{A}$ .
- We can take the sequence  $1/n + 1$  contained in  $(0, 1)$  which converges to 0, so  $0 \in \bar{A}$ .
- We can take the sequence  $1 - 1/n + 1$  contained in  $(0, 1)$  which converges to 1, so  $1 \in \bar{A}$ .

### Theorem 9.3.2 ► Properties of Closures of Sets

Let  $A \subseteq \mathbb{R}$ . Then:

- (i)  $A \subseteq \bar{A}$ ,
- (ii)  $\bar{A}$  is closed,
- (iii)  $A = \bar{A}$  if and only if  $A$  is closed,
- (iv)  $\bar{A} = \overline{\bar{A}}$ ,
- (v) if  $F \subseteq \mathbb{R}$  is closed and  $A \subseteq F$ , then  $\bar{A} \subseteq F$ , and
- (vi)  $\bar{A} = \bigcap \{F \subseteq \mathbb{R} : F \text{ is closed, and } A \subseteq F\}$

*Proof.* The proofs for i through vi are as follows:

- (i) Let  $x \in A$ , and let  $(x_n) := (x, x, x, \dots)$ . Then  $x_n \in A$  for all  $n \in \mathbb{N}$ , and  $\lim_{n \rightarrow \infty} x_n = x$ . Therefore,  $x \in \bar{A}$ .
- (ii) Let  $(x_n)$  be a sequence contained within  $\bar{A}$  that converges to some  $x \in \mathbb{R}$ . For each  $x_n \in \bar{A}$ , there exists  $y_n \in A$  such that  $|y_n - x_n| < 1/n$ . Then:

$$|y_n - x| = |y_n - x_n + x_n - x| \leq |y_n - x_n| + |x_n - x| < 1/n + |x_n - x|$$

Thus,  $y_n - x \rightarrow 0$ , so  $y_n \rightarrow x$ . Therefore,  $x \in \bar{A}$ .

- (iii) First, suppose  $A = \bar{A}$ . Then by (ii),  $\bar{A}$  is closed, so  $A$  is closed. Conversely, suppose  $A$  is closed. Then by (i),  $A \subseteq \bar{A}$ . Now we show  $\bar{A} \subseteq A$ . Let  $x \in \bar{A}$ . By definition, there exists a sequence  $(x_n)$  contained in  $A$  that converges to  $x$ . Since  $A$  is closed, then  $x \in A$ , so  $\bar{A} \subseteq A$ . Thus,  $\bar{A} = A$ .
- (iv) By (ii),  $\bar{A}$  is closed. By (iii),  $\bar{A} = \overline{\bar{A}}$ .
- (v) Let  $x \in \bar{A}$ . By definition, there exists a sequence  $(x_n)$  contained in  $A$  that converges to  $x$ . Since  $A \subseteq F$ , then  $(x_n)$  is also contained in  $F$ . Since  $F$  is closed, then  $x \in F$ .
- (vi) By (v), if  $F$  is closed and  $A \subseteq F$ , then  $\bar{A} \subseteq F$ . Therefore,  $\bar{A} \subseteq \bigcap \{F \subseteq \mathbb{R} : F \text{ is closed, and } A \subseteq F\}$ . By (ii),  $\bar{A}$  is closed, and by (i),  $A \subseteq \bar{A}$ . Thus, we have:

$$\bigcap \{F \subseteq \mathbb{R} : F \text{ is closed, and } A \subseteq F\} \subseteq \bar{A}$$

□

These properties can make it easier to prove statements about closures.

**Example 9.3.3 ► Using Properties of Closure**

If  $A \subseteq B$ , then  $\overline{A} \subseteq \overline{B}$ .

*Proof.* By (i),  $A \subseteq B \subseteq \overline{B}$ , and by (ii),  $\overline{B}$  is closed. Thus, by (v),  $\overline{A} \subseteq \overline{B}$ . □

The corresponding idea for open sets is the *interior* of a set.



# Compact Sets

In this chapter, we describe the idea of *compactness* for sets. This idea will prove tremendously useful in future chapters as it will enable us to bring a “finite quality” to an otherwise infinite set or idea. It turns out that, for metric spaces like the real numbers, compactness is equivalent to closed and bounded.

However, we will approach this idea from two (seemingly different) notions: sequential compactness which deals closed sets, and open cover compactness (or simply “compact”) that deals with open sets. Having these different approaches provides us with several avenues for proving things about compactness. We’ll dedicate a section to either approach, and ultimately show that these two ideas are equivalent in the real numbers (this is not generally true in all topological spaces).

## 10.1 Sequential Compactness

### Definition 10.1.1 ▶ Sequential Compactness

A set  $K \subseteq \mathbb{R}$  is *sequentially compact* if every sequence contained within  $K$  also has a subsequence that converges to an element of  $K$ .

### Example 10.1.2 ▶ Closed Intervals are Sequentially Compact

Let  $a, b \in \mathbb{R}$  where  $a \leq b$ . Then  $[a, b]$  is sequentially compact.

**Intuition:** We want to strictly use the definition of sequential compactness. So we will consider an arbitrary subsequence of  $[a, b]$ , and then find a subsequence that converges to something in  $[a, b]$ .

*Proof.* Let  $(x_n)$  be an arbitrary sequence entirely contained within  $[a, b]$ . Then  $(x_n)$  is bounded, so the Bolzano-Weierstrass Theorem guarantees the existence of a subsequence  $(x_{n_k})$  of  $(x_n)$  that  $x_{n_k}$  converges to some  $x \in \mathbb{R}$ . Since  $[a, b]$  is a closed set, then every sequence contained in  $[a, b]$  converges to a number in  $[a, b]$ . Then we know  $(x_{n_k})$  converges to some number in  $[a, b]$ .  $\square$

**Example 10.1.3 ▶ Open Sets are not Sequentially Compact**

$(0, 1]$  is not sequentially compact.

**Intuition:** Our strategy is to find a sequence contained within  $(0, 1]$  that converges to something outside of  $(0, 1]$ .

*Proof.* Consider the sequence  $(x_n) := 1/n$ . Then the sequence is entirely contained within  $(0, 1]$ , but it converges to 0 which is not in  $(0, 1]$ . Note that any arbitrary subsequence of  $(x_n)$  also converges to 0.

Reference to theorem 8.2.4 limits of subsequences

□

**Theorem 10.1.4 ▶ Sequentially compact just means closed and bounded**

Let  $E \subseteq \mathbb{R}$ . Then  $E$  is sequentially compact if and only if  $E$  is both closed and bounded.

*Proof.* First, suppose  $E$  is sequentially compact. Let  $(x_n)$  be an arbitrary sequence contained within  $E$  that converges to some  $x \in \mathbb{R}$ . Since  $E$  is sequentially compact, then there exists a subsequence  $(x_{n_k})$  of  $(x_n)$  such that  $(x_{n_k})$  converges to some  $y \in E$ . By (todo: Proposition 15.2 whatever this ref is), we have:

$$\lim_{k \rightarrow \infty} x_{n_k} = x$$

Since limits are unique (as proven in todo),  $x = y$ . Thus,  $(x_n)$  converges to  $y \in E$ , so  $E$  is closed.

To prove  $E$  is bounded, we proceed by contradiction. Suppose  $E$  is not bounded. Then, for each  $N \in \mathbb{N}$ , there exists  $x_n \in E$  such that  $|x_n| > n$ . Since  $E$  is sequentially compact, then there exists a subsequence  $(x_{n_k})$  of  $(x_n)$  such that  $x_{n_k}$  converges to a point in  $E$ . Note that  $|x_{n_k}| > n_k \geq k$ , so the sequence  $(x_{n_k})$  is unbounded and thus is divergent. This contradicts the fact that  $(x_{n_k})$  does converge. Thus, our supposition that  $E$  is not bounded was false, so  $E$  is in fact bounded.

Conversely, suppose that  $E$  is both closed and bounded. Let  $(x_n)$  be an arbitrary sequence entirely contained within  $E$ . Since  $E$  is bounded, then there exists  $M \in \mathbb{R}$  such that  $|x_n| \leq M$  for all  $n \in \mathbb{N}$ . By the Bolzano-Weierstrass Theorem, there exists a subsequence  $(x_{n_k})$  of  $(x_n)$  where  $(x_{n_k})$  converges to some  $x \in \mathbb{R}$ . Since  $E$  is closed and  $(x_{n_k})$  is contained within  $E$ , then its limit  $x$  must be in  $E$ . Therefore,  $E$  is sequentially compact. □

## 10.2 Open Cover Compactness

### Definition 10.2.1 ► Open Cover

Let  $A \subseteq \mathbb{R}$ . An **open cover** of  $A$  is a collection of open sets such that  $A$  is a subset of the union of that collection. We say that the collection **covers**  $A$ .

In other words, every number in  $A$  is in at least one of the open sets in the collection of open sets.

### Example 10.2.2 ► Open Cover Example #1

$[0, 1]$  is covered by  $\{B(x, 1/10) : x \in [0, 1]\}$ .

*Proof.* First note that  $B(x, 1/10)$  is open for all  $x \in [0, 1]$ . Let  $a \in [0, 1]$  be arbitrary. Then:

$$a \in B(a, 1/10) \in \{B(x, 1/10) : x \in [0, 1]\}$$

So it covers  $[0, 1]$ . □

### Example 10.2.3 ► Open Cover Example #2

$(0, 1)$  is covered by  $\{(x/2, 1) : x \in (0, 1)\}$ .

*Proof.* Note that  $(x/2, 1)$  is open for all  $x \in (0, 1)$ . Let  $a \in (0, 1)$ . Then:

$$a \in (a/2, 1) \in \{(x/2, 1) : x \in (0, 1)\}$$

So it covers  $(0, 1)$ . □

### Definition 10.2.4 ► Subcover

Given an open cover of a set, a **subcover** is a subset of the open cover that covers the set.

More formally, let  $A \subseteq \mathbb{R}$ , and let  $\{O_\lambda : \lambda \in \Lambda\}$  be an open cover of  $A$ . Then  $\{O_\lambda : \lambda \in \Lambda'\}$  is a **subcover** of  $\{O_\lambda : \lambda \in \Lambda\}$  if  $\Lambda' \subseteq \Lambda$  and  $A \subseteq \bigcup_{\lambda \in \Lambda'} O_\lambda$ .

In other words, a subcover is created by throwing away sets from the original cover, and the subcover still covers the original set. Also note that a cover is also one of its own subcovers. We say that a subcover is **finite** if there are only finitely many sets in the collection. Finiteness

in this context does not refer to the cardinality of the open sets in the collection, but rather the collection itself.

### Example 10.2.5 ▶ Open Cover Example #1 Revisited

Then open cover  $\{(x - 1/10, x + 1/10) : x \in [0, 1]\}$  of  $[0, 1]$  has a finite subcover.

For example, we can take  $\{B(0, 1/10), B(1/10, 1/10), \dots, B(1, 1/10)\}$  which only contains 11 open balls and is therefore a finite subcover of  $[0, 1]$ .

### Example 10.2.6 ▶ Open Cover Example #2 Revisited

The open cover  $\{(x/2, 1) : x \in (0, 1)\}$  of  $(0, 1)$  does not have a finite subcover.

*Proof.* Suppose for contradiction that there exists a finite subcover. Then, for some  $n \in \mathbb{N}$ , there exists  $x_1, x_2, \dots, x_n \in (0, 1)$  such that  $(0, 1) \subseteq \bigcup_{i=1}^n (x_i/2, 1)$ . Since there are finitely many “ $x$ ’s”, let  $y := \min\{x_1, \dots, x_n\}$ . Then, for all  $i \in \{1, 2, \dots, n\}$ , we have:

$$0 < \frac{y}{4} \leq \frac{x_i}{4} < \frac{x_i}{2}$$

so  $y/4 \notin (x_i/2, 1)$ . Thus, we found some  $y \in (0, 1)$  that is not in the supposed finite subcover. This contradicts the fact that the subcover must cover the entirety of  $(0, 1)$ . Therefore, there does not exist any finite subcover.  $\square$

### Definition 10.2.7 ▶ Open Cover Compactness

A set  $K \subseteq \mathbb{R}$  is **open cover compact** if every open cover of  $K$  has a finite subcover.

For example, the interval  $(0, 1)$  is not compact by Example 10.2.6. We found an open cover of  $(0, 1)$  that does not have a finite subcover.

### Example 10.2.8 ▶ Any finite set of points in $\mathbb{R}$ is open cover compact

*Proof.* Let  $A \subseteq \mathbb{R}$  be a finite set. Then  $A = \{x_1, x_2, \dots, x_n\}$  for some  $n \in \mathbb{N}$ . Let  $\{O_\lambda : \lambda \in \Lambda\}$  be an open cover of  $A$ . For each  $i \in \{1, 2, \dots, n\}$ , we have  $x_i \in A$ . Also note that  $A \subseteq \bigcup_{\lambda \in \Lambda} O_\lambda$ , so  $x_i$  is contained in some  $O_{\lambda_i}$  for some  $\lambda_i \in \Lambda$ . Then  $\{O_{\lambda_i} : i \in \{1, 2, \dots, n\}\}$  is a finite subcover of  $A$ , so  $A$  is compact.  $\square$

**Example 10.2.9 ▶ Open Cover Compactness Example #1**

Let  $A := \{1/n : n \in \{2, 3, 4, \dots\}\}$ . Then  $A$  is not open cover compact.

*Proof.* Consider the open cover:

$$\left\{ \left( \frac{1}{n+1}, \frac{1}{n-1} \right) : n \in \{2, 3, 4, \dots\} \right\}$$

For each  $n_0 \in \{2, 3, 4, \dots\}$ , note that  $1/n_0 \in (1/n_0 + 1, 1/n_0 - 1)$ , but if  $n \neq n_0$  then  $1/n_0 \notin (1/n + 1, 1/n - 1)$ . That is, we cannot remove any of the open intervals from the open cover, so there is no finite subcover. Therefore,  $A$  is not open cover compact.  $\square$

Surprisingly, we can add a single point to the set in the above example and make it a compact set.

**Example 10.2.10 ▶ Open Cover Compactness Example #2**

Let  $A := \{0\} \cup \{1/n : n \in \{2, 3, 4, \dots\}\}$ . Then  $A$  is open cover compact.

*Proof.* Let  $\{O_\lambda : \lambda \in \Lambda\}$  be an arbitrary open cover of  $A$ . Since  $0 \in A$ , then there must be some  $\lambda_0 \in \Lambda$  such that  $0 \in O_{\lambda_0}$ . Since this  $O_{\lambda_0}$  is an open set, then there exists some  $r > 0$  such that  $B(0, r) \subseteq O_{\lambda_0}$ . Let  $N \in \mathbb{N}$  such that  $N > 1/r$ . Then for any  $n > N$ , we have  $n > 1/r > 0$ . In particular,  $0 < 1/n < r$ . That is, for any  $n > N$ ,  $1/n \in B(0, r) \subseteq O_{\lambda_0}$ . For  $n \in \{2, 3, \dots, N\}$ , there exists some  $\lambda_n \in \Lambda$  such that  $1/n \in O_{\lambda_n}$ . We can now create a finite subcover as follows:

$$\{O_{\lambda_0}, O_{\lambda_2}, O_{\lambda_3}, \dots, O_{\lambda_N}\}$$

which is a finite subcover for  $A$ . Since our choice of the initial open cover was arbitrary, then every open cover of  $A$  has such a finite subcover. Therefore,  $A$  is open cover compact.  $\square$

**Theorem 10.2.11 ▶ Compactness implies boundedness**

If  $K \subseteq \mathbb{R}$  is open cover compact, then  $K$  is bounded.

*Proof.* Consider the following open cover of  $K$ :

$$O := \{(-n, n) : n \in \mathbb{N}\}$$

Since  $K$  is compact, there exists a finite subset of  $O$  that still covers  $K$  (i.e. a finite sub-

cover), which can be of the form:

$$\{(-n_i, n_i) : i \in \{1, 2, \dots, m\}\} \quad \text{such that} \quad m, n_1, n_2, \dots, n_m \in \mathbb{N}$$

Note  $K \subseteq \bigcup_{i=1}^m (-n_i, n_i)$ . Let  $N := \max\{n_1, n_2, \dots, n_m\}$ . Then:

$$K \subseteq \bigcup_{i=1}^m (-n_i, n_i) \subseteq (-N, N)$$

Therefore,  $K$  is bounded above by  $N$  and bounded below by  $-N$ , so  $K$  is bounded.  $\square$

### Theorem 10.2.12 ► Compactness implies closedness

If  $K \subseteq \mathbb{R}$  is open cover compact, then  $K$  is closed.

*Proof.* We will show that  $\mathbb{R} \setminus K$  is open. Let  $x \in \mathbb{R} \setminus K$ . For all  $r > 0$ , define  $O_r$  as:

$$O_r := \mathbb{R} \setminus [x - r, x + r]$$

Then for all  $r > 0$ ,  $O_r$  is open. Also,  $\bigcup_{r>0} O_r = \mathbb{R} \setminus \{x\}$ , so  $K \subseteq \mathbb{R} \setminus \{x\} = \bigcup_{r>0} O_r$ . That is,  $\{O_r : r > 0\}$  is an open cover for  $K$ . Since  $K$  is compact, there exists a finite subcover. Thus, for some  $n \in \mathbb{N}$ , there exists  $r_1, r_2, \dots, r_n$  such that  $K \subseteq \bigcup_{i=1}^n O_{r_i}$ . Let  $r := \min\{r_1, r_2, \dots, r_n\}$ . Then  $\bigcup_{i=1}^n O_{r_i} \subseteq O_r$ . Since  $K \subseteq O_r$ , we have:

$$\mathbb{R} \setminus K \supseteq O_r^c = [x - r, x + r] \supseteq (x - r, x + r)$$

so  $\mathbb{R} \setminus K$  is open. Therefore,  $K$  is closed.  $\square$

### Theorem 10.2.13 ► Closed subsets of compact sets are compact

Let  $K \subseteq \mathbb{R}$ . If  $K$  is open cover compact and  $E \subseteq K$  is closed, then  $E$  is open cover compact.

*Proof.* Let  $\{O_\lambda : \lambda \in \Lambda\}$  be an open cover for  $E$ . Since  $E$  is closed,  $\mathbb{R} \setminus E$  is open. Thus, the collection  $\{O_\lambda : \lambda \in \Lambda\} \cup \{\mathbb{R} \setminus E\}$  is an open cover for  $K$ . Since  $K$  is compact, there exists a finite subcover for  $K$  from  $\{O_\lambda : \lambda \in \Lambda\} \cup \{\mathbb{R} \setminus E\}$ . This subcover can either be:

1.  $\{O_{\lambda_i} : i \in \{1, 2, \dots, n\}\}$  for some  $n \in \mathbb{N}$ , in which case we have  $E \subseteq K \subseteq \bigcup_{i=1}^n O_{\lambda_i}$ , so  $E \subseteq \bigcup_{i=1}^n O_{\lambda_i}$ , or
2.  $\{O_{\lambda_i} : i \in \{1, 2, \dots, n\}\} \cup \{\mathbb{R} \setminus E\}$ , in which case we have  $E \subseteq K \subseteq \left(\bigcup_{i=1}^n O_{\lambda_i}\right) \cup \mathbb{R} \setminus E$ .

If  $x \in E$ , then  $x \notin \mathbb{R} \setminus E$ , so  $x \in \bigcup_{i=1}^n O_{\lambda_i}$ . Thus,  $E \subseteq \bigcup_{i=1}^n O_{\lambda_i}$ .  
 In either case, the collection  $\{O_{\lambda_i} : i \in \{1, 2, \dots, n\}\}$  is a finite subcover of  $E$ . Therefore,  $E$  is compact.  $\square$

### Theorem 10.2.14 ▶ Every closed interval is compact

Let  $a, b \in \mathbb{R}$  where  $a < b$ . Then  $[a, b]$  is open cover compact.

*Proof.* Let  $I := [a, b]$ , and let  $O := \{O_{\lambda} : \lambda \in \Lambda\}$  be an open cover of  $I$ . Suppose for contradiction there does not exist any finite subcover of  $I$  from  $O$ . Let  $I^l := \left[a, \frac{a+b}{2}\right]$ , and let  $I^r := \left[\frac{a+b}{2}, b\right]$ . Then  $I^l$  and/or  $I^r$  do not have a finite subcover from  $O$ . Choose such an interval and call it  $I_1$ . Repeat indefinitely for all  $n \in \mathbb{N}$ .

We claim there exists a sequence  $(I_j)$  of closed intervals such that:

1.  $I_j \subseteq I_{j-1} \subseteq I$ , where  $I_0 = I = [a, b]$ ,
2.  $\mathcal{L}(I_j) = \frac{\mathcal{L}(I)}{2^j}$ , where  $\mathcal{L}([c, d]) = d - c$  (i.e. the “length” of the interval), and
3. each  $I_j$  does not have a finite subcover from  $O$ .

For each  $j \in \mathbb{N}$ , let  $P(j)$  denote the statement: “all three properties hold for  $I_j$ .” Then the interval  $I_1$  defined above satisfies the three properties.

Maybe elaborate here

Now suppose that  $P(j)$  is true for some  $j \in \mathbb{N}$ . Then  $I_j$  does not have a finite subcover from  $O$ . Note that  $I_{j+1}$  is one of  $I_j^l$  or  $I_j^r$  which does not have a finite subcover from  $O$ .

1.  $I_{j+1} \subseteq I_j \subseteq I$ , so the first property holds.
2.  $\mathcal{L}(I_{j+1}) = \frac{1}{2} \mathcal{L}(I_j) = \frac{\mathcal{L}(I)}{2^{j+1}}$ , so the second property holds.
3.  $I_j$  does not have a finite subcover from  $O$ , so the third property holds.

Thus,  $P(j+1)$  is true. By the Principle of Induction,  $P(n)$  is true for all  $n \in \mathbb{N}$ . By the Nested Interval Property (todo: ref), we have:

$$\bigcap_{j \in \mathbb{N}} I_j \neq \emptyset$$

so there exists some  $x \in \bigcap_{j \in \mathbb{N}} I_j \subseteq [a, b]$ . Since  $O$  is an open cover for  $[a, b]$ , there exists an index  $\lambda_x \in \Lambda$  such that  $x \in O_{\lambda_x}$ . Since  $O_{\lambda_x}$  is open, there exists some radius  $r > 0$  such that  $B(x, r) \subseteq O_{\lambda_x}$ . Choose  $n \in \mathbb{N}$  such that  $\mathcal{L}(I_n) < r$   $\square$

**Theorem 10.2.15 ► Heine-Borel Theorem**

A set  $E \subseteq \mathbb{R}$  is open cover compact if and only if  $E$  is both closed and bounded.



# Index

## Definitions

2.1.1	Statement . . . . .	6	4.1.1	Peano Axioms for $\mathbb{N}$ . . . . .	22
2.1.2	Conjunction . . . . .	6	4.2.1	Integers $\mathbb{Z}$ . . . . .	24
2.1.3	Disjunction . . . . .	6	4.2.2	Even, Odd, Parity . . . . .	24
2.1.6	Negation . . . . .	7	4.3.1	Rational Numbers $\mathbb{Q}$ . . . . .	25
2.1.7	Implication . . . . .	7	4.3.3	Divides . . . . .	26
2.1.10	Logical Equivalence . . . . .	8	4.4.1	Field . . . . .	27
2.1.12	Converse . . . . .	9	4.5.1	Ordered Field . . . . .	29
2.1.14	Contrapositive . . . . .	9	4.5.3	Absolute Value . . . . .	29
3.1.1	Set . . . . .	13	4.6.1	Bounded Above, Bounded Be- low, Bounded . . . . .	30
3.1.2	Membership, $\in$ . . . . .	13	4.6.3	Maximum, Minimum . . . . .	31
3.1.3	Subset, $\subseteq$ . . . . .	13	4.6.4	Supremum . . . . .	31
3.1.4	Set Equality, $=$ . . . . .	13	4.6.7	Completeness . . . . .	32
3.1.5	Proper Subset, $\subsetneq$ . . . . .	14	4.6.9	Real Numbers $\mathbb{R}$ . . . . .	32
3.1.6	Empty Set ( $\emptyset$ ) . . . . .	14	5.0.1	Infimum . . . . .	34
3.1.9	Union . . . . .	14	7.0.1	Cardinality . . . . .	41
3.1.10	Intersection . . . . .	15	7.0.4	Power Set . . . . .	42
3.1.12	Set Minus . . . . .	15	7.0.6	Finite, Countably Infinite, Countable, Uncountable . . . . .	43
3.1.13	Complement . . . . .	16	7.1.1	Algebraic Number, Transcan- dental Number . . . . .	47
3.2.1	Cartesian Product . . . . .	17	8.0.1	Sequence . . . . .	49
3.2.2	Relation . . . . .	17	8.0.2	Convergence . . . . .	49
3.2.3	Function . . . . .	17	8.0.7	Divergence . . . . .	53
3.2.4	Function Image . . . . .	17	8.1.6	Increasing, Decreasing, Monotonic . . . . .	56
3.2.6	Function Inverse Image . . . . .	18	8.2.1	Subsequence . . . . .	57
3.3.1	Injective, One-to-one . . . . .	19	8.3.1	Limit Superior, Limit Inferior . . . . .	59
3.3.5	Surjective, Onto . . . . .	20	8.4.1	Cauchy Sequence . . . . .	63
3.3.8	Bijjective . . . . .	20	8.4.6	Completeness (in terms of Cauchy Sequences) . . . . .	64
3.3.9	Function Composition . . . . .	20			
3.3.11	Inverse Function . . . . .	21			

9.1.1	Open Set . . . . .	65
9.1.3	Open Ball . . . . .	65
9.2.1	Closed Set . . . . .	67
9.3.1	Closure of a Set . . . . .	69
10.1.1	Sequential Compactness . . . .	72
10.2.1	Open Cover . . . . .	74
10.2.4	Subcover . . . . .	74
10.2.7	Open Cover Compactness . . .	75

## Examples

1.0.1	Newton's Method . . . . .	4
1.0.2	Picard's Method . . . . .	5
2.1.4	Truth Table of Conjunction . .	7
2.1.5	Truth Table of Disjunction . .	7
2.1.8	Truth Table of Implication . .	8
2.1.9	Simple Statements . . . . .	8
2.1.11	Truth Table of Logical Equiv- alence . . . . .	9
2.1.13	Truth Table of Converse . . . .	9
2.1.15	Truth Table of Contrapositive .	10
2.2.2	Truth Table of Proof by Con- tradiction . . . . .	11
2.2.4	Logical Equivalence of Con- trapositive . . . . .	12
2.2.5	Proving Simple Logic State- ments . . . . .	12
3.1.8	Proving Simple Subset Relation	14
3.1.11	Indexed Sets . . . . .	15
3.2.5	Function Images . . . . .	17
3.2.7	Function Inverse Images . . .	18
3.3.3	Proving Injectivity . . . . .	19
3.3.4	Disproving Injectivity . . . . .	19
3.3.7	Proving Surjectivity . . . . .	20
3.3.12	. . . . .	21
4.1.4	Simple Proof by Induction . . .	23

4.2.4	Parity of Square . . . . .	25
4.3.5	. . . . .	26
4.3.6	$\sqrt{3}$ is not a Rational Number .	27
4.6.2	Upper and Lower Bounds . . .	31
4.6.6	Supremum of $(0, 1)$ . . . . .	32
5.0.5	. . . . .	35
7.0.2	Cardinality of $\mathbb{N}$ and $2\mathbb{N}$ . . . .	41
7.0.3	Cardinality of Intervals . . . .	41
7.0.5	Cardinality of $\mathbb{N}$ and $\mathcal{P}(N)$ . . .	43
8.0.4	$1/n$ converges to 0 . . . . .	50
8.0.5	. . . . .	50
8.0.6	. . . . .	52
8.0.8	Diverging Sequence . . . . .	53
8.1.5	Using the Properties . . . . .	56
8.2.2	Subsequences . . . . .	58
8.3.2	. . . . .	59
9.1.2	$[0, 1)$ is not open . . . . .	65
9.2.2	$(0, 1]$ is not closed . . . . .	67
9.3.3	Using Properties of Closure . .	71
10.1.2	Closed Intervals are Sequen- tially Compact . . . . .	72
10.1.3	Open Sets are not Sequen- tially Compact . . . . .	73
10.2.2	Open Cover Example #1 . . . .	74
10.2.3	Open Cover Example #2 . . . .	74
10.2.5	Open Cover Example #1 Re- visited . . . . .	75
10.2.6	Open Cover Example #2 Re- visited . . . . .	75
10.2.8	Any finite set of points in $\mathbb{R}$ is open cover compact . . . . .	75
10.2.9	Open Cover Compactness Ex- ample #1 . . . . .	76
10.2.10	Open Cover Compactness Ex- ample #2 . . . . .	76

## Techniques

2.2.1	Proof by Contradiction . . . . .	11
2.2.3	Proof by Contrapositive . . . . .	11
3.1.7	Proving a Subset Relation . . . . .	14
3.3.2	Proving a Function is Injective . . . . .	19
3.3.6	Proving a Function is Surjective . . . . .	20
4.1.3	Proof by Induction . . . . .	23
8.0.3	Proving Convergence . . . . .	50

## Lemmas

8.1.1	Approximating Zero . . . . .	53
8.1.4	Limit of a Constant Sequence . . . . .	55
8.2.3	Indices of Subsequences . . . . .	58
8.4.2	Convergent Sequences are Cauchy . . . . .	63
8.4.3	. . . . .	63
8.4.4	Cauchy Sequences are Bounded . . . . .	64
9.1.4	Open Intervals are Open Sets . . . . .	66
9.2.3	Closed Intervals are Closed Sets . . . . .	67

## Theorems

3.1.14	De Morgan's Laws for Sets . . . . .	16
3.3.10	Composition Preserves Injec- tivity and Surjectivity . . . . .	21
4.1.2	Principle of Induction (by the Peano Axioms) . . . . .	22
4.2.3	Parity Exclusivity . . . . .	24
4.3.2	$\sqrt{2}$ is not a Rational Number . . . . .	26
4.3.4	Division Algorithm . . . . .	26
4.4.2	Facts about Fields . . . . .	28
4.5.2	Facts about Ordered Fields . . . . .	29
4.5.4	Properties of Absolute Value . . . . .	30
4.5.5	Triangle Inequality . . . . .	30

4.6.5	Maximum is the Supremum . . . . .	31
4.6.8	$\mathbb{Q}$ is not complete . . . . .	32
4.6.10	$\sqrt{2}$ is a Real Number . . . . .	33
5.0.2	Existence of Infima in $\mathbb{R}$ . . . . .	34
5.0.3	Well-Ordering Principle . . . . .	34
5.0.4	Pushing Supremum . . . . .	35
5.0.6	Approximation Property of Suprema and Infima . . . . .	35
6.0.1	$\mathbb{N}$ is not Bounded Above . . . . .	37
6.0.2	Archimedean Principle . . . . .	37
6.0.3	Density of $\mathbb{Q}$ in $\mathbb{R}$ . . . . .	37
6.0.4	$\sqrt{2}$ is a Real Number . . . . .	38
6.0.5	Nested Interval Property . . . . .	39
7.0.7	$\mathcal{P}(\mathbb{N})$ is uncountable. . . . .	43
7.0.8	Subsets of Countable Sets are Countable . . . . .	44
7.0.9	Injectivity and Cardinality . . . . .	45
7.0.10	$ \mathbb{N} \times \mathbb{N}  =  \mathbb{N} $ . . . . .	45
7.0.11	Collection of Countable Sets . . . . .	45
7.0.12	$\mathbb{Q}$ is Countable . . . . .	46
7.0.13	$\mathbb{R}$ is Uncountable . . . . .	47
7.0.14	Irrational Numbers are Un- countable . . . . .	47
7.1.2	Schroeder-Bernstein . . . . .	48
7.1.3	Continuum Hypothesis . . . . .	48
8.1.2	Uniqueness of Limits . . . . .	53
8.1.3	Properties of Limits . . . . .	54
8.1.7	Monotone Sequence Theorem . . . . .	56
8.2.4	Limits of Subsequences . . . . .	58
8.3.3	. . . . .	60
8.3.4	Bolzano-Weierstrass Theorem . . . . .	61
8.3.5	Convergence iff $\limsup =$ $\liminf$ . . . . .	62
8.4.5	Cauchy Criterion . . . . .	64
9.1.5	Union of Open Sets is Open . . . . .	66

9.1.6	Finite Intersection of Open Sets is Open . . . . .	66	10.1.4	Sequentially compact just means closed and bounded . .	73
9.2.4	Intersection of Closed Sets is Closed . . . . .	68	10.2.11	Compactness implies bound- edness . . . . .	76
9.2.5	Finite Union of Closed Sets is Closed . . . . .	68	10.2.12	Compactness implies closed- ness . . . . .	77
9.2.6	Complement of an Open Set is Closed . . . . .	68	10.2.13	Closed subsets of compact sets are compact . . . . .	77
9.2.7	Complement of a Closed Set is Open . . . . .	68	10.2.14	Every closed interval is compact	78
9.3.2	Properties of Closures of Sets .	70	10.2.15	Heine-Borel Theorem . . . . .	79