

<u>Exan</u>	nple Transmission Route – Infe	ection from subpop j to i
To/From	Susceptible (S _i)	Vaccinated (V _i)
Infected (I _j)	S_i $\theta_{ij}I_jS_i$	$(1-e_{ff1})\beta_{ij}I_{j}V_{i}$
Vaccinated but Infected (Vi _j)	$(1-e_{ff2})\beta_{ij}Vi_{j}S_{i}$	$(1-e_{ff1})(1-e_{ff2})\beta_{ij}Vi_{j}V_{i}$

Per capita rate of re	covery	
Per capita rate of im	nmunity loss (vaccinated individuals)	
Per capita rate of immunity loss (from natural infection)		
Per capita rate of immunity loss (for those who have been infected/recovered and vaccinated)		
Vaccine Efficacy (pr	eventing infection)	
Vaccine Efficacy (pr	eventing onwards infectiousness)	
Compartment	Description (Proportion of population in)	
Compartment	Description (Proportion of population in)	
Compartment S_x	Description (Proportion of population in) Susceptibles in subpopulation x	
S_{x}	Susceptibles in subpopulation x	
S _x I _x	Susceptibles in subpopulation x Infectious individuals in subpopulation x	

Vaccinated and infectious individuals in subpopulation x

Vaccinated and recovered individuals in subpopulation x

Per capita rate of transmission from infectious subpopulation y to susceptible

Vaccination Rate (r_i)

 Vi_x

 Vr_x

Description

subpopulation x

Rate of Vaccination in subpopulation x

We model the vaccination rate as a function of the:

Total fraction of individuals in S, I and R compartments (available to

vaccinated) at the start of the vaccination period multiplied by the proportion divided by the duration of the vaccination period for the specific subgroup.

We assume that the rate of vaccination is constant (r_i^*1) , therefore the rate of vaccination in S, I and R compartments must be normalised to the total proportion of individuals in these three compartments.

Example equations sub-population i

$$\frac{dS_{i}}{dt} = \sigma_{1}V_{i} + \sigma_{2}R_{i} + \sigma_{3}(Rv_{i} + Vr_{i}) - \beta_{ii}S_{i}I_{i} - \beta_{ij}S_{i}I_{j} - \beta_{ik}S_{i}I_{k} - (1 - e_{ff2})\beta_{ii}S_{i}Vi_{i} - (1 - e_{ff2})\beta_{ij}S_{i}Vi_{j} - (1 - e_{ff2})\beta_{ik}S_{i}Vi_{k} - r_{i}\frac{S_{i}}{S_{i} + I_{i} + R_{i}}$$

$$\frac{dI_{i}}{dt} = \beta_{ii}S_{i}I_{i} + \beta_{ij}S_{i}I_{j} + \beta_{ik}S_{i}I_{k} + (1 - e_{ff2})\beta_{ii}S_{i}Vi_{i} + (1 - e_{ff2})\beta_{ij}S_{i}Vi_{j} + (1 - e_{ff2})\beta_{ik}S_{i}Vi_{k} - \gamma I_{i} - r_{i}\frac{I_{i}}{S_{i} + I_{i} + R_{i}}$$

$$\frac{dR_i}{dt} = \gamma I_i - r_i \frac{R_i}{S_i + I_i + R_i} - \sigma_2 R_i$$

$$\frac{dRv_i}{dt} = r_i \frac{I_i + R_i}{S_i + I_i + R_i} - \sigma_3 Rv_i$$

$$\frac{dV_i}{dt} = r_i \frac{S_i}{S_i + I_i + R_i} - (1 - e_{ff1})\beta_{ii}V_iI_i - (1 - e_{ff1})\beta_{ij}V_iI_j - (1 - e_{ff1})\beta_{ik}V_iI_k - (1 - e_{ff1})(1 - e_{ff2})\beta_{ii}V_iV_i - (1 - e_{ff1})(1 - e_{ff2})\beta_{ij}V_iV_i - (1 - e_{ff1})(1 - e_{ff2})\beta_{ik}V_iV_i - (1 - e_{ff1})(1 - e_{ff2})\beta_{ik}V_iV_i - (1 - e_{ff1})(1 - e_{ff2})\beta_{ij}V_iV_i - (1 - e_{ff1})(1 - e_{ff2})\beta_{ij}V_iV_i - (1 - e_{ff1})(1 - e_{ff2})\beta_{ij}V_iV_i - (1 - e_{ff1})(1 -$$

$$\frac{dVi_i}{dt} = (1 - e_{ff1})\beta_{ii}V_iI_i + (1 - e_{ff1})\beta_{ij}V_iI_j + (1 - e_{ff1})\beta_{ik}V_iI_k + (1 - e_{ff1})(1 - e_{ff2})\beta_{ii}V_iV_i + (1 - e_{ff1})(1 - e_{ff2})\beta_{ij}V_iV_i + (1 - e_{ff1})(1 - e_{ff2})\beta_{ik}V_iV_i + (1 - e_{ff1})(1 - e_{ff1})(1 - e_{ff1})\beta_{ik}V_iV_i + (1 - e_{ff1})(1 - e_{ff1})(1 - e_{ff1})\beta_{ik}V_iV_i + (1 - e_{ff1})(1 - e_{ff1})($$

$$\frac{dVr_i}{dt} = \gamma Vi_i - \sigma_3 Vr_i$$

Parameter	Description
r_{x}	Rate of Vaccination in subpopulation x
$oldsymbol{eta}_{xy}$	Per capita rate of transmission from infectious subpopulation y to susceptible subpopulation x
γ	Per capita rate of recovery
σ_1	Per capita rate of immunity loss (vaccinated individuals)
σ_2	Per capita rate of immunity loss (from natural infection)
σ_3	Per capita rate of immunity loss (for those who have been infected/recovered and vaccinated)
$e_{\it ff1}$	Vaccine Efficacy (preventing infection)

Vaccine Efficacy (preventing onwards infectiousness)

Compartment	Description (Proportion of population in)
S_{x}	Susceptibles in subpopulation x
I _x	Infectious individuals in subpopulation x
R_{x}	Recovered individuals in subpopulation x
Rv_x	Recovered or Infectious and subsequently vaccinated in subpopulation x
V_{x}	Vaccinated individuals in subpopulation x
Vi _x	Vaccinated and infectious individuals in subpopulation x
Vr _x	Vaccinated and recovered individuals in subpopulation x

BASELINE PARAMETERS

Model Details

We assume that 0.79% of the Scottish population is currently infected and 7.3% have already been infected and are now "Recovered". Each subpopulation is proportionately the same size.

Vaccine efficacy is modelled at 90% (both e_{ff1} and e_{ff2}) and coverage (P_i , P_j and P_k) aims for 90% of the entire subpopulation.

We currently assume a indefinite period of immunity (rate of immunity loss σ_1 , σ_2 and $\sigma_3 = 0$).

Initial Conditions

 $S_i = 0.3064$

 $S_i = 0.3064$

 $S_k = 0.3064$

 $I_i = 0.0071/3$

 $I_i = 0.0071/3$

 $I_{k} = 0.0071/3$

 $V_i = 0$

 $V_i = 0$

 $V_k = 0$

 $R_i = 0.073/3$

 $R_i = 0.073/3$

 $R_{k} = 0.073/3$

Subpopulation i

Target Coverage = 90%

Duration = 90 days

Trigger Date = day 0

Subpopulation j

Target Coverage = 90%

Duration = 90 days

Trigger Date = day 90

Subpopulation k

Target Coverage = 90%

Duration = 90 days Trigger Date = day 180

We explore 4 different scenarios:

Baseline

We model sequential vaccination of 3 sub-populations. Each vaccination schedule lasts 90 days and aims for 90% coverage of the available susceptibles, infecteds and recovereds in the vaccinated subpopulation at the beginning of the simulation.

After vaccination of each subpopulation, the sub-population is released from NPIs, with the R increasing from 1 to 4.2.

Full Release

We explore 3 different release scenarios:

1. First Group

We model a full release of the entire population (i, j and k) after the vaccination of the first sub-population (i) (after 90 days). This increases the R of the entire population from 1 to 4.2.

2. Middle Group

We model a full release of the entire population (i, j and k) after the vaccination of the second sub-population (j) (after 180 days). This increases the R of the entire population from 1 to 4.2.

3. Last Group

We model a full release of the entire population (i, j and k) after the vaccination of the final sub-population (k) (after 270 days). This increases the R of the entire population from 1 to 4.2.

Sequential Vaccination

We model sequential vaccination of 3 sub-populations. Each vaccination schedule lasts 90 days and aims for 90% coverage of the available susceptibles, infecteds and recovereds at the beginning of the simulation. After vaccination of each subpopulation, the sub-population is released from NPIs, with the R increasing from 1 to 4.2.


Attack Rate for fully susceptible individuals infected in sub-group i (excluding infections amongst those vaccinated) = **0.0084**


WAIFW Matrix (R) =
$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$
 During the pop i vaccination (t = 0-90)

WAIFW Matrix (R) = $\begin{pmatrix} 4.2 & 4.2 & 4.2 \\ 4.2 & 1 & 1 \\ 4.2 & 1 & 1 \end{pmatrix}$ During the pop j vaccination (t = 90-180)


WAIFW Matrix (R) = $\begin{pmatrix} 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 1 \end{pmatrix}$ During the pop k vaccination (t = 180-270)

WAIFW Matrix (R) = $\begin{pmatrix} 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \end{pmatrix}$ After final vaccination schedule (t = 270)





Time (Days)




Full Release - First Group (i)

We model a full release of the entire population (i, j and k) after the vaccination of the **first** sub-population (i). This increases the R of the entire population from 1 to 4.2.

Attack Rate for fully susceptible individuals infected in subgroup i (excluding infections amongst those vaccinated) = **0.032**

WAIFW Matrix (R) =
$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$
 Before first vaccination schedule (t < 90)

WAIFW Matrix (R) =
$$\begin{pmatrix} 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \end{pmatrix}$$
 After first vaccination schedule (t > 90)




Full Release - Second Group (j)

We model a full release of the entire population (i, j and k) after the vaccination of the **second** sub-population (j). This increases the R of the entire population from 1 to 4.2.

Attack Rate for fully susceptible individuals infected in sub-group i (excluding infections amongst those vaccinated) = **0.0081**

WAIFW Matrix (R) =
$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$
Before second vaccination schedule (t < 180)

WAIFW Matrix (R) =
$$\begin{pmatrix} 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \end{pmatrix}$$
 After second vaccination schedule (t > 180)




Full Release - Third Group (k)

We model a full release of the entire population (i, j and k) after the vaccination of the **last** sub-population (k). This increases the R of the entire population from 1 to 4.2.

Attack Rate for fully susceptible individuals infected in sub-group i (excluding infections amongst those vaccinated) = **0.0081**

WAIFW Matrix (R) =
$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$
 Before final vaccination schedule (t < 270)

WAIFW Matrix (R) =
$$\begin{pmatrix} 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \\ 4.2 & 4.2 & 4.2 \end{pmatrix}$$
 After final vaccination schedule (t > 270)



We now compare the total epidemic curve (the sum of the infectious compartment in each subpopulation) for 4 different scenarios:

- 1. With vaccination and sequential intervention release
- 2. Full release after vaccination of the first subpopulation (i)
- 3. Full release after vaccination of the second subpopulation (j)
- 4. Full release after vaccination of the last subpopulation (k)

The three shaded areas are the vaccination periods for subpopulation i, j and k respectively.

Sequential Vaccination

Attack Rate (i) - 0.009

Attack Rate (j) -0.012

Attack Rate (k) - 0.012

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0084**

Full Release after vaccination of i

Attack Rate (i) - 0.074

Attack Rate (j) - 0.173

Attack Rate (k) - 0.25

Attack Rate for fully susceptible individuals infected in sub-group i = **0.032**

Full Release after vaccination of j

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0081**


Full Release after vaccination of k

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0081**



HIGHER PRE-RELEASE R

Pre-release R is higher (R = 1.4) compared to baseline (R = 1.0)

We now compare the total epidemic curve (the sum of the infectious compartment in each subpopulation) for 4 different scenarios:

- 1. With vaccination and sequential intervention release
- 2. Full release after vaccination of the first subpopulation (i)
- 3. Full release after vaccination of the second subpopulation (j)
- 4. Full release after vaccination of the last subpopulation (k)

The three shaded areas are the vaccination periods for subpopulation i, j and k respectively.

Sequential Vaccination

Attack Rate (i) - 0.039

Attack Rate (j) - 0.062

Attack Rate (k) - 0.065

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0340**

Full Release after vaccination of i

Attack Rate (i) - 0.087

Attack Rate (j) - 0.204

Attack Rate (k) - 0.248

Attack Rate for fully susceptible individuals infected in sub-group i = **0.051**

Full Release after vaccination of j

Attack Rate (i) - 0.035

Attack Rate (j) - 0.057

Attack Rate (k) - 0.058

Attack Rate for fully susceptible individuals infected in sub-group i = 0.0317


Full Release after vaccination of k

Attack Rate (i) - 0.035

Attack Rate (j) - 0.057

Attack Rate (k) - 0.058

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0317**



PARTIAL RELAXATION

R is only released from 1 to 1.8 (25% to the baseline R release of 4.2)

We now compare the total epidemic curve (the sum of the infectious compartment in each subpopulation) for 4 different scenarios:

- 1. With vaccination and sequential intervention release
- 2. Full release after vaccination of the first subpopulation (i)
- 3. Full release after vaccination of the second subpopulation (j)
- 4. Full release after vaccination of the last subpopulation (k)

The three shaded areas are the vaccination periods for subpopulation i, j and k respectively.

Sequential Vaccination

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**

Full Release after vaccination of i

Attack Rate (i) - 0.009

Attack Rate (j) - 0.012

Attack Rate (k) - 0.013

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0083**

Full Release after vaccination of j

Attack Rate (i) - 0.009

Attack Rate (i) -0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = 0.008


Full Release after vaccination of k

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.008**



PARTIAL RELAXATION

R is only released from 1 to 2.6 (halfway to the baseline R release of 4.2)

We now compare the total epidemic curve (the sum of the infectious compartment in each subpopulation) for 4 different scenarios:

- 1. With vaccination and sequential intervention release
- 2. Full release after vaccination of the first subpopulation (i)
- 3. Full release after vaccination of the second subpopulation (j)
- 4. Full release after vaccination of the last subpopulation (k)

The three shaded areas are the vaccination periods for subpopulation i, j and k respectively.

Sequential Vaccination

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**

Full Release after vaccination of i

Attack Rate (i) - 0.015

Attack Rate (j) - 0.025

Attack Rate (k) - 0.042

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0113**

Full Release after vaccination of j

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**


Full Release after vaccination of k

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**



DECREASED INTER-GROUP MIXING

R on the non-diagonal elements of the WAIFW matrix is half relative to a baseline NPI release.

We now compare the total epidemic curve (the sum of the infectious compartment in each subpopulation) for 4 different scenarios:

- 1. With vaccination and sequential intervention release
- 2. Full release after vaccination of the first subpopulation (i)
- 3. Full release after vaccination of the second subpopulation (j)
- 4. Full release after vaccination of the last subpopulation (k)

The three shaded areas are the vaccination periods for subpopulation i, j and k respectively.

Sequential Vaccination

Attack Rate (i) - 0.009

Attack Rate (j) -0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**

Full Release after vaccination of i

Attack Rate (i) -0.03

Attack Rate (j) - 0.061

Attack Rate (k) - 0.157

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0183**

Full Release after vaccination of j

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**


Full Release after vaccination of k

Attack Rate (i) - 0.009

Attack Rate (j) - 0.011

Attack Rate (k) - 0.011

Attack Rate for fully susceptible individuals infected in sub-group i = **0.0082**




LOSS OF NATURAL IMMUNITY, VACCINE IMMUNITY AND BOTH

Waning immunity is modelled in those who are natural infected (σ_2 = 1/6 months⁻¹), vaccinated (σ_1 = 1/6 months⁻¹) and then both are modelled in tandem (both σ_1 = 1/6 months⁻¹ and σ_2 = 1/6 months⁻¹)

Waning Natural Immunity $(\sigma_2 = 1/6 \text{ months}^{-1})$

Effects of Vaccination (ALL)



Scenario - Seg Release - After i - After j - After k

Sequential Vaccination

Attack Rate (i) -0.009Attack Rate (j) - 0.012Attack Rate (k) - 0.013 Attack Rate for fully susceptibles in i - 0.0088

Full Release after i vacc

Attack Rate (i) - 0.088 Attack Rate (j) -0.212Attack Rate (k) - 0.302Attack Rate for fully susceptibles in i - 0.0374

Full Release after j vacc


Attack Rate (i) -0.009Attack Rate (j) -0.012Attack Rate (k) - 0.012Attack Rate for fully susceptibles in i - 0.0084

Full Release after k vacc

Attack Rate (i) - 0.009 Attack Rate (j) -0.012Attack Rate (k) - 0.012Attack Rate for fully susceptibles in i - 0.0084

Waning Vaccine Immunity $(\sigma_1 = 1/6 \text{ months}^{-1})$

Effects of Vaccination (ALL)



Scenario - Seq Release - After i - After j - After k

Sequential Vaccination

Attack Rate (i) - 0.179 Attack Rate (j) -0.133Attack Rate (k) - 0.187Attack Rate for fully susceptibles in i - 0.159

Full Release after i vacc

Attack Rate (i) - 0.164 Attack Rate (j) -0.221Attack Rate (k) - 0.279Attack Rate for fully susceptibles in i - 0.1277

Full Release after i vacc


Attack Rate (i) - 0.208 Attack Rate (i) -0.177Attack Rate (k) - 0.146Attack Rate for fully susceptibles in i - 0.194

Full Release after k vacc

Attack Rate (i) - 0.064 Attack Rate (i) -0.06Attack Rate (k) - 0.05Attack Rate for fully susceptibles in i - 0.0621

Waning Natural and Vaccine Immunity (both $\sigma_1 = 1/6$ months⁻¹ and $\sigma_2 = 1/6$ months⁻¹)





Scenario - Seq Release - After i - After j - After k

Sequential Vaccination

Attack Rate (i) -0.307Attack Rate (i) - 0.244 Attack Rate (k) - 0.287 Attack Rate for fully susceptibles in i - 0.2779

Full Release after i vacc

Attack Rate (i) - 0.233 Attack Rate (j) - 0.282Attack Rate (k) - 0.35Attack Rate for fully susceptibles in i - 0.1886

Full Release after j vacc

Attack Rate (i) - 0.273 Attack Rate (i) – 0.239 Attack Rate (k) - 0.214 Attack Rate for fully susceptibles in i - 0.255


Full Release after k vacc

Attack Rate (i) -0.205Attack Rate (j) -0.193Attack Rate (k) - 0.163 Attack Rate for fully susceptibles in i - 0.19897

VACCINE INDUCED BLOCKING OF COVID-19 TRANSMISSION

We model varying levels of vaccine induced transmission blocking. We model variations in reductions to vaccinated individuals becoming infected (e_{ff1}) and transmission from vaccinated individuals (e_{ff2}).

We model 3 levels of transmission blocking: 50%, 75% and 90% (baseline).



Attack Rate (i) - 0.05

Attack Rate (j) -0.052

Attack Rate (k) - 0.052

susceptibles in i - 0.0152

Attack Rate for fully

Transmission Blocking: 50%


Sequential Vaccination Attack Rate (i) - 0.136 Attack Rate (i) -0.125Attack Rate (k) - 0.135Attack Rate for fully susceptibles in i - 0.027

Full Release after i vacc Attack Rate (i) - 0.226 Attack Rate (j) -0.26Attack Rate (k) - 0.281

Full Release after k vacc Attack Rate (i) - 0.01 Attack Rate (j) -0.012Attack Rate (k) - 0.012 Attack Rate for fully Attack Rate for fully susceptibles in i - 0.035 susceptibles in i - 0.0085

Transmission Blocking: 75% $(e_{ff1} \text{ and } e_{ff2} = 0.75)$

Effects of Vaccination (ALL)



Scenario - Seg Release - After i - After i - After k

Sequential Vaccination

Attack Rate (i) - 0.011 Attack Rate (j) -0.013Attack Rate (k) - 0.013Attack Rate for fully susceptibles in i - 0.0087

Full Release after i vacc

Attack Rate (i) -0.134Attack Rate (j) - 0.207Attack Rate (k) - 0.261Attack Rate for fully susceptibles in i - 0.0332

Full Release after j vacc


Attack Rate (i) - 0.009 Attack Rate (i) -0.011Attack Rate (k) - 0.011Attack Rate for fully susceptibles in i - 0.0083

Full Release after k vacc

Attack Rate (i) -0.009Attack Rate (i) - 0.011 Attack Rate (k) - 0.011Attack Rate for fully susceptibles in i - 0.0082

Transmission Blocking: 90%

 $(e_{ff1} \text{ and } e_{ff2} = 0.9)$ Effects of Vaccination (ALL)



Sequential Vaccination Attack Rate (i) - 0.009 Attack Rate (j) -0.012Attack Rate (k) - 0.012 Attack Rate for fully susceptibles in i - 0.0084

Full Release after i vacc

Attack Rate (i) -0.074Attack Rate (j) - 0.173Attack Rate (k) - 0.25Attack Rate for fully susceptibles in i - 0.032

Full Release after j vacc

Attack Rate (i) - 0.009 Attack Rate (j) -0.011Attack Rate (k) - 0.011 Attack Rate for fully susceptibles in i - 0.0082

Full Release after k vacc

Attack Rate (i) -0.009Attack Rate (j) -0.011Attack Rate (k) - 0.011 Attack Rate for fully susceptibles in i - 0.0082

SENSITIVITY TO VACCINE COVERAGE

We model varying levels of vaccine coverage. Coverage is varied from 90% (baseline), 70% to 50%.

Vaccine Coverage = 50% Vaccine Coverage = 70% Vaccine Coverage = 90% Effects of Vaccination (ALL) Effects of Vaccination (ALL) Effects of Vaccination (ALL) 0.250.25 0.25 -Attack Rate (Baseline) = 0.477 Attack Rate (Baseline) = 0.227 Attack Rate (Baseline) = 0.032 Attack Rate (After i vac, t > 90) = 0.724 Attack Rate (After i vac, t > 90) = 0.628 Attack Rate (After i vac, t > 90) = 0.496 Attack Rate (After j vac, t > 180) = 0.484 Attack Rate (After j vac, t > 180) = 0.069 Attack Rate (After j vac, t > 180) = 0.031 Attack Rate (After k vac, t > 270) = 0.041 Attack Rate (After k vac, t > 270) = 0.034 Attack Rate (After k vac, t > 270) = 0.031 0.20 0.20 0.20 0.15 0.15 Prevalence Prevalence Prevalence 0.10 0.10 0.10 0.05 0.05 0.05 200 300 300 300 200 Time (Days) Time (Days) Time (Days) Scenario - Seq Release - After i - After j - After k Scenario - Seq Release - After i - After j - After k Scenario - Seq Release - After i - After j - After k

Sequential Vaccination

Attack Rate (i) – 0.154 Attack Rate (j) – 0.142 Attack Rate (k) – 0.182 Attack Rate for fully susceptibles in i - **0.131**

Full Release after i vacc

Attack Rate (i) – 0.182 Attack Rate (j) – 0.256 Attack Rate (k) – 0.286 Attack Rate for fully susceptibles in i - **0.147**

Full Release after j vacc

Attack Rate (i) – 0.151 Attack Rate (j) – 0.153 Attack Rate (k) – 0.181 Attack Rate for fully susceptibles in i - **0.128**

Full Release after k vacc

Attack Rate (i) – 0.012 Attack Rate (j) – 0.014 Attack Rate (k) – 0.015 Attack Rate for fully susceptibles in i - **0.0120**

Sequential Vaccination

Attack Rate (i) – 0.067 Attack Rate (j) – 0.067 Attack Rate (k) – 0.092 Attack Rate for fully susceptibles in i - **0.053**

Full Release after i vacc

Attack Rate (i) -0.13Attack Rate (j) -0.223Attack Rate (k) -0.274Attack Rate for fully susceptibles in i - **0.088**

Full Release after j vacc

Attack Rate (i) – 0.021 Attack Rate (j) – 0.024 Attack Rate (k) – 0.024 Attack Rate for fully susceptibles in i - **0.0186**

Full Release after k vacc

Attack Rate (i) -0.01Attack Rate (j) -0.012Attack Rate (k) -0.012Attack Rate for fully susceptibles in i - **0.0094**

Sequential Vaccination

Attack Rate (i) -0.009Attack Rate (j) -0.012Attack Rate (k) -0.012Attack Rate for fully susceptibles in i -0.0084

Full Release after i vacc

Attack Rate (i) – 0.073 Attack Rate (j) – 0.173 Attack Rate (k) – 0.25 Attack Rate for fully susceptibles in i - **0.032**

Full Release after j vacc

Attack Rate (i) – 0.008 Attack Rate (j) – 0.011 Attack Rate (k) – 0.011 Attack Rate for fully susceptibles in i - **0.008**

Full Release after k vacc

Attack Rate (i) -0.009Attack Rate (j) -0.011Attack Rate (k) -0.011Attack Rate for fully susceptibles in i - **0.008**