Exercise # 1. Numerical methods for ODES.

Alexandre Rodrigues (2039952)

November 20, 2021

${\bf Intro}$

Methods

Answers

Question 1

$$y(t) = e^{-5t} (1)$$




Figure 1: Absolute error in function of time using Forward Euler method to compute y(1)

We got a maximum error of $4.2916\times 10^{10}...$




Figure 2: Absolute error in function of time using RK4 method to compute y(1)

We got a maximum error of $4.3146 \times 10^{10}...$

Comment the different behavior observed by the numerical method.

The Simpson's method has an empty stability region as proved by: ... We can notice the difference in the initial conditions in our results. The FE calculation for y(2) is better then the RK4 calculation given the best final error. This is, although, not that relevant, the difference is of about $0.5 \times 10^{-10}\%$.

Question 2

The exact solution can be found as:

$$y(t) = \frac{1}{10t+1} \tag{2}$$




Figure 3: LogLog plot of the error as a function of the number of steps.

h	error	
3.125000×10^{-2}	2.291844×10^{-7}	
1.562500×10^{-2}	1.785763×10^{-8}	
7.812500×10^{-3}	1.160234×10^{-9}	
3.906250×10^{-3}	7.312862×10^{-11}	
1.953125×10^{-3}	4.579586×10^{-12}	
9.765625×10^{-4}	2.863750×10^{-13}	

The error reduces with the increase in the number of steps due to the decrease of h as expected in theory. . . .

Question 3

Question 4

Method	Number of steps	Error	CPU time elapsed
ODE45	9445	1.155269×10^{-5}	8.791882s
CN	100	4.467899×10^{-3}	208.038764s
CN	1000	4.441078×10^{-4}	514.197773s
CN	10000	4.438412×10^{-5}	3086.958397s
BDF3	100	4.482679×10^{-3}	188.455409s
BDF3	1000	4.442484×10^{-4}	557.765280s
BDF3	10000	4.438552×10^{-5}	3399.768021s

Question 5




Figure 4: Evolution of the number of preys and predators.

Results

Outputs