

Course Notes

Introduction to Probability

Alex Rutar

BSM Fall 2018

Contents

1	Fundamentals	3
1.1	Basic Principles	3
1.1.1	Probability Spaces	3
1.1.2	Ω	3
1.1.3	\mathcal{F}	4
1.1.4	\mathbb{P}	4
1.1.5	Consequences	4
1.1.6	Examples with Finite Uniform Probabilities	6
1.2	Conditional Probability	8
1.2.1	Basic Principles	8
1.2.2	Bayes' Formula	10
1.3	Independent Events	12
1.3.1	Definitions	12
1.3.2	Independent Trials	12
1.3.3	Random Walks	13

Chapter 1

Fundamentals

1.1 Basic Principles

1.1.1 Probability Spaces

A probability space is a triple $(\Omega, \mathcal{F}, \mathbb{P})$.

1.1.2 Ω

Ω is a set, called the sample space, and $\omega \in \Omega$ are called outcomes and $A \subset \Omega$ are called events.

Ex. 1.1.1 A horserace with 3 horses, a, b, c , has $\Omega = \{(a, b, c), (a, c, b), \dots, (c, b, a)\}$. Then $|\Omega| = 6$ and $A = \{a \text{ wins the race}\} = \{(a, b, c), (a, c, b)\}$.

Ex. 1.1.2 Roll two fair dice, a white die and a yellow die. Then $\Omega = \{(1, 1), (1, 2), \dots, (6, 6)\}$ and $|\Omega| = 36$.

Ex. 1.1.3 Continue flipping a coin until there is a head. Then

$$\Omega = \{(H), (T, H), (T, T, H), \dots\}$$

Then define

$$A = \{\text{there are an even number of rolls}\} = \{(T, H), (T, T, T, H), \dots\}$$

Ex. 1.1.4 Consider $\Omega = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 100\}$. Then $A = \{\text{you score 50 points}\} = \{(x, y) \mid x^2 + y^2 \leq 1\}$.

Def'n. 1.1.5 If $A \cap B = \emptyset$, we say that A and B are **mutually exclusive** events. If $A \subset B$, we say that A **implies** B .

Write $A^c = \Omega \setminus A$. Recall distributivity, the deMorgan relations, etc.

1.1.3 \mathcal{F}

\mathcal{F} is a collection of subsets of Ω , which denote the events that we consider.

- If Ω is countable, then typically \mathcal{F} is just the collection of all subsets of Ω .
- If Ω is a domain in \mathbb{R}^n , then it is a strict subset of \mathbb{R}^n .

In any case, \mathcal{F} has to be closed under the following operations:

1. $\Omega \in \mathcal{F}$
2. If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$
3. If $A_1, A_2, \dots \in \mathcal{F}$, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$.

in other words, that \mathcal{F} is a σ -algebra.

1.1.4 \mathbb{P}

Finally, $\mathbb{P}: \mathcal{F} \rightarrow \mathbb{R}$ is a function that satisfies 3 axioms:

1. For any $A \in \mathcal{F}$, then $\mathbb{P}(A) \geq 0$
2. $\mathbb{P}(\Omega) = 1$
3. (σ -additivity) Let A_1, A_2, A_3, \dots be a sequence of mutually exclusive events. Then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$$

1.1.5 Consequences

- $\mathbb{P}(A^c) + \mathbb{P}(A) = \mathbb{P}(A \cup A^c) = \mathbb{P}(\Omega) = 1$.
- If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$ since $\mathbb{P}(B) = \mathbb{P}((A^c \cap B) \cup (A \cap B)) = \mathbb{P}(A^c \cap B) + \mathbb{P}(A \cap B) = \mathbb{P}(A^c \cap B) + \mathbb{P}(A)$
- For any A, B , we have

$$\mathbb{P}(A \cup B) = \mathbb{P}((A^c \cap B) \cup (A \cap B) \cup (A \cap B^c)) = \mathbb{P}(A^c \cap B) + \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^c) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$

Similarly,

$$\mathbb{P}(A \cup B \cup C) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) - \mathbb{P}(A \cap B) - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C) + \mathbb{P}(A \cap B \cap C)$$

which generalizes arbitrarily:

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) = \sum_{r=1}^n (-1)^{r+1} \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_r})$$

PROOF We have already proved the base case for $n = 2$, so assume the formula holds for a union of n events. Then

$$\mathbb{P}(A_1 \cup \dots \cup A_n \cup A_{n+1}) = \mathbb{P}(A_1 \cup \dots \cup A_n) + \mathbb{P}(A_{n+1}) - \mathbb{P}((A_1 \cup \dots \cup A_n) \cap A_{n+1})$$

We can distribute the first and third terms using the induction hypothesis, and the result follows. \square

Def'n. 1.1.6 We say D_1, D_2, \dots is a **decreasing** sequence of events of $D_{k+1} \subset D_k$. We say D_1, D_2, \dots is a **increasing** sequence of events of $D_{k+1} \supset D_k$.

Let $\lim_{n \rightarrow \infty} D_n = \bigcap_{n=1}^{\infty} D_n$ and $\lim_{n \rightarrow \infty} I_n = \bigcup_{n=1}^{\infty} I_n$.

Prop. 1.1.7 σ -additivity implies that for any increasing sequence,

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} I_n\right) = \lim_{n \rightarrow \infty} \mathbb{P}(I_n)$$

and similarly for any decreasing sequence

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} D_n\right) = \lim_{n \rightarrow \infty} \mathbb{P}(D_n)$$

PROOF Note that (2) implies (1): if D_k is a decreasing sequence, then $I_k = D_k^c$ is an increasing sequence and

$$\left(\lim_{n \rightarrow \infty} D_n\right)^c = \left(\bigcap_{n=1}^{\infty} D_n\right)^c = \bigcup_{n=1}^{\infty} I_n = \lim_{n \rightarrow \infty} I_n$$

and taking probabilities,

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} D_n\right) = 1 - \mathbb{P}\left(\lim_{n \rightarrow \infty} I_n\right) = 1 - \lim_{n \rightarrow \infty} \mathbb{P}(I_n) = \lim_{n \rightarrow \infty} \mathbb{P}(D_n)$$

To prove that σ -additivity implies (1), let I_1, I_2, \dots be increasing. Let $A_1 = I_1$ and for $k \geq 2$ let $A_k = I_k \setminus I_{k-1}$. Then A_1, A_2, \dots are mutually exclusive and for any $k \geq 1$,

$$\bigcup_{k=1}^K A_k = I_K$$

Thus

$$\bigcup_{k=1}^{\infty} A_k = \lim_{n \rightarrow \infty} I_n$$

Now note that $\mathbb{P}(I_K) = \sum_{k=1}^K \mathbb{P}(A_k)$ while

$$\begin{aligned} \mathbb{P}\left(\lim_{n \rightarrow \infty} I_n\right) &= \mathbb{P}\left(\bigcup_{k=1}^{\infty} A_k\right) \\ &= \sum_{k=1}^{\infty} \mathbb{P}(A_k) \\ &= \lim_{K \rightarrow \infty} \sum_{k=1}^K \mathbb{P}(A_k) \\ &= \lim_{K \rightarrow \infty} \mathbb{P}(I_K) \end{aligned}$$

\square

1.1.6 Examples with Finite Uniform Probabilities

We assume that $\Omega = \{\omega_1, \omega_2, \dots, \omega_N\}$ and $\mathbb{P}(\{\omega_i\}) = \mathbb{P}(\{\omega_j\})$. Then $\mathbb{P}(\{\omega_i\}) = \frac{1}{N}$ and $\mathbb{P}(A) = |A|/N$.

Ex. 1.1.8 In an urn there are 6 blue balls and 5 red balls. Draw 3 balls out of this 11. What is the chance that among the 3 there are exactly 2 blue balls and 1 red ball?

Let us pretend that the balls are labelled, 1 through 11, and set Ω to be all the ordered triples of disjoint elements. Then $A = \{\text{exactly 2 blue and 1 red}\}$, and note that $A = A^1 \cup A^2 \cup A^3$ where A^i has a red in position i and blue in the other two positions. Now, $|A^i| = 5 \cdot 6 \cdot 5$, so $|A| = 3 \cdot 6 \cdot 5 \cdot 6$ and

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|} = \frac{3 \cdot 6 \cdot 5 \cdot 6}{11 \cdot 10 \cdot 9}$$

We now suppose that $\Omega = \{\Lambda \subset \{1, \dots, 11\} \mid |\Lambda| = 3\}$, so $|\Omega| = \binom{11}{3}$. Now

$$A = \{\Lambda_1 \cup \Lambda_2 \mid \Lambda_1 \subset \{1, \dots, 6\}, |\Lambda_1| = 2, \Lambda_2 \subset \{7, \dots, 11\}, |\Lambda_2| = 1\}$$

So $|A| = \binom{6}{2} \cdot 5$.

Ex. 1.1.9 Consider a group of N people. What is the chance that there is at least one pair among them who have the same birthday?

Define $\Omega = \{(i_1, i_2, \dots, i_N) \mid i_j \in \{1, \dots, 365\}\}$. We want $A = \{\text{there is at least one common birthday}\}$. We can write

$$A^c = \{(i_1, \dots, i_N) \in \Omega \mid i_j \neq i_k \forall j \neq k\}$$

Then $|A^c| = 365 \cdot 364 \cdots (365 - N + 1)$ and

$$P_N = \mathbb{P}(A) = 1 - \mathbb{P}(A^c) = 1 - \frac{365 \cdot 364 \cdots (365 - N + 1)}{365^N}$$

Ex. 1.1.10 Suppose we have N people at a party. The following day, everyone leaves one after another, and chooses a single phone from a pile. What is the chance that nobody chooses her own phone?

Define $\Omega = \{(i_1, \dots, i_N) \mid \text{permutations of } \{1, \dots, N\}\}$, so $\omega = (i_1, \dots, i_k)$ means person k chooses phone i_k . Then $|\Omega| = N!$. Fix $B = \{\text{nobody picks her/his phone}\}$. Define $A_1 = \{\text{person 1 picks his phone}\}$, so $|A_1| = (N - 1)!$, and similarly for A_2 , etc. Then $B = A_1^c \cap A_2^c \cdots \cap A_N^c = (A_1 \cup \dots \cup A_N)^c$, and $\mathbb{P}(A_i) = \frac{1}{N}$. Now in general,

$$\mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) = \frac{(N - k)!}{N!}$$

for i_k distinct. Thus we now have

$$\begin{aligned} \mathbb{P}(B) &= 1 - \mathbb{P}(A_1 \cup A_2 \cup \dots \cup A_N) \\ &= 1 - \sum_{r=1}^N (-1)^{r+1} \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq N} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_r}) \\ &= \sum_{r=1}^N (-1)^{r+1} \binom{N}{r} \frac{(N - r)!}{N!} \\ &= \sum_{r=1}^N (-1)^{r+1} \frac{1}{r!} \end{aligned}$$

so that

$$\mathbb{P}(B) = 1 + \sum_{r=1}^N (-1)^r \frac{1}{r!} = \sum_{r=0}^N (-1)^r \frac{1}{r!}$$

Thus $\lim_{N \rightarrow \infty} \mathbb{P}(B) = \frac{1}{e}$.

Ex. 1.1.11 (Round table seating) Consider a round table with 20 seats, and 10 married couples sit. What is the chance that no couples sit together?

Define $\Omega = \{\text{permutations of } \{1, \dots, 20\} / \sim\}$ where $(i_1, \dots, i_{20}) \sim (i_{20}, i_1, \dots, i_{19})$. Then $|\Omega| = 19!$. Define $B = \{\text{no couples together} = A_1^c \cap A_2^c \cap \dots \cap A_{10}^c\}$, where

$$A_k = \{\text{the } k\text{th woman sits next to her spouse}\}$$

so that

$$\mathbb{P}(B) = 1 - \mathbb{P}(A_1 \cup \dots \cup A_{10})$$

Note that

$$\mathbb{P}(A_i) = \frac{18! \cdot 2}{19!} = \frac{2}{19}$$

by “joining” the couple together, arranging them around the table, and permuting the couple internally. This generalizes to

$$\mathbb{P}(A_{i_1} \cap \dots \cap A_{i_r}) = \frac{2^r (19-r)!}{19!}$$

Then by inclusion-exclusion,

$$\mathbb{P}(B) = 1 - \binom{10}{1} \cdot \frac{18! \cdot 2}{19!} + \binom{10}{2} \frac{17! \cdot 2^2}{19!} - \binom{10}{3} \frac{16! \cdot 2^3}{19!} \dots + \binom{10}{10} \frac{9! \cdot 2^{10}}{19!} \approx 0.339$$

Ex. 1.1.12 (Poker hand probabilities) A poker hand is a straight if the 5 cards are of increasing value and not all of the same suit, starting with A, 2, 3, 4, ..., 10.

Define $\Omega = \{5 \text{ element subsets of the } 52 \text{ cards}\}$. Then $|\Omega| = \binom{52}{5}$. Thus

$$\mathbb{P}(\text{straight}) = \frac{10 \cdot (4^5 - 4)}{\binom{52}{5}}$$

$$\mathbb{P}(\text{full house}) = \frac{13 \cdot 12 \cdot \binom{4}{3} \cdot \binom{4}{2}}{\binom{52}{5}}$$

Ex. 1.1.13 (Bridge hand probabilities) In bridge, each of the 4 players get 13 cards. Let $\Omega = \{13 \text{ cards that North gets}\}$.

$$\mathbb{P}(\text{North receives all spaces}) = \frac{1}{\binom{52}{13}}$$

$\mathbb{P}(\text{North does not receive all 4 suits of any value}) = \mathbb{P}(\text{There is some value such that all suits are at N})$

Let $V_k = \{\text{North gets all four suits of value } k\}$. Then

$$\mathbb{P}(V_1) = \frac{\binom{48}{9}}{\binom{52}{13}}$$

$$\mathbb{P}(V_1 \cap V_2) = \frac{\binom{44}{5}}{\binom{52}{13}}$$

$$\mathbb{P}(V_1 \cap V_2 \cap V_4) = \frac{\binom{40}{1}}{\binom{52}{13}}$$

Thus

$$1 - \mathbb{P}(V_1 \cup V_2 \cup \dots \cup V_{13}) = 1 - \frac{\binom{48}{9}}{\binom{52}{13}} \cdot 13 + \binom{13}{2} \frac{\binom{44}{5}}{\binom{52}{13}} - \binom{13}{3} \frac{40}{\binom{52}{5}}$$

What is the chance that each player receives one ace? There are

$$\frac{52!}{13!13!13!13!}$$

possible hands. There are $4!$ ways to arrange the aces, which gives

$$\mathbb{P}(E) = \frac{4! \binom{48}{12,12,12,12}}{\binom{52}{13,13,13,13}}$$

1.2 Conditional Probability

1.2.1 Basic Principles

Suppose we roll two fair dice. Then $\mathbb{P}(\text{the sum is } 10) = \frac{3}{36} = \frac{1}{12}$. Suppose instead that the white dice is rolled first, and it turns up 6. Now the probability that the sum is 10 is now $1/6$.

Def'n. 1.2.1 Given an event E with $\mathbb{P}(E) > 0$, for any event F , let $\mathbb{P}(F|E) = \frac{\mathbb{P}(F \cap E)}{\mathbb{P}(E)}$. We call this the **conditional probability of F given E** .

Prop. 1.2.2 Fix E with $\mathbb{P}(E) > 0$ and consider $\mathbb{P}(\cdot|E) : \mathcal{F} \rightarrow \mathbb{R}$. This function satisfies the axioms of probability.

PROOF 1. $\mathbb{P}(F|E) \geq 0$ for all $F \in \mathcal{F}$.

$$2. \mathbb{P}(\Omega|E) = \frac{\mathbb{P}(E \cap \Omega)}{\mathbb{P}(E)} = 1$$

3. If F_1, F_2, \dots are mutually exclusive, then

$$\begin{aligned}\mathbb{P}\left(\bigcup_{i=1}^{\infty} F_i | E\right) &= \frac{\mathbb{P}\left(\left(\bigcup_{i=1}^{\infty} F_i\right) \cap E\right)}{\mathbb{P}(E)} \\ &= \frac{\mathbb{P}\left(\bigcup_{i=1}^{\infty} (E \cap F_i)\right)}{\mathbb{P}(E)} \\ &= \sum_{n=1}^{\infty} \frac{\mathbb{P}(F_n \cap E)}{\mathbb{P}(E)} \\ &= \sum_{n=1}^{\infty} \mathbb{P}(F_n | E)\end{aligned}\quad \square$$

Prop. 1.2.3 We have $\mathbb{P}(E \cap F) = \mathbb{P}(F|E) \cdot \mathbb{P}(E)$, and more generally

$$\mathbb{P}(E_n \cap E_{n-1} \cap \dots \cap E_1) = \mathbb{P}(E_n | E_{n-1} \cap \dots \cap E_1) \dots \mathbb{P}(E_3 | E_2 \cap E_1) \mathbb{P}(E_2 | E_1) \mathbb{P}(E_1)$$

PROOF This follows by induction from the definition of conditional probability. \square

Ex. 1.2.4 Andrew and Bob play for the college basketball team. They get two T-shirts each, in closed bags. Any T-shirt can be black or white, with 50-50 chance. Andrew prefers black, but Bob has no preference. The following day, Andrew shows up with a black shirt on. What is the chance that Andrew's other shirt is black?

SOL'N We have $\Omega = \{(B, B), (B, W), (W, B), (W, W)\}$ which is reduced to $\{(B, B), (B, W), (W, B)\}$, so the answer is $1/3$. To make this transparent, consider

$$\begin{aligned}A_1 &= \{\text{Andrew has at least one black shirt}\} \\ A_2 &= \{\text{Both of Andrew's shirts are black}\} \\ A_3 &= \{\text{Andrew has a black shirt on}\}\end{aligned}$$

so in Andrew's case, $A_1 = A_3$ and $\mathbb{P}(A_2 | A_3) = \mathbb{P}(A_2 | A_1)$.

Ex. 1.2.5 (Polya's Urn) Initially, we have two balls, 1 red, 1 blue, in the urn. For the first draw, pick one, check its color, and put it back and put another ball of the same color into the urn.

1. What is $\mathbb{P}(\text{the first three balls are red, blue, red (in this order)})$.

SOL'N 1. Let R_i, B_i denote the i^{th} draw is red or blue respectively. Then

$$\mathbb{P}(R_3 \cap B_2 \cap R_1) = \mathbb{P}(R_3 | B_2 \cap R_1) \mathbb{P}(B_2 | R_1) \mathbb{P}(R_1) = \frac{1}{2} \frac{1}{3} \frac{1}{2} = \frac{1}{12}$$

Ex. 1.2.6 What is $\mathbb{P}(\text{in bridge, each of the players gets one ace})$?

SOL'N Write

$$\begin{aligned}
 &E_4 \\
 &\cap \\
 &E_3 = \{\text{Aces of spaces, hearts, and diamonds are at 3 different players.}\} \\
 &\cap \\
 &E_2 = \{\text{Aces of spaces, hearts, and diamonds are at 2 different players.}\} \\
 &\cap \\
 &E_1 = \Omega
 \end{aligned}$$

so that $\mathbb{P}(E_4) = \mathbb{P}(E_4 \cap E_3 \cap E_2 \cap E_1) = \mathbb{P}(E_4|E_3)\mathbb{P}(E_3|E_2)\mathbb{P}(E_2|E_1)\mathbb{P}(E_1)$.

1.2.2 Bayes' Formula

Ex. 1.2.7 Consider an insurance company, which classifies people into accident prone drivers (30%) and non-accident-prone drivers, (70%). For accident prone drivers, the chance of being involved in an accident within a year is 0.2, while for non-accident-prone drivers, the chance of being involved in an accident is 0.1. Now suppose we have a new policyholder.

1. What is the probability that the policyholder is involved in an accident within a year?
2. The policyholder was involved in an accident?

SOL'N 1. $B = \{\text{accident in 2018}\}$, $A = \{\text{the policyholder is accident prone}\}$. Then

$$\mathbb{P}(B) = \mathbb{P}(B \cap A) + \mathbb{P}(B \cap A^c) = \mathbb{P}(B|A)\mathbb{P}(A) + \mathbb{P}(B|A^c)\mathbb{P}(A^c) = 0.2 \cdot 0.3 + 0.1 \cdot 0.7 = 0.13$$

2. Now

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B|A)\mathbb{P}(A) + \mathbb{P}(B|A^c)\mathbb{P}(A^c)} = \frac{0.2 \cdot 0.3}{0.13} = \frac{6}{13}$$

Prop. 1.2.8 Suppose $A_1, A_2, \dots, A_n \in \mathcal{F}$ form a partition of Ω . Given such a partition, for any $B \in \mathcal{F}$,

$$\mathbb{P}(B) = \sum_{i=1}^n \mathbb{P}(B \cap A_i) = \sum_{i=1}^n \mathbb{P}(B|A_i) \cdot \mathbb{P}(A_i)$$

Then for any $k \in [n]$,

$$\mathbb{P}(A_k|B) = \frac{\mathbb{P}(B \cap A_k)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A_k) \cdot \mathbb{P}(A_k)}{\sum_{i=1}^n \mathbb{P}(B|A_i) \cdot \mathbb{P}(A_i)}$$

Ex. 1.2.9 Roll a fair dice. There is a urn with one white ball in it. If the die turns up 1, 3, or 5, put one black ball into the urn. If it turns up 2 or 4, put 3 black and 5 white, and if it turns up 6, put 5 black and 5 white.

SOL'N Write

$$\begin{aligned}
 A_1 &= \{1, 3 \text{ or } 5 \text{ rolled}\} \\
 A_2 &= \{2 \text{ or } 4 \text{ rolled}\} \\
 A_3 &= \{6 \text{ rolled}\}B &= \{\text{black ball rolled}\}
 \end{aligned}$$

so that

$$\begin{aligned}\mathbb{P}(A_3|B) &= \frac{\mathbb{P}(B|A_3)\mathbb{P}(A_3)}{\mathbb{P}(B|A_1) \cdot \mathbb{P}(A_1) + \mathbb{P}(B|A_2) \cdot \mathbb{P}(A_2) + \mathbb{P}(B|A_3) \cdot \mathbb{P}(A_3)} \\ &= \frac{5/6 \cdot 1/6}{1/2 \cdot 1/2 + 3/4 \cdot 1/3 + 5/6 \cdot 1/6} \\ &= \frac{5}{23}\end{aligned}$$

Ex. 1.2.10 There is a blood test for a rare but serious disease. Only 1/10000 people have this disease. Suppose the test is 100% effective, so if someone is tested ill, it is positive with 100% chance. Suppose there is also a 1% chance of false positive.

A new patient is tested, and tests positive. What are the odds that she has the disease?

Sol'n Let $A = \{\text{the person is ill}\}$ and $B = \{\text{the test is positive}\}$. Then

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B|A)\mathbb{P}(A) + \mathbb{P}(B|A^c)\mathbb{P}(A^c)} = \frac{1 \cdot 0.0001}{1 \cdot 0.0001 + 0.01 \cdot 0.9999}$$

Ex. 1.2.11 (Monty Hall paradox) There are three doors: one of them hides a prize, and two hide nothing. Pick a door. The announcer then reveals another door not containing a prize. Is it better to stay or switch?

Sol'n Write $A_i = \{\text{door } i \text{ hides the prize}\}$, and $B_2 = \{\text{door 2 is opened}\}$. Then

$$\mathbb{P}(A_1|B_2) = \frac{\mathbb{P}(B_2|A_1)\mathbb{P}(A_1)}{\mathbb{P}(B_2|A_1)\mathbb{P}(A_1) + \mathbb{P}(B_2|A_2)\mathbb{P}(A_2) + \mathbb{P}(B_2|A_3)\mathbb{P}(A_3)} = \frac{1/2 \cdot 1/3}{1/2 \cdot 1/3 + 0 + 1/3 \cdot 1/3}$$

but

$$\mathbb{P}(A_3|B_2) = \frac{\mathbb{P}(B_2|A_3)\mathbb{P}(A_3)}{\mathbb{P}(B_2|A_1)\mathbb{P}(A_1) + \mathbb{P}(B_2|A_2)\mathbb{P}(A_2) + \mathbb{P}(B_2|A_3)\mathbb{P}(A_3)} = \frac{1 \cdot 1/3}{1/2 \cdot 1/3 + 0 + 1/3 \cdot 1/3}$$

so it is better to switch!

Ex. 1.2.12 There is an inspection, which is 60% sure of the guilt of a certain suspect. The suspect is left-handed. There is new evidence: the criminal is left handed. Say 20% of the population is left handed; how certain should the inspector now be?

Sol'n Write $C = \{\text{the suspect is the criminal}\}$ and $C^c = \{\text{the criminal is someone else}\}$. Then $\mathbb{P}(C) = 0.6$ and $\mathbb{P}(C^c) = 0.4$. Let $L = \{\text{the criminal is left-handed}\}$. Then

$$\mathbb{P}(C|L) = \frac{\mathbb{P}(L|C)\mathbb{P}(C)}{\mathbb{P}(L)} \quad \mathbb{P}(C^c|L) = \frac{\mathbb{P}(L|C^c)\mathbb{P}(C^c)}{\mathbb{P}(L)}$$

Here, we can compute the “odds”:

$$\frac{\mathbb{P}(C|L)}{\mathbb{P}(C^c|L)} = \frac{\mathbb{P}(L|C)\mathbb{P}(C)}{\mathbb{P}(L|C^c)\mathbb{P}(C^c)}$$

Now $\mathbb{P}(L|C) = 1$, but $\mathbb{P}(L|C^c) = \mathbb{P}(L) = 0.2$, since the probability is taken a priori. Now a priori, the odds are given by $\mathbb{P}(C)/\mathbb{P}(C^c) = 0.6/0.4$, scaled by the factor $\mathbb{P}(L|C)/\mathbb{P}(L|C^c) = 5$ given updated information. Thus $\mathbb{P}(C|L) = 15/17$.

1.3 Independent Events

1.3.1 Definitions

Def'n. 1.3.1 The events A and B are **independent** if $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.

Ex. 1.3.2 Draw a card from a deck of 52. Let

$$A = \{\text{it is a spade}\}, \quad B = \{\text{it is an ace}\}, \quad C = \{\text{it is a heart}\}$$

We have

$$\mathbb{P}(A) = \frac{1}{4}, \quad \mathbb{P}(B) = \frac{1}{13}, \quad \mathbb{P}(A \cap B) = \frac{1}{52}$$

so A and B are independent. Similarly, B and C are independent. However, $\mathbb{P}(A \cap C) = 0 \neq 1/4$ so A and C are not independent.

Rmk. 1.3.3 Exclusive events are quite different than independence: in fact, they are (in a sense) the opposite. Let $\mathbb{P}(A) > 0$. Then A and B are independent iff $\mathbb{P}(B|A) = \mathbb{P}(B)$. Similarly, A and B are exclusive iff $\mathbb{P}(B|A) = 0$.

Ex. 1.3.4 Roll two fair dice, the yellow and the white die. Then

$$\begin{aligned} A &= \{\text{the sum is 7}\} \\ B &= \{\text{the sum is 10}\} \\ C &= \{\text{the yellow die turns up 6}\} \\ D &= \{\text{the white die turns up 6}\} \end{aligned}$$

We have $\mathbb{P}(A) = 1/6$, $\mathbb{P}(C) = 1/6$. Then $\mathbb{P}(A \cap C) = 1/36 = 1/6 \cdot 1/6$ so A and C are independent. Similarly, C and D are independent and A and D are independent. Thus A, C, D are pairwise independent but not independent as a triple.

Def'n. 1.3.5 The events A_1, A_2, \dots are **independent (as a collection)** if, for any choice of indices $1 \leq i_1 < i_2 < \dots < i_k \leq n$, then

$$\mathbb{P}(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = \mathbb{P}(A_{i_1})\mathbb{P}(A_{i_2}) \dots \mathbb{P}(A_{i_k})$$

1.3.2 Independent Trials

We have two parameters: $n \geq 1$, which is the number of trials, and $p \in (0, 1)$, which is the chance of success for an individual trial. Then $A_k = \{\text{the } k^{\text{th}} \text{ trial is a success}\}$ so that $\mathbb{P}(A_k) = p$ and the events A_1, \dots, A_n are independent. Our framework is to consider the space $\Omega \times \Omega \times \dots \times \Omega$.

Ex. 1.3.6 Roll a fair die 10 times. Then $A_k = \{\text{the } k^{\text{th}} \text{ roll is a 6}\}$. Then we have

- $\mathbb{P}(\text{all } n \text{ trials are successful}) = \mathbb{P}(A_1 \cap \dots \cap A_n) = p^n$
- $\mathbb{P}(\text{there is at least one success}) = 1 - (1 - p)^n$
- $\mathbb{P}(\text{there are exactly } k \text{ success out of } n \text{ trials}) = \binom{n}{k} p^k (1 - p)^{n-k}$

Consider the case now where n is countable (infinite number of trials). Let $S = \{\text{all trials are successful}\}$ define and $S_n = \{\text{the first } n \text{ trials are successful}\}$. Then $S = \bigcap_{n=1}^{\infty} S_n$ so

$$\mathbb{P}(S) = \lim_{n \rightarrow \infty} \mathbb{P}(S_n) = \lim_{n \rightarrow \infty} p^n = 0$$

Ex. 1.3.7 Repeatedly roll two fair dice until the sum is either 5 or 7. What is the probability that the sum is 5 when we stop?

Let $A_i = \{\text{rolls less than } i \text{ are not 5 or 7, roll } i \text{ is 5}\}$. Since $\mathbb{P}(\text{roll is 5 or 7}) = 1/6 + 1/9$, we have $\mathbb{P}(\text{roll is not}) = 13/18$. Thus

$$\mathbb{P}(A_i) = \left(\frac{13}{18}\right)^{i-1} \frac{5}{18}$$

so that

$$\mathbb{P}(A) = \frac{1}{9} \sum_{i=0}^{\infty} \left(\frac{13}{18}\right)^i = \frac{1}{9} \frac{1}{1 - \frac{13}{18}} = \frac{2}{5}$$

We have an alternate solution: note that A_1, B_1, C_1 partition the sample space. By the law of total probability,

$$\begin{aligned} \mathbb{P}(D) &= \mathbb{P}(D|A_1)\mathbb{P}(A_1) + \mathbb{P}(D|A_2)\mathbb{P}(A_2) + \mathbb{P}(D|C_1)\mathbb{P}(C_1) \\ &= \mathbb{P}(B_1) + \mathbb{P}(C_1)\mathbb{P}(D) \end{aligned}$$

so that

$$\mathbb{P}(D) = \frac{\mathbb{P}(B_1)}{1 - \mathbb{P}(C_1)} = \frac{\mathbb{P}(B_1)}{\mathbb{P}(A_1) + \mathbb{P}(B_1)}$$

1.3.3 Random Walks

We first see the gambling interpretation. Suppose we have two players, A has initial capital k and B has initial capital $N - k$. At each round, a coin is flipped. If it is a head, then B gives A 1 dollar, and if it is a tail, A gives B 1 dollar. Repeat this until someone runs out of money.



Let $\mathbb{P}_k^{(N)} = \mathbb{P}(\text{when starting at position } j, \text{ the probability that eventually } A \text{ wins})$. We have $P_0 = 0, P_N = 1$. Then for $1 \leq k \leq N - 1$, we have

$$P_k = \mathbb{P}\{\text{ending at } N \text{ when starting at } k | \text{first flip is H}\} \cdot \frac{1}{2} + \mathbb{P}\{\text{end at } N \text{ if start at } k | \text{first flip is T}\} \cdot \frac{1}{2}$$

which can be written

$$\mathbb{P}_k = P_{k+1} \frac{1}{2} + P_{k-1} \frac{1}{2} \Rightarrow \frac{1}{2} (P_k - P_{k-1}) = \frac{1}{2} (P_{k+1} - P_k)$$

so, for any $1 \leq k \leq N$, $P_k - P_{k-1} = d$ and

$$1 = P_N - P_0 = P_n - P_{N-1} + P_{N-1} - P_{N-2} + \cdots + (P_1 - P_0) = N \cdot d$$

so $d = 1/N$ and

$$P_k = P_k - P_0 = \sum_{j=1}^k (P_j - P_{j-1}) = kd = \frac{k}{N}$$