

Spatial Statistical Modeling of Wildfire Incidence in Arizona

STAT574E | Group 3

Raymond Owino, Alex Salce, Matthew Wallace

December 2024

Executive Summary

The *Wildland Fire Incident Locations* dataset we selected for study (provided by the National Interagency Fire Center) contains spatial coordinates of a wildfire's origin and its resulting size in acreage (among several other useful attributes) for all wildfire incidence in the US as archived in the IRWIN system since 2014. Our research goals included the study of factors that may influence the resulting size of a wildfire, and to build spatial models that may aid as a tool for wildfire risk assessment and resource allocation, forecasting, and other possible real-world applications. We were also interested whether wildfire incidence, when treated as a point pattern, occurs as a completely spatially random process, or if there really is inhomogeneity in spatial intensity for incidence and thus truly higher risk in some spatial regions than others.

Attributes contained in the dataset allowed us to take three different approaches to spatial statistical modeling. Using the incident size as a continuous response, we built a spatial linear model that predicts wildfire size based on spatial and selected covariate inputs. Filtering the data using incident size as well as other incident attributes also allowed us to treat wildfire incidence data as a point process. We fit a Log-Gaussian Cox Process (LGCP) spatial model to study spatial intensity, as well as a binary response spatial logistic regression model to map spatially the regions of highest probability of large wildfire (≥ 1000 acres) incidence in the region of study.

Our modeling efforts studied Coconino County (CC) in Northern Arizona, which has the highest overall wildfire incidence as compared to the rest of the state. We gathered data for incidence proximity to roads, environmental, and census data corresponding to spatial coordinates to use as covariates for our models to support our research questions. These covariates improved performance of our models by AIC metrics. **Our LGCP model indicated that wildfires are not CSR, and there is thus greater risk of wildfire incidence in certain regions of CC.** Our models require some further refinements, but serve as a solid foundation and proof of concept for our real-world application goals.

Introduction

Wildfire incidence data describes the spatial origin of a wildfire, and is one of many ways that wildfires can be studied; it is also a type of data that lends itself well to be studied by methods in spatial statistical analyses. Spatial statistical models can be valuable in real applications for wildfire prevention, giving inferential insights into wildfire risk as well as tools to aid in resource allocation for mitigation efforts and planning. Our project seeks to build models that could be useful for these kinds of efforts, and our research questions aim to provide insight to what kinds of environmental and human factors can influence the size of a resulting wildfire based upon where it originates spatially.

Wildfire Incidence Data

Accompanying this paper is `wfigs_az_sf_EPSG32612.RData` which imports `wfigs_az_sf` to the R environment, an `sf` object with 18089 observations of wildfire incidence in the state of Arizona as well as corresponding covariate data for each entry (to be discussed later). This data was originally acquired via Wildland Fire Incident Locations

from the National Interagency Fire Center. It contains spatial point data indicating the origin of each wildfire recorded in the IRWIN database (since 2014), and includes many useful data attributes for each entry. The attributes from this dataset that we used were

- x and y | Spatial coordinates in lat/lon
- IncidentSize | Size of the resulting wildfire in acres
- FireCause | Human, Natural, Unknown, Undetermined
- FireDiscoveryDateTime | Date & time of incident reporting
- IncidentTypeCategory | WF (wildfire) or RX (prescribed burn)

Other covariate data that was captured for each incidence point will be discussed later in the report.

Research Questions

Our research questions were driven by our curiosities about the ways in which we could approach modeling the data. We were also motivated in part by our findings during our explorations, and we wanted to evaluate whether real data we felt could influence wildfire size could serve as useful covariate information for our models. Additionally, we wanted to evaluate spatial randomness for different types of incidence to be sure what we are investigating is more than a completely spatially random process. In general, we found this to be a particularly interesting dataset because it seemed as though there was more than one way to approach its analysis with tools we have learned in this course.

In what ways can we approach spatial modeling of this data to produce useful insights? [RQ1]

Our overall goal is to find models that could support wildfire risk assessment, tools for resource allocation, and prediction for wildfire size for forecasting. This got us thinking about how we may approach our data in different ways. During our data explorations, we rationalized different ways we could model the dataset. We determined that we could use the data to build the following models that we will detail throughout the report.

- Spatial Linear Model - used to predict wildfire size for input coordinates and conditions found to affect size.
- Log-Gaussian Cox Process (LGCP) - used to determine whether a type of wildfire (e.g. large human caused fires) have characteristics of spatial randomness, or if they are not random, and we can study where higher-risk areas of incidence are spatially.
- Binary response GLM logistic regression - model to be used to assess risk in terms of the model's chosen risk factors, with outputs expressed as probabilities of the incidence in question spatially.

Can we find useful covariate data that can improve our models? [RQ2]

Based on combinations of findings from data explorations and our intuition about what could influence wildfire size, we chose to collect covariate data for the following factors.

- Proximity to roads - Is a fire's size influenced by how accessible it is via roads? Do more remote fires tend to get bigger?
- Environmental factors - can factors like temperature and precipitation influence wildfire size? Can terrain factors like forest and grass coverage, or local steepness influence?
- Population density - do areas of more concentrated human settlement influence wildfire incidence?

Are the patterns of human or non human caused fires spatially CSR, or do they exhibit an inhomogeneous spatial intensity? [RQ3]

An extension of the LGCP model, we wanted to investigate whether certain types of fires occur spatially randomly, or if they exhibit characteristics to the contrary? For example, we may find large human caused fires are not spatially random, and a risk model from an LGCP fit could help determine higher-risk areas of such incidence spatially.

Exploratory Data Analysis

Wildfire Incidence, Coconino County

The full set of incidence data for Arizona contained 18089 total observations, 10174 of which Human caused, 5409 Naturally caused, and 2210 unknown In effort to keep our data manageable, we opted to discard unknown cause data, and filter down to Coconino County, which has the highest count of wildfire incidence in the state with 3824 incidents.

Proximity to roads

.

Environmental factors

.

Population density

.

Statistical Analyses

Fixed, Continuous Response

Spatial Linear Model

Point Process

Log-Gaussian Cox Process

Binary Response GLM Logistic Regression

Conclusions