

```
/* import data */
PROC IMPORT DATAFILE="/home/u63048916/STAT571B/Homework/Homework 3/Q4-40.csv"
  DBMS=CSV
  OUT=mont4_40
  REPLACE;
  GETNAMES=YES;
RUN;

title 'ANOVA - missing data';
proc glm data=mont4_40;
class Car Additive;
model Mileage = Additive Car;
lsmeans Additive / alpha=0.05 adjust=tukey;
output out=diag r=res p=pred;
run;

/* check normality */
proc univariate data=diag normal;
var res;
qqplot res / normal (mu=est sigma=est);
Run;

/* check outliers */
data outlier;
set diag;
stdres=res/0.954257;
run;

proc print data=outlier;
run;

/* check constant variance using graph*/
title 'residual plot: res vs predicted value ';
proc sgplot data=diag;
scatter x=pred y=res;
refline 0;
run;

title 'residual plot: res vs Additive ';
proc sgplot data=diag;
scatter x=Additive y=res;
refline 0;
run;

title 'residual plot: res vs Car ';
proc sgplot data=diag;
scatter x=Car y=res;
refline 0;
run;

title 'Post ANOVA';
proc glm data=mont4_40;
class Car Additive;
model Mileage = Additive Car;
lsmeans Additive / tdiff pdiff alpha=0.05 adjust=bon;
lsmeans Additive / pdiff alpha=0.05 adjust=tukey;
output out=diag r=res p=pred;
run;
```

sis of these data given from Problem 4.17.



**4.40.** An engineer is studying the mileage performance characteristics of five types of gasoline additives. In the road test he wishes to use cars as blocks; however, because of a

time constraint, he must use an incomplete block design. He runs the balanced design with the five blocks that follow. Analyze the data from this experiment (use  $\alpha = 0.05$ ) and draw conclusions.

Additive	Car				
	1	2	3	4	5
1		17	14	13	12
2	14	14		13	10
3	12		13	12	9
4	13	11	11	12	
5	11	12	10		8