

A de-novo RNA biomarker of ICB response that does not survive independent replication




Fig 1. HOOK — de-novo antigen axis separates response in Gide (AUROC 0.91) but collapses on held-out Riaz (0.40)

Internal claim tested on WES-derived proxies: co-variation is burden-family only, not RNA-phenotype




Fig 2. MECHANISM — WES-proxy phenotypes share no low-rank structure beyond mutational burden (perm p=0.78)

Co-variation test is well-powered for a latent RNA-state at proxy resolution




Fig 3. POWER — co-variation test well-powered ($\sim 23\%$ detectable) vs observed $\sim 2\%$: wrong instrument, not underpowered

Pooled cross-validated ROC, ICB response baselines




Fig 4. THE BAR — approved biomarkers reproduce the modest ceiling (TMB/clinical 0.62, TIDE 0.53)

WES-derived neoantigen proxies add no signal over TMB




Fig 5. EXTERNAL NULL — no WES-proxy phenotype adds incremental AUROC over the TMB floor

Raw-read de-novo transcriptome organizes ICB response (Gide 2019 pilot, n=12)




Fig 6. RAW-READ CANDIDATE — de-novo antigen program separates responders in a balanced 12-sample pilot

Why the de-novo antigen axis is infiltration-driven (n=40)





Fig 7. WHY IT FAILS — antigen axis is infiltration-driven ($\rho=0.77$; every antigen gene 0.54-0.79 infiltration-correlated)




Fig 8. TEARDOWN — antigen axis fails to transfer to BOTH held-out cohorts (Gide 0.87 → Riaz 0.36, Hugo 0.58)