

MTHE 212 — Linear Algebra

WINTER 2026

Based on lectures by A. Shaltut – Queen's University

Notes written by Alex Lévesque

These notes are my own interpretations of the course material and they are not endorsed by the lecturers.

Feel free to reach out if you point out any errors.

Contents

1	Preface	2
2	Quiz 2 Practice 1	3
2.1	Topic 1: Direct Sums & Dimension Formulas	3
2.2	Topic 2: Definitions & Basic Proofs	3
2.3	Topic 3: Null Space, Range, & Rank-Nullity	4
3	Quiz 2 Practice 2	5
3.1	Topic 1: Direct Sums & Dimension Formulas	5
3.2	Topic 2: Definitions & Basic Proofs	5
3.3	Topic 3: Null Space, Range & Subspace Equality	6

1 Preface

Grading Scheme:

Textbook:

Comments:

-

2 Quiz 2 Practice 1

2.1 Topic 1: Direct Sums & Dimension Formulas

Focus: Equivalence of $U \cap W = \{0\}$ and direct sums; applying $\dim(U + W) = \dim U + \dim W - \dim(U \cap W)$.

Question 1.1 (Concrete Application):

Consider the vector space $\mathcal{P}_3(\mathbb{R})$ of polynomials with degree at most 3. Let U and W be the following subspaces:

$$U = \{p(x) \in \mathcal{P}_3(\mathbb{R}) \mid p(0) = 0 \text{ and } p'(0) = 0\}$$

$$W = \{ax + b \mid a, b \in \mathbb{R}\}$$

- (a) Find the dimension of U and the dimension of W .
- (b) Determine the dimension of $U \cap W$.
- (c) Use the dimension formula to find $\dim(U + W)$. Is the sum $U + W$ a direct sum? Explain why or why not using the appropriate equivalence.

Question 1.2 (Abstract/Theoretical):

Let V be a finite-dimensional vector space with $\dim V = 10$. Let U and W be subspaces of V such that $\dim U = 6$ and $\dim W = 5$.

- (a) Prove that $U \cap W \neq \{0\}$.
- (b) What are the possible values for $\dim(U \cap W)$?
- (c) Prove that there does not exist a subspace Z such that $U \oplus Z = V$ and $W \oplus Z = V$ simultaneously (Hint: Use dimensions).

2.2 Topic 2: Definitions & Basic Proofs

Focus: Basis, Linear Independence, Span, Injectivity, Surjectivity, Isomorphisms.

Question 2.1 (Linear Independence & Injectivity):

Let V and W be vector spaces and let $T \in \mathcal{L}(V, W)$ be an injective linear map.

Prove that if v_1, \dots, v_n is a linearly independent list of vectors in V , then the list Tv_1, \dots, Tv_n is linearly independent in W .

(Note: This is a classic “follow the definitions” proof that Axler emphasizes).

Question 2.2 (Surjectivity & Spanning):

Let V and W be finite-dimensional vector spaces and $T \in \mathcal{L}(V, W)$.

Prove that T is surjective if and only if T maps a spanning list of V to a spanning list of W .

(Specifically: Show that if $\text{span}(v_1, \dots, v_n) = V$, then $\text{span}(Tv_1, \dots, Tv_n) = \text{range}(T)$).

2.3 Topic 3: Null Space, Range, & Rank-Nullity

Focus: Direct sums involving null/range, proving subspace equality ($A \subseteq B$ and $B \subseteq A$), Rank-Nullity Theorem.

Question 3.1 (Subspace Equality via Double Inclusion):

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear map defined by $T(x, y, z) = (x - y, y - z, z - x)$.

- (a) Find a basis for $\text{null}(T)$.
- (b) Let $S = \text{span}((1, 1, 1))$. Prove that $\text{null}(T) = S$ by showing that $S \subseteq \text{null}(T)$ and $\text{null}(T) \subseteq S$.

Question 3.2 (Rank-Nullity & Direct Sums):

Let V be a finite-dimensional vector space and $T \in \mathcal{L}(V, V)$ be a linear operator.

- (a) State the Rank-Nullity Theorem (Fundamental Theorem of Linear Maps).
- (b) Suppose that $\text{range}(T) \cap \text{null}(T) = \{0\}$. Prove that $V = \text{range}(T) \oplus \text{null}(T)$.

(Hint: Use the Rank-Nullity theorem to show that the dimension of the sum equals the dimension of V , then use the property of direct sums).

3 Quiz 2 Practice 2

3.1 Topic 1: Direct Sums & Dimension Formulas

Prof's Hint: “Understand the equivalence between the intersection being zero and the sum being direct.”

Question 1.1 (Concrete Intersection - Practice for the Exam):

Consider the vector space \mathbb{R}^4 . Let U and W be the following subspaces:

$$U = \{(x, y, z, w) \in \mathbb{R}^4 \mid x + y = 0 \text{ and } z = 2w\}$$

$$W = \text{span}((1, -1, 0, 0), (0, 0, 1, 1))$$

- (a) Find a basis for U and state $\dim U$.
- (b) Find a basis for W and state $\dim W$.
- (c) Find a basis for $U \cap W$. (Hint: Set a general vector in W equal to a general vector in U , or plug the generic vector of W into the equations for U).
- (d) Is $U + W$ a direct sum? Explain using the intersection result.

Question 1.2 (The Equivalence Proof):

Let U and W be subspaces of a finite-dimensional vector space V .

Prove that $U + W$ is a direct sum if and only if $\dim(U + W) = \dim U + \dim W$.

(Note: This directly tests the “equivalence” your professor mentioned. You will likely need to use the full dimension formula $\dim(U + W) = \dim U + \dim W - \dim(U \cap W)$ to prove this).

3.2 Topic 2: Definitions & Basic Proofs

Prof's Hint: “Comfortable proving basic facts that follow directly from these definitions (injective, surjective, basis).”

Question 2.1 (Isomorphisms & Basis):

Let V and W be finite-dimensional vector spaces and let $T \in \mathcal{L}(V, W)$ be an isomorphism (injective and surjective).

Prove that if (v_1, \dots, v_n) is a basis of V , then (Tv_1, \dots, Tv_n) is a basis of W .

(Hint: You must prove two things: that the list is linearly independent (uses injectivity) and that it spans W (uses surjectivity).)

Question 2.2 (Injectivity via Null Space):

Let $T \in \mathcal{L}(V, W)$. Prove that T is injective if and only if $\text{null}(T) = \{0\}$.

*(Note: This is a fundamental “definition check” in Axler. You need to prove the implication in both directions:

1. If T is injective, then only 0 maps to 0.
 2. If the only thing that maps to 0 is 0, then $T(u) = T(v) \Rightarrow u = v$.)*
-

3.3 Topic 3: Null Space, Range & Subspace Equality

Prof’s Hint: “Prove that two subspaces are equal ($A = B$) by showing $A \subseteq B$ and $B \subseteq A$.”

Question 3.1 (Subspace Equality Challenge):

Let $T : \mathcal{P}_2(\mathbb{R}) \rightarrow \mathcal{P}_3(\mathbb{R})$ be the linear map defined by $T(p(x)) = x \cdot p(x)$.

Let S be the subspace defined by $S = \{q(x) \in \mathcal{P}_3(\mathbb{R}) \mid q(0) = 0\}$.

- (a) Find the range of T .
- (b) Prove that $\text{range}(T) = S$ using the double inclusion method.

(i.e., Show every polynomial in the range has a zero constant term, and show every polynomial with a zero constant term is in the range).

Question 3.2 (Rank-Nullity & Intersection):

Let V be a finite-dimensional vector space and let $T \in \mathcal{L}(V, V)$.

Suppose that $\text{range}(T) = \text{null}(T)$.

- (a) Prove that $\dim V$ must be an even number.
- (b) Give an example of such a map T on \mathbb{R}^2 .

(Hint: Use the Rank-Nullity Theorem $\dim V = \dim \text{null}(T) + \dim \text{range}(T)$ and substitute the equality given)