

## Homework 1

Name: Yun-Ping Du

## Question 1

A. Convert the following numbers to their decimal representation. Show your work.

1.  $10011011_2$ *Solution.*

$$\begin{aligned} 10011011_2 &= 2^7 + 2^4 + 2^3 + 2^1 + 2^0 \\ &= 128 + 16 + 8 + 2 + 1 \\ &= \mathbf{155} \end{aligned}$$

■

2.  $456_7$ *Solution.*

$$\begin{aligned} 456_7 &= 4 \times 7^2 + 5 \times 7^1 + 6 \times 7^0 \\ &= 196 + 35 + 6 \\ &= \mathbf{237} \end{aligned}$$

■

3.  $38A_{16}$ *Solution.*

$$\begin{aligned} 38A_{16} &= 3 \times 16^2 + 8 \times 16^1 + 10 \times 16^0 \\ &= 768 + 128 + 10 \\ &= \mathbf{906} \end{aligned}$$

■

4.  $2214_5$ *Solution.*

$$\begin{aligned} 2214_5 &= 2 \times 5^3 + 2 \times 5^2 + 1 \times 5^1 + 4 \times 5^0 \\ &= 250 + 50 + 5 + 4 \\ &= \mathbf{309} \end{aligned}$$

■

B. Convert the following numbers to their binary representation:

1.  $69_{10}$

*Solution.* Recursively divide the quotients and collect the remainders

$$\left. \begin{array}{r|l} 2)69 & 1 \\ 2)34 & 0 \\ 2)17 & 1 \\ 2)8 & 0 \\ 2)4 & 0 \\ 2)2 & 0 \\ 2)1 & 1 \end{array} \right\} \mathbf{1000101_2}$$

Thus the binary representation of  $69_{10}$  is  $\mathbf{1000101_2}$ . ■

2.  $485_{10}$

*Solution.* Recursively divide the quotients and collect the remainders

$$\left. \begin{array}{r|l} 2)485 & 1 \\ 2)242 & 0 \\ 2)121 & 1 \\ 2)60 & 0 \\ 2)30 & 0 \\ 2)15 & 1 \\ 2)7 & 1 \\ 2)3 & 1 \\ 2)1 & 1 \end{array} \right\} \mathbf{111100101_2}$$

Thus the binary representation of  $485_{10}$  is  $\mathbf{111100101_2}$ . ■

3.  $6D1A_{16}$

*Solution.* We first obtain the binary representation of each digit

$$6_{16} = 0110_2$$

$$D_{16} = 1101_2$$

$$1_{16} = 0001_2$$

$$A_{16} = 1010_2$$

Thus the binary representation of  $6D1A_{16}$  is  $\mathbf{110110100011010_2}$ . ■

C. Convert the following numbers to their hexadecimal representation:

1.  $1101011_2$

*Solution.* Since we have

$$0110_2 = 6_{16}$$

$$1011_2 = B_{16}$$

thus the hexadecimal representation of  $1101011_2$  is  **$6B_{16}$** . ■

2.  $895_{10}$

*Solution.* Recursively divide the quotients and collect the remainders

$$895 \div 16 = 55 \text{ R}15 \rightarrow F_{16}$$

$$55 \div 16 = 3 \text{ R}7 \rightarrow 7_{16}$$

$$3 \div 16 = 0 \text{ R}3 \rightarrow 3_{16}$$

Thus the hexadecimal representation of  $895_{10}$  is  **$37F_{16}$** . ■

## Question 2

Solve the following, do all calculation in the given base. Show your work.

1.  $7566_8 + 4515_8 =$

*Solution.*

$$\begin{array}{r} \phantom{00}111 \\ 7566_8 \\ + 4515_8 \\ \hline 14303_8 \end{array}$$

■

2.  $10110011_2 + 1101_2 =$

*Solution.*

$$\begin{array}{r} \phantom{0000}111111 \\ 10110011_2 \\ + 00001101_2 \\ \hline 11000000_2 \end{array}$$

■

3.  $7A66_{16} + 45C5_{16} =$

*Solution.*

$$\begin{array}{r} \phantom{00}11 \\ 7A66_{16} \\ + 45C5_{16} \\ \hline C02B_{16} \end{array}$$

■

4.  $3022_5 - 2433_5 =$

*Solution.*

$$\begin{array}{r} \phantom{00}241 \\ 3022_5 \\ - 2433_5 \\ \hline 34_5 \end{array}$$

■

### Question 3

A. Convert the following numbers to their 8-bit two's complement representation. Show your work.

1.  $124_{10}$

*Solution.* We first obtain the binary representation of  $124_{10}$

$$\left. \begin{array}{r|l} 2 \overline{)124} & 0 \\ 2 \overline{)62} & 0 \\ 2 \overline{)31} & 1 \\ 2 \overline{)15} & 1 \\ 2 \overline{)7} & 1 \\ 2 \overline{)3} & 1 \\ 2 \overline{)1} & 1 \end{array} \right\} \mathbf{1111100_2}$$

Thus the 8-bit two's complement representation of  $124_{10}$  is **01111100**. ■

2.  $-124_{10}$

*Solution.* To get the 8-bit two's complement representation of a negative number, we first "flip" the bits (swapping 0s and 1s) of its positive counterpart; the value of 1 is then added to the resulting value (ignoring the overflow). We already know that  $124_{10}$  is represented by

$$01111100$$

Flip the bits and we have

$$01111100 \rightarrow 10000011$$

Finally, add 1 to the representation and we have

$$10000011_2 + 1_2 = 10000100_2$$

Thus the 8-bit two's complement representation of  $-124_{10}$  is **10000100**. ■

3.  $109_{10}$

*Solution.* We first obtain the binary representation of  $109_{10}$

$$\left. \begin{array}{r|l} 2 \overline{)109} & 1 \\ 2 \overline{)54} & 0 \\ 2 \overline{)27} & 1 \\ 2 \overline{)13} & 1 \\ 2 \overline{)6} & 0 \\ 2 \overline{)3} & 1 \\ 2 \overline{)1} & 1 \end{array} \right\} \mathbf{1101101_2}$$

Thus the 8-bit two's complement representation of  $109_{10}$  is **01101101**. ■

4.  $-79_{10}$

*Solution.* We first obtain the binary representation of  $79_{10}$

$$\left. \begin{array}{r|l} 2)79 & 1 \\ 2)39 & 1 \\ 2)19 & 1 \\ 2)9 & 1 \\ 2)4 & 0 \\ 2)2 & 0 \\ 2)1 & 1 \end{array} \right\} \mathbf{1001111_2}$$

Flip the bits and we have

$$01001111 \rightarrow 10110000$$

Finally, add 1 to the representation and we have

$$10110000_2 + 1_2 = 10110001_2$$

Thus the 8-bit two's complement representation of  $-79_{10}$  is **10110001**. ■

B. Convert the following numbers (represented as 8-bit two's complement) to their decimal representation. Show your work.

1.  $00011110_{8\text{-bit } 2\text{'s comp}}$

*Solution.* Since the leftmost bit is 0, we directly convert the binary representation to the decimal representation

$$\begin{aligned} 11110_2 &= 2^4 + 2^3 + 2^2 + 2^1 \\ &= 16 + 8 + 4 + 2 \\ &= 30 \end{aligned}$$

Thus the decimal representation of  $00011110_{8\text{-bit } 2\text{'s comp}}$  is **30**. ■

2.  $11100110_{8\text{-bit } 2\text{'s comp}}$

*Solution.* Since the leftmost bit is 1, we know that the number is negative. To convert a negative number into its decimal representation, we first subtract 1 from the 8-bit two's complement, flip the bits of the result, and then convert the flipped bits to its decimal representation.

$$11100110_2 - 1_2 = 11100101_2$$

Flip the bits

$$11100101 \rightarrow 00011010$$

Finally we have

$$\begin{aligned} 11010_2 &= 2^4 + 2^3 + 2^1 \\ &= 16 + 8 + 2 \\ &= 26 \end{aligned}$$

Thus the decimal representation of  $11100110_{8\text{-bit } 2\text{'s comp}}$  is **-26**. ■

3. 00101101<sub>8-bit 2's comp</sub>

*Solution.* Since the leftmost bit is 0, we directly convert the binary representation to the decimal representation

$$\begin{aligned}101101_2 &= 2^5 + 2^3 + 2^2 + 2^0 \\&= 32 + 8 + 4 + 1 \\&= 45\end{aligned}$$

Thus the decimal representation of 00101101<sub>8-bit 2's comp</sub> is **45**. ■

4. 10011110<sub>8-bit 2's comp</sub>

*Solution.* Since the leftmost bit is 1, we know that the number is negative. To convert a negative number into its decimal representation, we first subtract 1 from the 8-bit two's complement, flip the bits of the result, and then convert the flipped bits to its decimal representation.

$$10011110_2 - 1_2 = 10011101_2$$

Flip the bits

$$10011101 \rightarrow 01100010$$

Finally we have

$$\begin{aligned}1100010_2 &= 2^6 + 2^5 + 2^1 \\&= 64 + 32 + 2 \\&= 98\end{aligned}$$

Thus the decimal representation of 10011110<sub>8-bit 2's comp</sub> is **-98**. ■

## Question 4

Solve the following questions from the Discrete Math zyBook:

1. Exercise 1.2.4

(b) *Solution.*

$p$	$q$	$\neg(p \vee q)$
T	T	F
T	F	F
F	T	F
F	F	T

(c) *Solution.*

$p$	$q$	$r$	$r \vee (p \wedge \neg q)$
T	T	T	T
T	T	F	F
T	F	T	T
T	F	F	T
F	T	T	T
F	T	F	F
F	F	T	T
F	F	F	F

2. Exercise 1.3.4

(b) *Solution.*

$p$	$q$	$(p \rightarrow q) \rightarrow (q \rightarrow p)$
T	T	T
T	F	T
F	T	F
F	F	T

(d) *Solution.*

$p$	$q$	$(p \leftrightarrow q) \oplus (p \leftrightarrow \neg q)$
T	T	T
T	F	T
F	T	T
F	F	T



## Question 5

Solve the following questions from the Discrete Math zyBook:

1. Exercise 1.2.7

(b) *Solution.*  $(B \wedge D) \vee (B \wedge M) \vee (D \wedge M)$  ■

(c) *Solution.*  $B \vee (D \wedge M)$  ■

2. Exercise 1.3.7

(b) *Solution.*  $(s \vee v) \rightarrow p$  ■

(c) *Solution.*  $p \rightarrow y$  ■

(d) *Solution.*  $p \leftrightarrow (s \wedge y)$  ■

(e) *Solution.*  $p \rightarrow (s \vee y)$  ■

3. Exercise 1.3.9

(c) *Solution.*  $c \rightarrow p$  ■

(d) *Solution.*  $p \rightarrow c$  ■

## Question 6

Solve the following questions from the Discrete Math zyBook:

1. Exercise 1.3.6

(b) *Solution.* If Joe is eligible for the honors program, then he maintains a B average. ■

(c) *Solution.* If Rajiv can go on the roller coaster, then he is at least four feet tall. ■

(d) *Solution.* If Rajiv is at least four feet tall, he can go on the roller coaster. ■

2. Exercise 1.3.10 ( $p$ : T,  $q$ : False,  $r$ : unknown)

(c)  $(p \vee r) \leftrightarrow (q \wedge r)$

*Solution.* False.  $(p \vee r)$  is true because  $p$  is true.  $(q \wedge r)$  is false because  $q$  is false. Thus the biconditional proposition is false regardless of the truth value of  $r$ . ■

(d)  $(p \wedge r) \leftrightarrow (q \wedge r)$

*Solution.* Unknown. Since  $q$  is false,  $(q \wedge r)$  is false. However, the truth value of  $(p \wedge r)$  depends on the truth value of  $r$ . ■

(e)  $p \rightarrow (r \vee q)$

*Solution.* Unknown. The hypothesis is true, but the truth value of the conclusion depends on the truth value of  $r$ . ■

(f)  $(p \wedge q) \rightarrow r$

*Solution.* True. The hypothesis is false, thus the conditional proposition is true regardless of the truth value of  $r$ . ■

## Question 7

Solve Exercise 1.4.5, sections b – d, from the Discrete Math zyBook:

- (b)
- If Sally did not get the job, then she was late for interview or did not update her resume.
  - If Sally updated her resume and was not late for her interview, then she got the job.

*Solution.* Logically equivalent.

- $\neg j \rightarrow (l \vee \neg r)$
- $(r \wedge \neg l) \rightarrow j$

$j$	$l$	$r$	$\neg j \rightarrow (l \vee \neg r)$	$(r \wedge \neg l) \rightarrow j$
T	T	T	T	T
T	T	F	T	T
T	F	T	T	T
T	F	F	T	T
F	T	T	T	T
F	T	F	T	T
F	F	T	F	F
F	F	F	T	T

■

- (c)
- If Sally got the job then she was not late for her interview.
  - If Sally did not get the job, then she was late for her interview.

*Solution.* Not logically equivalent.

- $j \rightarrow \neg l$
- $\neg j \rightarrow l$

$j$	$l$	$j \rightarrow \neg l$	$\neg j \rightarrow l$
T	T	F	T
T	F	T	T
F	T	T	T
F	F	T	F

■

- (d)
- If Sally updated her resume or she was not late for her interview, then she got the job.
  - If Sally got the job, then she updated her resume and was not late for her interview.

*Solution.* Not logically equivalent.

- $(r \vee \neg l) \rightarrow j$
- $j \rightarrow (r \wedge \neg l)$

$j$	$l$	$r$	$(r \vee \neg l) \rightarrow j$	$j \rightarrow (r \wedge \neg l)$
T	T	T	T	F
T	T	F	T	F
T	F	T	T	T
T	F	F	T	F
F	T	T	F	T
F	T	F	T	T
F	F	T	F	T
F	F	F	F	T

■

## Question 8

Solve the following questions from the Discrete Math zyBook:

### 1. Exercise 1.5.2

(c)  $(p \rightarrow q) \wedge (p \rightarrow r) \equiv p \rightarrow (q \wedge r)$

*Solution.*

$$\begin{aligned}(p \rightarrow q) \wedge (p \rightarrow r) &\equiv (\neg p \vee q) \wedge (\neg p \vee r) \\ &\equiv \neg p \vee (q \wedge r) \\ &\equiv p \rightarrow (q \wedge r)\end{aligned}$$

■

(f)  $\neg(p \vee (\neg p \wedge q)) \equiv \neg p \wedge \neg q$

*Solution.*

$$\begin{aligned}\neg(p \vee (\neg p \wedge q)) &\equiv \neg((p \vee \neg p) \wedge (p \vee q)) \\ &\equiv \neg(T \wedge (p \vee q)) \\ &\equiv \neg(p \vee q) \\ &\equiv \neg p \wedge \neg q\end{aligned}$$

■

(i)  $(p \wedge q) \rightarrow r \equiv (p \wedge \neg r) \rightarrow \neg q$

*Solution.*

$$\begin{aligned}(p \wedge q) \rightarrow r &\equiv \neg(p \wedge q) \vee r \\ &\equiv (\neg p \vee \neg q) \vee r \\ &\equiv \neg p \vee r \vee \neg q \\ &\equiv \neg(p \wedge \neg r) \vee \neg q \\ &\equiv (p \wedge \neg r) \rightarrow \neg q\end{aligned}$$

■

### 2. Exercise 1.5.3

(c)  $\neg r \wedge (\neg r \rightarrow p)$

*Solution.*

$$\begin{aligned}\neg r \wedge (\neg r \rightarrow p) &\equiv \neg r \vee (\neg(\neg r) \vee p) \\ &\equiv \neg r \vee (r \vee p) \\ &\equiv (\neg r \vee r) \vee p \\ &\equiv T \vee p \\ &\equiv T\end{aligned}$$

■

(d)  $\neg(p \rightarrow q) \rightarrow \neg q$

*Solution.*

$$\begin{aligned}\neg(p \rightarrow q) \rightarrow \neg q &\equiv \neg(\neg p \vee q) \rightarrow \neg q \\ &\equiv \neg\neg(\neg p \vee q) \vee \neg q \\ &\equiv \neg p \vee q \vee \neg q \\ &\equiv \neg p \vee \mathbf{T} \\ &\equiv \mathbf{T}\end{aligned}$$



## Question 9

Solve the following questions from the Discrete Math zyBook:

1. Exercise 1.6.3

(c) *Solution.*  $\exists x (x = x^2)$



(d) *Solution.*  $\forall x (x \leq x^2)$



2. Exercise 1.7.4

(b) *Solution.*  $\forall x (\neg S(x) \wedge W(x))$



(c) *Solution.*  $\forall x (S(x) \rightarrow \neg W(x))$



(d) *Solution.*  $\exists x (S(x) \wedge W(x))$



## Question 10

Solve the following questions from the Discrete Math zyBook:

1. Exercise 1.7.9

	$P(x)$	$Q(x)$	$R(x)$
$a$	T	T	F
$b$	T	F	F
$c$	F	T	F
$d$	T	T	F
$e$	T	T	T

(c)  $\exists x ((x = c) \rightarrow P(x))$

*Solution.* True. Example:  $a$ .

■

(d)  $\exists x (Q(x) \wedge R(x))$

*Solution.* True. Example:  $e$ .

■

(e)  $Q(a) \wedge P(d)$

*Solution.* True.

■

(f)  $\forall x ((x \neq b) \rightarrow Q(x))$

*Solution.* True.

■

(g)  $\forall x (P(x) \vee R(x))$

*Solution.* False. Counterexample:  $c$ .

■

(h)  $\forall x (R(x) \rightarrow P(x))$

*Solution.* True.

■

(i)  $\exists x (Q(x) \vee R(x))$

*Solution.* True. Example:  $a$ .

■



2. Exercise 1.9.2

$P$	1	2	3	$Q$	1	2	3	$S$	1	2	3
1	T	F	T	1	F	F	F	1	F	F	F
2	T	F	T	2	T	T	T	2	F	F	F
3	T	T	F	3	T	F	F	3	F	F	F

(b)  $\exists x \forall y Q(x, y)$

*Solution.* True. Let  $x = 2$ , then  $Q(x, 1)$ ,  $Q(x, 2)$ , and  $Q(x, 3)$  are all true. ■

(c)  $\exists x \forall y P(y, x)$

*Solution.* True. Let  $x = 1$ , then  $P(1, x)$ ,  $P(2, x)$ , and  $P(3, x)$  are all true. ■

(d)  $\exists x \exists y S(x, y)$

*Solution.* False. There is no pair  $(x, y)$  such that  $S(x, y)$  is true. ■

(e)  $\forall x \exists y Q(x, y)$

*Solution.* False. Let  $x = 1$ , then there is no  $y$  such that  $Q(1, y)$  is true. ■

(f)  $\forall x \exists y P(x, y)$

*Solution.* True. When  $x = 1$ , let  $y = 1$ . When  $x = 2$ , let  $y = 1$ . When  $x = 3$ , let  $y = 1$ . ■

(g)  $\forall x \forall y P(x, y)$

*Solution.* False. Counterexample:  $(x, y) = (1, 2)$ . ■

(h)  $\exists x \exists y Q(x, y)$

*Solution.* True. Example:  $(x, y) = (2, 1)$ . ■

(i)  $\forall x \forall y \neg S(x, y)$

*Solution.* True. ■

## Question 11

Solve the following questions from the Discrete Math zyBook:

### 1. Exercise 1.10.4

- (c) There are two numbers whose sum is equal to their product.

*Solution.*  $\exists x \exists y (x + y = xy)$  ■

- (d) The ratio of every two positive numbers is also positive.

*Solution.*  $\forall x \forall y ((x > 0 \wedge y > 0) \rightarrow x/y > 0)$  ■

- (e) The reciprocal of every positive number less than one is greater than one.

*Solution.*  $\forall x ((x > 0) \wedge (x < 1) \rightarrow 1/x > 1)$  ■

- (f) There is no smallest number.

*Solution.*  $\forall x \exists y (x > y)$  ■

- (g) Every number besides 0 has a multiplicative inverse.

*Solution.*  $\forall x \exists y ((x \neq 0) \rightarrow xy = 1)$  ■

### 2. Exercise 1.10.7

- $P(x, y)$ :  $x$  knows  $y$ 's phone number. (A person may or may not know their own phone number.)
- $D(x)$ :  $x$  missed the deadline.
- $N(x)$ :  $x$  is a new employee.

- (c) There is at least one new employee who missed the deadline.

*Solution.*  $\exists x (N(x) \wedge D(x))$  ■

- (d) Sam knows the phone number of everyone who missed the deadline.

*Solution.*  $\forall x (D(x) \rightarrow P(\text{Sam}, x))$  ■

- (e) There is a new employee who knows everyone's phone number.

*Solution.*  $\exists x \forall y (N(x) \wedge P(x, y))$  ■

- (f) Exactly one new employee missed the deadline.

*Solution.*  $\exists x \forall y ((N(x) \wedge D(x)) \wedge (((y \neq x) \wedge N(y)) \rightarrow \neg D(y)))$  ■

3. Exercise 1.10.10

- (c) Every student has taken at least one class besides Math 101.

*Solution.*  $\forall x \exists y ((y \neq \text{Math 101}) \wedge T(x, y))$  ■

- (d) There is a student who has taken every math class besides Math 101.

*Solution.*  $\exists x \forall y ((y \neq \text{Math 101}) \rightarrow T(x, y))$  ■

- (e) Everyone besides Sam has taken at least two different math classes.

*Solution.*  $\forall x \exists y \exists z ((x \neq \text{Sam}) \rightarrow ((y \neq z) \wedge T(x, y) \wedge T(x, z)))$  ■

- (f) Sam has taken exactly two math classes.

*Solution.*  $\exists x \exists y \forall z ((x \neq y) \wedge T(\text{Sam}, x) \wedge T(\text{Sam}, y) \wedge (((z \neq x) \wedge (z \neq y)) \rightarrow \neg T(\text{Sam}, z)))$  ■

## Question 12

Solve the following questions from the Discrete Math zyBook:

1. Exercise 1.8.2

- $P(x)$ :  $x$  was given the placebo
- $D(x)$ :  $x$  was given the medication
- $M(x)$ :  $x$  had migraines

(b) Every patient was given the medication or the placebo or both.

*Solution.*

- $\forall x (D(x) \vee P(x))$
- Negation:  $\neg \forall x (D(x) \vee P(x))$
- Applying De Morgan's law:  $\exists x (\neg D(x) \wedge \neg P(x))$
- English: There is a patient who was not given the medication and the placebo. ■

(c) There is a patient who took the medication and had migraines.

*Solution.*

- $\exists x (D(x) \wedge M(x))$
- Negation:  $\neg \exists x (D(x) \wedge M(x))$
- Applying De Morgan's law:  $\forall x (\neg D(x) \vee \neg M(x))$
- English: Every patient either did not took the medication or did not have migraines (or both). ■

(d) Every patient who took the placebo had migraines.

*Solution.*

- $\forall x (P(x) \rightarrow M(x))$
- Negation:  $\neg \forall x (P(x) \rightarrow M(x))$
- Applying De Morgan's law:  $\exists x (P(x) \wedge \neg M(x))$
- English: There is a patient who took the placebo and did not have migraines. ■

(e) There is a patient who had migraines and was given the placebo.

*Solution.*

- $\exists x (M(x) \wedge P(x))$
- Negation:  $\neg \exists x (M(x) \wedge P(x))$
- Applying De Morgan's law:  $\forall x (\neg M(x) \vee \neg P(x))$
- English: Every patient either did not had migraines or was not given the placebo (or both).

■

## 2. Exercise 1.9.4

(c)  $\exists x \forall y (P(x, y) \rightarrow Q(x, y))$

*Solution.*

$$\begin{aligned} \neg \exists x \forall y (P(x, y) \rightarrow Q(x, y)) &\equiv \forall x \exists y \neg (\neg P(x, y) \vee Q(x, y)) \\ &\equiv \forall x \exists y (P(x, y) \wedge \neg Q(x, y)) \end{aligned}$$

■

(d)  $\exists x \forall y (P(x, y) \leftrightarrow P(y, x))$

*Solution.*

$$\begin{aligned} \neg \exists x \forall y (P(x, y) \leftrightarrow P(y, x)) &\equiv \forall x \exists y \neg ((P(x, y) \rightarrow P(y, x)) \wedge (P(y, x) \rightarrow P(x, y))) \\ &\equiv \forall x \exists y \neg ((\neg P(x, y) \vee P(y, x)) \wedge (\neg P(y, x) \vee P(x, y))) \\ &\equiv \forall x \exists y (\neg(\neg P(x, y) \vee P(y, x)) \vee \neg(\neg P(y, x) \vee P(x, y))) \\ &\equiv \forall x \exists y ((P(x, y) \wedge \neg P(y, x)) \vee (P(y, x) \wedge \neg P(x, y))) \end{aligned}$$

■

(e)  $\exists x \exists y P(x, y) \wedge \forall x \forall y Q(x, y)$

*Solution.*

$$\begin{aligned} \neg(\exists x \exists y P(x, y) \wedge \forall x \forall y Q(x, y)) &\equiv \neg \exists x \exists y P(x, y) \vee \neg \forall x \forall y Q(x, y) \\ &\equiv \forall x \forall y \neg P(x, y) \vee \exists x \exists y \neg Q(x, y) \end{aligned}$$

■