Figure 10. Access and Quality

Rank (highest to lowest)	1	2	3	4	5	6	7	8	9	10	11	Mean
Access, %												
Able to get same- or next-day appointment[a]	NLD 77	Australia 67	UK 57	France 56	Germany 53	US 51	Sweden 49	Canada 43	CHE NA	Denmark NA	Japan NA	57 (n = 8)
2-mo Wait time to see specialist	Canada 39	UK 19	Sweden 19	Australia 13	CHE 9	NLD 7	US 6	France 4	Germany 3	Denmark NA	Japan NA	13 (n = 9)
Adequate time with regular (primary) physician	Germany 88	UK 86	NLD 85	CHE 84	Australia 83	US 81	Canada 79	Sweden 78	France NA	Denmark NA	Japan NA	83 (n = 8)
Perceptions, %												
System works well	Germany 60	CHE 58	France 54	UK 44	Sweden 44	Australia 44	Canada 35	US 19	NLD NA	Denmark NA	Japan NA	45 (n = 8)
Fundamental changes needed	Canada 55	US 53	UK 46	Sweden 46	Australia 46	France 41	Germany 37	CHE 37	NLD NA	Denmark NA	Japan NA	45 (n = 8)
Complete rebuild of health system needed	US 23	Sweden 10	Canada 9	UK 7	France 4	Australia 4	Germany 3	CHE 3	NLD NA	Denmark NA	Japan NA	8 (n = 8)
Prevention												
Measles immunization, % of children	Sweden 98	Japan 98	Germany 97	NLD 96	UK 93	CHE 93	Australia 93	US 92	France 91	Denmark 91	Canada 90	94
Breast cancer screening, % of women aged 50-69 y[b]	Denmark 84	US 81	NLD 79	UK 76	Sweden 75	Canada 72	Germany 71	Australia 55	France 52	CHE 47	Japan 41	67
Clinical outcomes												
30-d Stroke mortality per 1000 patients[c]	Canada 10.0	Sweden 9.6	Australia 9.3	UK 9.2	France 7.9	CHE 6.9	Germany 6.4	US 4.2	NLD NA	Denmark NA	Japan NA	7.9 (n = 8)
30-d Mortality per 1000 patients with acute myocardial infarction	Germany 8.7	Sweden 8.3	CHE 7.7	UK 7.6	France 7.2	Canada 6.7	US 5.5	Australia 4.1	NLD NA	Denmark NA	Japan NA	7.0 (n = 8)
Foreign body left per 100 000 discharges	CHE 12.3	Canada 8.6	Australia 8.6	France 6.2	UK 6.1	Germany 5.5	Sweden 4.6	US 4.1	NLD NA	Denmark NA	Japan NA	7.0 (n = 8)
Obstetric trauma without instrument per 100 deliveries	Canada 3.1	UK 2.8	Sweden 2.8	CHE 2.6	Denmark 2.6	NLD 2.5	Australia 2.4	Germany 2.1	US 1.5	France 0.6	Japan NA	2.3 (n = 10)
Avoidable hospitalizations												
Diabetes hospitalizations per 100 000 population[d]	Germany 218.3	US 191.0	Japan 162.3	France 150.6	Australia 141.1	Denmark 113.4	Sweden 96.0	Canada 93.7	UK 72.8	CHE 72.6	NLD 69.8	125.6
Diabetes hospitalizations as a ratio of population with diabetes[e]	Japan 2.80	Australia 2.80	Germany 2.40	US 2.00	Sweden 1.90	Denmark 1.80	UK 1.70	Canada 1.30	France 1.20	NLD 1.20	CHE 1.20	1.85
Asthma hospitalizations per 100 000 population[f]	US 89.7	UK 71.0	Australia 64.8	Denmark 50.6	NLD 36.0	Japan 34.7	France 29.6	Germany 28.7	CHE 27.5	Sweden 19.0	Canada 14.6	42.4
Asthma hospitalizations as a ratio of population with asthma[g]	US 1.20	UK 1.00	France 0.80	Denmark 0.80	Germany 0.70	NLD 0.70	Australia 0.60	CHE 0.40	Sweden 0.30	Japan 0.30	Canada 0.20	0.64

NA indicates not applicable. CHE indicates Switzerland; NLD, the Netherlands. See eTable 6 in Supplement 2 for data ordered by country.

Mean is the unweighted arithmetic mean of the countries with data; (n = x) under a mean gives the number of countries contributing when data were not available (NA) for all 11.

[a] Able to get same- or next-day appointment when sick, excluding those who did not need to see a physician or nurse.

[b] Women aged 40 to 49 years in Sweden.

[c] Thirty-day stroke mortality after hospital admission for ischemic stroke.

[d] Limited to a primary diagnosis of diabetes.

[e] Diabetes hospitalizations are limited to persons aged 15 years or older, whereas the denominator (population with diabetes) is both adult and pediatric.

[f] Limited to a primary diagnosis of asthma.

[g] Asthma hospitalizations are limited to persons aged 15 years or older, whereas the denominator (population with asthma) is both adult and pediatric.