

The Relative Eigenvector and Pure Tensor Problems

Alex Ryba,
Queens College, CUNY

Groups, Nilpotence and Tensors 2023
Colorado State University, 04/29/2023

Relative Eigenvectors

Relative Eigenvectors

Let A and B be matrices with the same size. A relative eigenvector v with relative eigenvalue λ satisfies $vB = \lambda vA$.

Relative Eigenvectors

Let A and B be matrices with the same size. A relative eigenvector v with relative eigenvalue λ satisfies $vB = \lambda vA$.

Problem (Relative Eigenvectors)

Find the relative eigenvectors and the corresponding relative eigenspaces for matrices A and B .

Relative Eigenvectors

Let A and B be matrices with the same size. A relative eigenvector v with relative eigenvalue λ satisfies $vB = \lambda vA$.

Problem (Relative Eigenvectors)

Find the relative eigenvectors and the corresponding relative eigenspaces for matrices A and B .

The Eigenvector Problem: B is square and $A = I$.

Relative Eigenvectors

Let A and B be matrices with the same size. A relative eigenvector v with relative eigenvalue λ satisfies $vB = \lambda vA$.

Problem (Relative Eigenvectors)

Find the relative eigenvectors and the corresponding relative eigenspaces for matrices A and B .

The Eigenvector Problem: B is square and $A = I$.

Definition

If matrices A, B over k have same size, v is a relative eigenvector if vA and vB lie in a 1-dimensional space.

The difference is that we add the nullspace of A to the set of relative eigenvectors.

Simultaneous Eigenvectors

Simultaneous Eigenvectors

Definition

A *simultaneous eigenvector* v for A_1, A_2, \dots, A_n satisfies:

$$vA_1 = \lambda_1 v, \quad vA_2 = \lambda_2 v, \quad \dots, \quad vA_n = \lambda_n v.$$

Simultaneous Eigenvectors

Definition

A simultaneous eigenvector v for A_1, A_2, \dots, A_n satisfies:

$$vA_1 = \lambda_1 v, \quad vA_2 = \lambda_2 v, \quad \dots, \quad vA_n = \lambda_n v.$$

Alternative: $vI, vA_1, vA_2, \dots, vA_n$ span a 1-dimensional space.

Simultaneous Eigenvectors

Definition

A simultaneous eigenvector v for A_1, A_2, \dots, A_n satisfies:

$$vA_1 = \lambda_1 v, \quad vA_2 = \lambda_2 v, \quad \dots, \quad vA_n = \lambda_n v.$$

Alternative: $vI, vA_1, vA_2, \dots, vA_n$ span a 1-dimensional space.

Problem (Relative Eigenvectors)

Given $a_i : X \rightarrow Y$, $1 \leq i \leq n$, find all vectors x such that xa_1, xa_2, \dots, xa_n lie in a 1-dimensional space.

Simultaneous Eigenvectors

Definition

A simultaneous eigenvector v for A_1, A_2, \dots, A_n satisfies:

$$vA_1 = \lambda_1 v, \quad vA_2 = \lambda_2 v, \quad \dots, \quad vA_n = \lambda_n v.$$

Alternative: $vI, vA_1, vA_2, \dots, vA_n$ span a 1-dimensional space.

Problem (Relative Eigenvectors)

Given $a_i : X \rightarrow Y$, $1 \leq i \leq n$, find all vectors x such that xa_1, xa_2, \dots, xa_n lie in a 1-dimensional space.

Alternative: there exist scalars $\lambda_1, \lambda_2, \dots, \lambda_n$ and a vector y with:

$$xa_1 = \lambda_1 y, \quad xa_2 = \lambda_2 y, \quad \dots, \quad xa_n = \lambda_n y.$$

Simultaneous Eigenvectors

Definition

A simultaneous eigenvector v for A_1, A_2, \dots, A_n satisfies:

$$vA_1 = \lambda_1 v, \quad vA_2 = \lambda_2 v, \quad \dots, \quad vA_n = \lambda_n v.$$

Alternative: $v, vA_1, vA_2, \dots, vA_n$ span a 1-dimensional space.

Problem (Relative Eigenvectors)

Given $a_i : X \rightarrow Y$, $1 \leq i \leq n$, find all vectors x such that xa_1, xa_2, \dots, xa_n lie in a 1-dimensional space.

Alternative: there exist scalars $\lambda_1, \lambda_2, \dots, \lambda_n$ and a vector y with:

$$xa_1 = \lambda_1 y, \quad xa_2 = \lambda_2 y, \quad \dots, \quad xa_n = \lambda_n y.$$

If a_1 has a right inverse r , this reduces to a simultaneous eigenvector problem. Here, $yr = \frac{xa_1 r}{\lambda_1} = \frac{x}{\lambda_1}$. Hence, x is a simultaneous eigenvector for the maps $a_2 r, a_3 r, \dots, a_n r$.

Pure Tensors

Pure Tensors

A typical element in $Y \otimes Z$ is a finite sum $\sum y_i \otimes z_i$, where $y_i \in Y, z_i \in Z$.

Pure Tensors

A typical element in $Y \otimes Z$ is a finite sum $\sum y_i \otimes z_i$, where $y_i \in Y, z_i \in Z$.

Definition

A pure tensor in $Y \otimes Z$ is a special element $y \otimes z$.

Pure Tensors

A typical element in $Y \otimes Z$ is a finite sum $\sum y_i \otimes z_i$, where $y_i \in Y, z_i \in Z$.

Definition

A pure tensor in $Y \otimes Z$ is a special element $y \otimes z$.

Problem (Pure Tensors)

Given k -vector spaces Y and Z and $X \leq Y \otimes Z$. Find all pure tensors in $X \otimes \bar{k}$.

Pure Tensors

A typical element in $Y \otimes Z$ is a finite sum $\sum y_i \otimes z_i$, where $y_i \in Y, z_i \in Z$.

Definition

A pure tensor in $Y \otimes Z$ is a special element $y \otimes z$.

Problem (Pure Tensors)

Given k -vector spaces Y and Z and $X \leq Y \otimes Z$. Find all pure tensors in $X \otimes \bar{k}$.

Let b_1, \dots, b_n be a basis of the dual space of Z . We have maps:

$$a_i = 1 \otimes b_i : Y \otimes Z \rightarrow Y \otimes k \cong Y.$$

A pure tensor $x = y \otimes z$ is a relative eigenvector for the maps a_1, \dots, a_n . Because images $xa_i = (a_i(z))y$ are proportional. Conversely, if an element of $Y \otimes Z$ has proportional images under these maps then it is a pure tensor.

Group Theoretic Applications

Group Theoretic Applications

In a recent J. Alg. paper, I prove that an absolutely irreducible kG -module V either has a tensor decomposition $V \cong U \otimes W$ or is induced iff $A = \text{Hom}_k(V, V)$ has a proper G -invariant subalgebra B .

Group Theoretic Applications

In a recent J. Alg. paper, I prove that an absolutely irreducible kG -module V either has a tensor decomposition $V \cong U \otimes W$ or is induced iff $A = \text{Hom}_k(V, V)$ has a proper G -invariant subalgebra B .

Then the G -invariant algebra $C = C_A(B)$ is non-trivial.

Group Theoretic Applications

In a recent J. Alg. paper, I prove that an absolutely irreducible kG -module V either has a tensor decomposition $V \cong U \otimes W$ or is induced iff $A = \text{Hom}_k(V, V)$ has a proper G -invariant subalgebra B .

Then the G -invariant algebra $C = C_A(B)$ is non-trivial.

B and C contain commuting non-trivial irreducible G -submodules I and J .

Group Theoretic Applications

In a recent J. Alg. paper, I prove that an absolutely irreducible kG -module V either has a tensor decomposition $V \cong U \otimes W$ or is induced iff $A = \text{Hom}_k(V, V)$ has a proper G -invariant subalgebra B .

Then the G -invariant algebra $C = C_A(B)$ is non-trivial.

B and C contain commuting non-trivial irreducible G -submodules I and J .

Options for I and J are parameterized by $v_I \in \bar{k}^m$, $v_J \in \bar{k}^n$, where m and n are the multiplicities of I and J in the module $\text{Soc}(A)$.

Group Theoretic Applications

In a recent J. Alg. paper, I prove that an absolutely irreducible kG -module V either has a tensor decomposition $V \cong U \otimes W$ or is induced iff $A = \text{Hom}_k(V, V)$ has a proper G -invariant subalgebra B .

Then the G -invariant algebra $C = C_A(B)$ is non-trivial.

B and C contain commuting non-trivial irreducible G -submodules I and J .

Options for I and J are parameterized by $v_I \in \bar{k}^m$, $v_J \in \bar{k}^n$, where m and n are the multiplicities of I and J in the module $\text{Soc}(A)$.

I and J commute iff $v_I \otimes v_J$ lies in a computable subspace $X \leq k^m \otimes k^n$, a Pure Tensor Problem.

Group Theoretic Applications

In a recent J. Alg. paper, I prove that an absolutely irreducible kG -module V either has a tensor decomposition $V \cong U \otimes W$ or is induced iff $A = \text{Hom}_k(V, V)$ has a proper G -invariant subalgebra B .

Then the G -invariant algebra $C = C_A(B)$ is non-trivial.

B and C contain commuting non-trivial irreducible G -submodules I and J .

Options for I and J are parameterized by $v_I \in \bar{k}^m$, $v_J \in \bar{k}^n$, where m and n are the multiplicities of I and J in the module $\text{Soc}(A)$.

I and J commute iff $v_I \otimes v_J$ lies in a computable subspace $X \leq k^m \otimes k^n$, a Pure Tensor Problem.

A similar approach detects embeddings of finite groups into exceptional groups of Lie type.

Gröbner Basis solves the Pure Tensor Problem

Gröbner Basis solves the Pure Tensor Problem

The Pure Tensor Problem: Identify pure tensors in $X \leq Y \otimes Z$.

Gröbner Basis solves the Pure Tensor Problem

The Pure Tensor Problem: Identify pure tensors in $X \leq Y \otimes Z$.

Let $\dim(X) = l$, $\dim(Y) = m$, $\dim(Z) = n$, and pick b_Y, b_Z as bases of Y and Z .

Gröbner Basis solves the Pure Tensor Problem

The Pure Tensor Problem: Identify pure tensors in $X \leq Y \otimes Z$.

Let $\dim(X) = l, \dim(Y) = m, \dim(Z) = n$, and pick b_Y, b_Z as bases of Y and Z .

An $l \times mn$ matrix M_x gives a basis of X relative to $b_Y \otimes b_Z$.

Gröbner Basis solves the Pure Tensor Problem

The Pure Tensor Problem: Identify pure tensors in $X \leq Y \otimes Z$.

Let $\dim(X) = l$, $\dim(Y) = m$, $\dim(Z) = n$, and pick b_Y, b_Z as bases of Y and Z .

An $l \times mn$ matrix M_X gives a basis of X relative to $b_Y \otimes b_Z$.

The column nullspace N_X of M_X is an $nm \times (nm - l)$ matrix with: $M_X N_X = 0$.

Gröbner Basis solves the Pure Tensor Problem

The Pure Tensor Problem: Identify pure tensors in $X \leq Y \otimes Z$.

Let $\dim(X) = l, \dim(Y) = m, \dim(Z) = n$, and pick b_Y, b_Z as bases of Y and Z .

An $l \times mn$ matrix M_X gives a basis of X relative to $b_Y \otimes b_Z$.

The column nullspace N_X of M_X is an $nm \times (nm - l)$ matrix with: $M_X N_X = 0$.

The pure tensors in $X \otimes \bar{k}$ satisfy:

$$((y_1, y_2, \dots, y_m) \otimes (z_1, z_2, \dots, z_n)) N_X = 0.$$

Gröbner Basis solves the Pure Tensor Problem

The Pure Tensor Problem: Identify pure tensors in $X \leq Y \otimes Z$.

Let $\dim(X) = l, \dim(Y) = m, \dim(Z) = n$, and pick b_Y, b_Z as bases of Y and Z .

An $l \times mn$ matrix M_X gives a basis of X relative to $b_Y \otimes b_Z$.

The column nullspace N_X of M_X is an $nm \times (nm - l)$ matrix with: $M_X N_X = 0$.

The pure tensors in $X \otimes \bar{k}$ satisfy:

$$((y_1, y_2, \dots, y_m) \otimes (z_1, z_2, \dots, z_n)) N_X = 0.$$

$mn - l$ (quadratic) polynomial conditions on $n + m$ unknowns.

Linear Method

The Relative Eigenvector Problem for 2 maps: Given a and $b : X \rightarrow Y$, compute the relative eigenvectors with $vb = \lambda va$.

The Relative Eigenvector Problem for 2 maps: Given a and $b : X \rightarrow Y$, compute the relative eigenvectors with $vb = \lambda va$.

A (linear) recursion: In the base case, map a has a right inverse r . We have the eigenvector problem for the map br .

Otherwise, write N for the non-zero nullspace of a and M for its image Nb . Both a and b map N to M , so there are induced maps $[a], [b] : X/N \rightarrow Y/M$. A relative eigenvector x for a and b maps to a relative eigenvector $x + N$ for $[a]$ and $[b]$. The following lemma shows that relative eigenvectors lift back from $[a]$ and $[b]$ to a and b .

Otherwise, write N for the non-zero nullspace of a and M for its image Nb . Both a and b map N to M , so there are induced maps $[a], [b] : X/N \rightarrow Y/M$. A relative eigenvector x for a and b maps to a relative eigenvector $x + N$ for $[a]$ and $[b]$. The following lemma shows that relative eigenvectors lift back from $[a]$ and $[b]$ to a and b .

Lemma

Suppose that $x + N$ is a relative eigenvector for $[a]$ and $[b]$ with $(x + N)[b] = \lambda(x + N)[a]$. Then the set of lifting vectors $x^\dagger \in x + N \subset X$ for which $x^\dagger b = \lambda x^\dagger a$ is a (non-empty) coset of $N(a) \cap N(b)$.

Otherwise, write N for the non-zero nullspace of a and M for its image Nb . Both a and b map N to M , so there are induced maps $[a], [b] : X/N \rightarrow Y/M$. A relative eigenvector x for a and b maps to a relative eigenvector $x + N$ for $[a]$ and $[b]$. The following lemma shows that relative eigenvectors lift back from $[a]$ and $[b]$ to a and b .

Lemma

Suppose that $x + N$ is a relative eigenvector for $[a]$ and $[b]$ with $(x + N)[b] = \lambda(x + N)[a]$. Then the set of lifting vectors $x^\dagger \in x + N \subset X$ for which $x^\dagger b = \lambda x^\dagger a$ is a (non-empty) coset of $N(a) \cap N(b)$.

Proof. Let x^* be any lift of $x + N$ to X . Let $y = x^*a$. We have

$$(x + N)[b] = \lambda(x + N)[a] = \lambda(y + M).$$

So, $x^*b = \lambda y + m$, with $m \in M = Nb$. Pick $n \in N$ with $nb = m$. Let $x^\dagger = x^* - n$. Then $x^\dagger a = y$ and $x^\dagger b = x^*b - nb = \lambda y + m - m = \lambda x^\dagger a$.



NP Hardness

The Relative Eigenvector Problem: Given

$a_1, a_2, \dots, a_n : X \rightarrow Y$, compute the set of vectors x such that xa_1, xa_2, \dots, xa_n is contained in a 1-space. This is NP-Hard.

The Relative Eigenvector Problem: Given

$a_1, a_2, \dots, a_n : X \rightarrow Y$, compute the set of vectors x such that xa_1, xa_2, \dots, xa_n is contained in a 1-space. This is NP-Hard.

Problem (Relative Eigenvector Decision)

Given a set of matrices with entries in the field k , does there exist a relative eigenvector whose first coordinate is 1 and whose other coordinates belong to k ?

The Relative Eigenvector Problem: Given

$a_1, a_2, \dots, a_n : X \rightarrow Y$, compute the set of vectors x such that xa_1, xa_2, \dots, xa_n is contained in a 1-space. This is NP-Hard.

Problem (Relative Eigenvector Decision)

Given a set of matrices with entries in the field k , does there exist a relative eigenvector whose first coordinate is 1 and whose other coordinates belong to k ?

This clearly belongs to NP. To prove that it's in NPC we can construct a reduction from:

Problem (CNF SAT)

Is there a truth assignment that satisfies a boolean formula given in conjunctive normal form (CNF)?

Problem (CNF SAT)

Is there a truth assignment that satisfies a boolean formula given in conjunctive normal form (CNF)?

A formula in CNF is a conjunction of clauses, each of which is a disjunction of terms. The terms are selected from n variables and their negations. The following is an example with 2 clauses, 4 variables and 5 terms:

$$(X_1 \vee \overline{X_2} \vee X_3) \wedge (\overline{X_1} \vee X_4). \quad (1)$$

Problem (CNF SAT)

Is there a truth assignment that satisfies a boolean formula given in conjunctive normal form (CNF)?

A formula in CNF is a conjunction of clauses, each of which is a disjunction of terms. The terms are selected from n variables and their negations. The following is an example with 2 clauses, 4 variables and 5 terms:

$$(X_1 \vee \overline{X_2} \vee X_3) \wedge (\overline{X_1} \vee X_4). \quad (1)$$

From a formula that uses n variables, m clauses and ℓ terms we construct an instance of the Relative Eigenvector Decision Problem that uses $1 + n + \ell + m$ matrices with size $(\ell + 1)(2n + 1) \times (2n + 2)$ and entries in the field F_2 .

Questions

Is the following NPC?

Problem (Relative Eigenvector Decision)

Given a set of matrices does there exist a relative eigenvector?

Is the following NPC?

Problem (Relative Eigenvector Decision)

Given a set of matrices does there exist a relative eigenvector?

Is the following NPC? Does it lead to special relative eigenvector problems?

Problem (Tensor Decomposition)

Given an absolutely irreducible matrix group, does there exist a tensor decomposition?