

Foundations of Machine Learning — Homework Assignment 1

Anirudhan J Rajagopalan
N18824115
ajr619

October 11, 2015

D. Kernels

1

Given: Kernel, K is defined by $K(x, y) = \sum_{i=1}^N \cos^n(x_i^2 - y_i^2)$ for all $(X, Y) \in \mathbb{R}^N \times \mathbb{R}^N$

Solution: We know that

$$\cos(x_i^2 - y_i^2) = \sin(x_i^2) \cdot \sin(y_i^2) + \cos(x_i^2) \cdot \cos(y_i^2) \quad (1)$$

This can be written as a dot product of two vectors

$$\phi(x_i) = \begin{bmatrix} \cos(x_i^2) \\ \sin(x_i^2) \end{bmatrix} \quad \text{and} \quad \phi(y_i) = \begin{bmatrix} \cos(y_i^2) \\ \sin(y_i^2) \end{bmatrix} \quad (2)$$

We know that if K can be written as $\langle \phi(x_i), \phi(y_i) \rangle$, then it is a PDS@.

Also, $\langle \phi(x_i), \phi(y_i) \rangle$ is a positive scalar. When a scalar is raised to a positive power (n in our case) and summed with N other positive scalar, we get a positive scalar as our answer. Hence

$$K(x, y) = \sum_{i=1}^N \cos^n(x_i^2 - y_i^2) \text{ is PDS.}$$