

# Quantum Theory for Mathematicians

Erik Herrera

January 21, 2025

**Problem 1.** Consider a particle moving in the real line in the presence of a force coming from a potential function  $V$ . Given some value  $E_0$  for the energy of the particle, suppose that  $V(x) < E_0$  for all  $x$  in some closed interval  $[x_0, x_1]$ . Then a particle with initial position  $x_0$  and particle moving in the real line in the presence of a force coming from a potential function  $V$ . Given some value  $E_0$  for the energy of the particle, suppose that  $V(x) < E_0$  for all  $x$  in some closed interval  $[x_0, x_1]$ . Then a particle with initial position  $x_0$  and positive initial velocity will continue to move to the right until it reaches  $x_1$ . Using (2.6), show that the time needed to travel from  $x_0$  to  $x_1$  is given by

$$t = \int_{x_0}^{x_1} \sqrt{\frac{m}{2(E_0 - V(y))}} dy$$

Note: This shows that we can solve Newton's equation in  $\mathbb{R}^1$  more or less explicitly for time as a function of position, which in principle determines the position as a function of time.

*Proof.* The total energy of the above particle is given by

$$E_0 = \frac{1}{2}m\dot{x}^2 + V(x)$$

Solving for velocity as a function of position gives

$$\dot{x} = \sqrt{\frac{2(E_0 - V(x))}{m}}$$

Since  $V(x) < E_0$ , this velocity is positive on the entire interval  $[x_0, x_1]$ . By the inverse function theorem, we may conclude that the time is a function of position and moreover that

$$\frac{dt}{dx} = \sqrt{\frac{m}{2(E_0 - V(x))}}$$

The total time elapsed between  $x_0$  and  $x_1$  is then given by

$$\int_{t_0}^{t_1} dt = \int_{x_0}^{x_1} \frac{dt}{dx} dx = \int_{x_0}^{x_1} \sqrt{\frac{m}{2(E_0 - V(y))}} dy$$

■

**Problem 3.** Consider the equation of motion of a pendulum of length  $L$ ,

$$\frac{d^2\theta}{dt^2} + \frac{g}{L} \sin \theta = 0$$

where  $g$  is the acceleration of gravity. Here  $\theta$  is the angle between the pendulum and the negative  $y$ -axis in the plane. This system has a stable equilibrium at  $\theta = 0$  and an unstable equilibrium at  $\theta = \pi$ . Consider initial conditions of the form  $\theta(0) = \pi - \delta, \dot{\theta}(0) = 0$ , for  $0 < \delta < \pi/4$ . Fix some angle  $\theta_0$  and let  $T(\delta)$  denote the time it takes for the pendulum with the given initial conditions to reach the angle  $\theta_0$ . (Here  $\theta_0$  represents an arbitrarily chosen cutoff point at which the pendulum is no longer "close" to  $\theta = \pi$ .) Show that  $T(\delta)$  grows only logarithmically as  $\delta$  tends to zero. Note: Logarithmic growth of  $T$  as a function of  $\delta$  corresponds to exponential decay of  $\delta$  as a function of  $T$ . Thus, if we want  $T$  to be large, we must choose  $\delta$  to be very small.

*Proof.* The total energy for the pendulum is given by

$$E = \frac{1}{2}m \left( L \frac{d\theta}{dt} \right)^2 - mgL \cos \theta$$

This is the same as the Hamiltonian for a particle with mass  $mL^2$  and potential energy  $mgL \cos \theta$ . Therefore, by the result of problem 1, the time to travel is given by

$$T(\delta) = - \int_{\pi-\delta}^{\theta_0} \sqrt{\frac{mL^2}{2(-mgL \cos(\pi - \delta) + mgL \cos \theta)}} d\theta = \sqrt{\frac{L}{2g}} \int_{\pi-\delta}^{\theta_0} \frac{-1}{\sqrt{\cos \theta - \cos(\pi - \delta)}} d\theta$$

Rewriting this we get

$$T(\delta) = \sqrt{\frac{L}{2g}} \int_{\delta}^{\pi-\theta_0} \frac{1}{\sqrt{\cos(\delta) - \cos(\theta)}} d\theta$$

Since we are only interested in the case of small  $\delta$ , we may instead examine the approximation

$$T(\delta) \approx \sqrt{\frac{L}{2g}} \int_{\delta}^{\pi-\theta_0} \frac{1}{\sqrt{1 - \cos(\theta)}} d\theta = \sqrt{\frac{L}{4g}} \int_{\delta}^{\pi-\theta_0} \frac{1}{\sin(\theta/2)} d\theta$$

Thus, we have the approximation

$$T(\delta) \approx \sqrt{\frac{L}{g}} \left( \log \left( \tan \left( \frac{\pi - \theta_0}{4} \right) \right) - \log \left( \tan \left( \frac{\delta}{4} \right) \right) \right)$$

By using a Taylor expansion for  $\tan(\delta/4)$ , we can see that

$$T(\delta) \sim -\sqrt{\frac{L}{g}} \log(\delta)$$

This shows that  $T(\delta)$  grows logarithmically in  $\delta$  as  $\delta \rightarrow 0$ .

■

**Problem 8.** Consider a particle moving in  $\mathbb{R}^n$  with a velocity-dependent force law given by

$$\mathbf{F}(\mathbf{x}, \mathbf{v}) = -\nabla V(\mathbf{x}) + \mathbf{F}_2(\mathbf{x}, \mathbf{v})$$

where the velocity-dependent term  $\mathbf{F}_2$  acts perpendicularly to the velocity of the particle. (That is, we assume that  $\mathbf{v} \cdot \mathbf{F}_2(\mathbf{x}, \mathbf{v}) = 0$  for all  $\mathbf{x}$  and  $\mathbf{v}$ .) Let  $E$  denote the usual energy function  $E(\mathbf{x}, \mathbf{v}) = \frac{1}{2}m|\mathbf{v}|^2 + V(\mathbf{x})$ , unmodified by the presence of the velocity-dependent term in the force. Show that  $E$  is conserved.

*Proof.* Notice that

$$\begin{aligned} \frac{dE}{dt} &= m\dot{\mathbf{x}} \cdot \ddot{\mathbf{x}} + \nabla V \cdot \dot{\mathbf{x}} \\ &= \dot{\mathbf{x}} \cdot (m\ddot{\mathbf{x}} + \nabla V \cdot \mathbf{x}) \\ &= \dot{\mathbf{x}} \cdot (\mathbf{F}(\mathbf{x}, \mathbf{v}) + \nabla V \cdot \mathbf{x}) \\ &= \dot{\mathbf{x}} \cdot \mathbf{F}_2(\mathbf{x}, \mathbf{v}) \end{aligned}$$

If  $\mathbf{F}_2(\mathbf{x}, \mathbf{v})$  is always orthogonal to  $\dot{\mathbf{x}}$ , then  $\frac{dE}{dt} = 0$ .

■

**Problem 18.** Determine the Hamiltonian flow on  $\mathbb{R}^2$  generated by the function  $f(x, p) = xp$ .

*Proof.* We calculate the gradient of the Hamiltonian to be

$$\nabla f(x, p) = (p, x)$$

By Hamilton's equations, we then have

$$\dot{x} = p, \quad \dot{p} = -x$$

These have the unique solutions

$$x(t) = x_0 e^t, \quad p(t) = p_0 e^{-t}$$

■