

3A Exercise 2

Suppose $f_1 = e^{-|x|}$, $f_2 = e^{-\frac{|x|}{2}}$, $f_3 = e^{-\frac{|x|}{3}}$, \dots

Then we have

$$\lim_{k \rightarrow \infty} f_k(x) = \lim_{k \rightarrow \infty} e^{-\frac{|x|}{k}} = 0$$

Now we calculate the limit of the integral:

$$\lim_{k \rightarrow \infty} \int f_k \, d\lambda = \lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} e^{-\frac{|x|}{k}} \, dx = \lim_{k \rightarrow \infty} 2 \int_0^{\infty} e^{-\frac{x}{k}} \, dx = \infty$$

3A Exercise 3

WLOG suppose $b \geq a$:

$$\begin{aligned}
 |g(a) - g(b)| &= \left| \int_{(-\infty, a)} f \, d\lambda - \int_{(-\infty, b)} f \, d\lambda \right| \\
 &= \left| \int_{(-\infty, a)} f \, d\lambda - \int_{(-\infty, a)} f \, d\lambda - \int_{(a, b)} f \, d\lambda \right| \\
 &= \left| \int_{(a, b)} f \, d\lambda \right| \\
 &\leq \lambda((a, b)) \sup_{(a, b)} |f|
 \end{aligned}$$

where the last inequality holds by [3.25 of Measure, Integration & Real Analysis](#).

Since $f \in \mathcal{L}^1(\lambda)$, then we can assume $\sup_{(a, b)} |f| = M < \infty$.

Then to have $|g(a) - g(b)| < \epsilon$, we must have $(b - a)M < \epsilon$.

Therefore, for fixed $\epsilon > 0$ it suffices that we choose $\delta > \frac{\epsilon}{M}$ to have $|g(a) - g(b)| < \epsilon$ for all $a, b \in \mathbb{R}$ such that $|b - a| < \delta$. And by doing so, g is uniformly continuous on \mathbb{R} .

3A Exercise 5

Let $f_1 = f\chi_{[-1,1]}$, $f_2 = f\chi_{[-2,2]}$, $f_3 = f\chi_{[-3,3]}$, \dots be sequence of Borel measurable functions. We can see that

$$\lim_{k \rightarrow \infty} f_k(x) = f(x)$$

for any $x \in X$.

Also $|f|$ is Borel measurable function from X to $[0, \infty]$ satisfying two conditions:

- $\int |f| d\lambda < \infty$
- $|f_k(x)| \leq |f(x)|$

for arbitrary $k \in \mathbb{Z}^+$ and any $x \in X$.

By using Dominated Convergence Theorem we have:

$$\lim_{k \rightarrow \infty} \int f_k d\lambda = \int f d\lambda$$

3A Exercise 12

Let $f_1 = \frac{(1-x)\cos(x^{-1})}{\sqrt{x}}$, $f_2 = \frac{(1-x)^2\cos(x^{-2})}{\sqrt{x}}$, $f_3 = \frac{(1-x)^3\cos(x^{-3})}{\sqrt{x}}$, \dots be sequence of measurable functions from $(0, 1)$ to $[-\infty, \infty]$. We can see that

$$\lim_{k \rightarrow \infty} f_k(x) = 0$$

for any $x \in (0, 1)$.

Now let $g_j(x) = \frac{1}{\sqrt{x}}\chi_{(\frac{1}{j}, 1)}$, where g_j is a non-negative increasing sequence of functions from $(0, 1)$ to $[0, \infty]$. We also have

$$g(x) = \lim_{j \rightarrow \infty} g_j(x)$$

by using Monotone Convergence Theorem we have

$$\lim_{j \rightarrow \infty} \int_{(0,1)} g_j d\mu = \int_{(0,1)} g d\mu$$

Suppose $g(x) = \frac{1}{\sqrt{x}}$. Then g is a Borel measurable function from $(0, 1)$ to $[0, \infty]$ satisfying two conditions:

- $\int_0^1 |g| d\mu = \int_{(0,1)} g d\mu = \lim_{j \rightarrow \infty} \int_{(0,1)} g_j d\mu = \lim_{j \rightarrow \infty} \int_0^1 g_j = \lim_{j \rightarrow \infty} \int_0^{\frac{1}{j}} x^{-1/2} dx = \lim_{j \rightarrow \infty} 2 - 2\sqrt{\frac{1}{j}} = 2 < \infty$
- $|f_k(x)| \leq |g(x)|$

for arbitrary $k \in \mathbb{Z}^+$ and any $x \in X$.

By using Dominated Convergence Theorem we have:

$$\lim_{k \rightarrow \infty} \int_{(0,1)} f_k d\lambda = \int_{(0,1)} 0 d\lambda = 0$$