



Managing Heart Attacks and Strokes

Every Minute Matters — Quick Recognition and Requesting Immediate Help

"Emergencies that require quickly recognizing the signs, requesting help immediately, and reassuring the person"



Why Do We Care About This Topic?

Delaying help can increase the severity of the condition



**Early
Recognition**
of the Warning Signs



Reducing Delay
in Requesting Help




**Preventing
Mistakes**
Such as Giving
Medications



**Reassuring the
Person**
and Organizing the
Area



**A Quick
Response**
to Emergencies



**A Culture of
First Aid**
in the Community

Remember: When a heart attack or stroke is suspected, don't wait for the symptoms to improve, and call 999 immediately



What Is Meant by a Stroke?

Blood flow to part of the brain being affected



**Sudden
weakness in
the limbs**



**Difficulty
speaking**



**Loss of balance
or dizziness**



**A sudden
change in
vision**



**A sudden
severe
headache**



**Confusion in
responsiveness**

Note: Only one sign may appear — take the case seriously and request help immediately



The FAST Rule for Recognizing a Stroke

A quick test for the most important signs

F	A	S	T
Face	Arm	Speech	Time
Face	Arm	Speech	Time
			
Drooping or weakness in one side of the face?	Difficulty raising one of the arms?	Difficulty or slurring in speech?	Call 999 immediately at any sign

Awareness Message: Don't wait for all the signs to appear — one sign may be enough to request help



Other Signs That May Be Associated with a Stroke

In addition to FAST



**A sudden loss
of balance**



**Sudden
difficulty
walking**



**Blurred or lost
vision**



**Severe,
unusual
dizziness**



**A sudden
severe
headache**



**Numbness on
one side of the
body**



**Confusion or
not
understanding**

Correct Action: Call 999 immediately, and state the symptom start time if known



What Is Meant by a Heart Attack?

Signs that may indicate cardiac symptoms requiring immediate intervention



Pain or pressure in the center of the chest

1

Pain spreading to the arm or jaw

2

Shortness of breath

3

Heavy sweating or paleness

4

Nausea or dizziness

5

Severe, unusual fatigue

6

Note: Symptoms may vary from person to person — don't wait for all the signs to appear



When Do We Call the Emergency Number 999?

Call immediately when any sign is suspected

Loss of consciousness or weak responsiveness

5

Sudden weakness in the limbs

1

A sudden severe headache

6

Difficulty speaking or understanding

2

Sudden dizziness or loss of balance

7

Pain or pressure in the chest

3

Sudden abnormal symptoms

8

Unusual shortness of breath

4

Important Message: Waiting or driving the person yourself can delay the arrival of appropriate care



What Do You Say When Calling Emergency Services?

Clear information speeds up the assessment



The precise location



The person's approximate age



The symptoms that appeared



The symptom start time



Are they conscious and breathing?



Is there chest pain?



Any known health condition

Important Message: Don't end the call unless the emergency operator asks you to



The Correct Response to a Stroke

Protecting the person from falling or choking

Monitor breathing and responsiveness

5

Reassure them: Help is coming

6

State the symptom start time

7

Call 999 immediately

1

Have them sit or lie down safely

2

No food or drink

3

No medications without guidance

4

Note: Even if the symptoms improve, the condition must not be ignored



Handling Chest Pain or a Suspected Cardiac Event

Treat it as an emergency



**Call 999
immediately**



**Sit them in a
comfortable
position**



**Encourage
them to rest**



**Reassure them
and stay with
them**



**Monitor
breathing and
responsiveness**



**No
medications
without
guidance**



**Follow the
operator's
instructions**

Important Message: Don't let the person drive a vehicle or transport themselves when serious symptoms are present



If the Person Loses Consciousness

A serious sign requiring quick action



Make sure the area is safe

1

Call 999 immediately

2

Check for normal breathing

3

No food or drink

4

Follow the operator's CPR instructions

5

Ask for an AED to be brought if available

6

Don't leave the person alone

7

Note: Loss of consciousness with difficulty breathing is a very serious emergency



Don't Give Food, Drink, or Medications

It can be dangerous if they're unable to swallow



No water or food



No medications without guidance



Don't force them to stand



Don't let them drive



Don't delay requesting help

Important Message: Your main role is requesting help, reassuring the person, and monitoring their condition



The Importance of Knowing When the Symptoms Started

Among the most important information the specialists need



When did the symptoms start?

1

When was the last time they were normal?

2

Did they appear suddenly or gradually?

3

Did they improve then return?

4

Are there any illnesses or medications?

5

Awareness Message: Recording or remembering the time can help specialists make the right decision



Handling the Person Calmly

Reassurance is part of first aid



Speak in a calm voice



Say: Help is on the way



Don't ask too many questions



Disperse the crowd



Protect their privacy



No filming or posting

Note: Calmness doesn't mean delay — call emergency services first, then reassure the person



Cases Inside the Home

The family is the first to notice the change



Don't wait for the symptoms to improve



Call 999 immediately



Clear the way for the ambulance



Prepare health information



Don't transport them yourself



Keep children away from the scene

Important Message: The family's speed in recognizing and requesting help can help reduce delay



Cases at Work or in Public Places

Organization to facilitate the arrival of help



**Someone calls
999**



**Someone
guides the
ambulance
teams**



**Provide space
and privacy**



**No crowding or
filming**



**Have the
symptom start
time ready**



**Don't move
them except in
case of danger**

Awareness Message: Dividing roles reduces chaos and helps with a quick response



Common Mistakes to Avoid

They may delay help or increase the danger



Waiting for the symptoms to disappear



Interpreting them as ordinary fatigue



Giving food or drink



Medications without guidance



Allowing them to drive a vehicle



Moving an unstable person



Crowding or filming

Important Message: Delay is the biggest mistake when a heart attack or stroke is suspected



Prevention and Reducing Risk Factors

General guidelines for a healthy lifestyle



**Monitoring
blood pressure
and sugar**



**Adhering to
medications**



**Appropriate
physical
activity**



A balanced diet



**Avoiding
smoking**



**A healthy
weight**



**Seeing the
doctor**

Note: These guidelines are general and don't replace consulting a doctor or regular health follow-up



The Role of Those Present at the Scene

Organized cooperation to coordinate the response



Someone calls
999



Someone
reassures the
person



Someone
disperses the
onlookers



Someone
clears the way



Someone
gathers the
symptom
timing



Someone
brings an AED
if needed

Important Message: Organized cooperation is better than confusion or crowding

Do's and Don'ts with Heart Attacks and Strokes

Correct Practices and Others to Avoid

Don't ✕

Don't wait for the symptoms to improve

1

No food or drink

2

No medications without guidance

3

Don't let them drive a vehicle

4

Don't crowd around the person

5

Don't film or post the case

6

Don't move them if they're unstable

7

Do ✓

Call 999 immediately

1

Use the FAST rule to recognize it

2

Help them sit safely

3

Monitor breathing and responsiveness

4

State the symptom start time

5

Reassure them and provide privacy

6

Follow the emergency operator's instructions

7

Interactive Scenarios

Practical Application of What You've Learned

Scenario 2

A person feels severe pressure in the chest and shortness of breath

Correct Action: Call 999 immediately, sit them in a comfortable position, and don't give them medications without guidance

Scenario 1

A person showed sudden drooping in the face and difficulty speaking

Correct Action: Call 999 immediately, reassure the person, and record the symptom start time

Scenario 4

People gathered around the person and one of them started filming

Correct Action: Disperse the onlookers, stop the filming, and provide privacy and space for the person

Scenario 3

The person says the symptoms are easing and wants to wait

Correct Action: Don't wait, call 999, because the symptoms easing doesn't mean the danger is gone



Test Your Knowledge

Quick Questions to Assess Your Understanding

Q1: What does the FAST rule mean for a stroke?

Answer: Face, Arm, Speech, and Time — checking the signs and immediately calling 999



Q2: Do we wait for the symptoms to improve when a stroke is suspected?

Answer: No, you must call 999 immediately even if the symptoms improve



Q3: What do we do with chest pain and shortness of breath?

Answer: Call 999 immediately, sit the person in a comfortable position, and monitor their condition



Q4: Why do we determine the symptom start time?

Answer: Because the information helps specialists assess the case and make the right decision





Thank You

Quick recognition of the signs and not waiting can save a life

FAST to recognize it • Don't wait • Call 999 • No food or medication • State the symptom time

"In heart attacks and strokes, every minute matters"