



Managing Low Blood Sugar

Calm, Observing the Signs, and Safe Help

"A condition that can occur suddenly and requires early recognition, safe action, and calling emergency services in serious cases"



Why Do We Care About Low Blood Sugar?

Awareness protects the person before the condition deteriorates



**Early
Recognition**
of the Signs



**Helping the
Person**
Before Deterioration




**Preventing
Mistakes**
Such as Feeding the
Unconscious



**Supporting
Those Affected**
Everywhere



Knowing When
to Call 999



A Culture of
Safe First Aid

Remember: Low blood sugar may seem minor, but it can become an emergency if accompanied by loss of consciousness or severe weakness



What Is Meant by Low Blood Sugar?

A drop in blood sugar below the level appropriate for the body



People with diabetes



Delaying meals



Physical exertion without preparation



Blood sugar regulation medications



Illness or exhaustion



Cases requiring specialists

Note: This presentation is for awareness and does not replace the doctor's guidance or the person's individual treatment plan



Who Is Most at Risk of Low Blood Sugar?

Groups that need follow-up and awareness



People with diabetes



Users of diabetes medications



Children with diabetes



The elderly



Those who delay meals



Exerting effort without food

Awareness Message: Knowing the person's health condition — if they disclose it — helps with quick action during an emergency

Early Signs of Low Blood Sugar

Detecting the condition before it deteriorates



Severe hunger



Trembling or shaking



Sweating



Dizziness or weakness



Headache



Sudden anxiety or tension



Difficulty concentrating



Paleness or unusual fatigue

Correct Action: If the person is conscious and able to swallow, help them calmly and follow their health plan if known



Signs of Severe Low Blood Sugar

They call for requesting help immediately



Confusion or unusual behavior

1

Difficulty speaking and understanding

2

Severe weakness preventing them from standing

3

Severe drowsiness or loss of consciousness

4

Seizures

5

Inability to swallow

6

Rapid deterioration of the condition

7

Urgent Action: Call 999 in case of loss of consciousness, seizures, inability to swallow, or clear deterioration



Securing the Area Before Helping

Protecting the person and those around them



**No vehicle
traffic or fall
hazards**



**No sharp
objects nearby**



**A well-
ventilated
place**



**Enough space
around them**



**Disperse the
onlookers**



**Someone to
call emergency
services**

Important Message: Securing the area reduces the chance of an additional injury, especially with dizziness or weakness



Handling a Conscious Person Able to Swallow

Safe help according to their health plan

Monitor the condition's improvement

5

Request help if there's no improvement

6

Call 999 when in doubt

7

Have them sit in a safe place

1

Ask them about diabetes or their plan

2

Help them with a fast sugar source

3

Don't leave them alone

4

Note: Don't force the person to consume anything if they can't swallow normally




General Examples of a Fast Sugar Source

For a conscious person able to swallow



Sweetened juice

If suitable for their plan



Glucose tablets

If available to them



A sugary drink

Depending on the case



Suitable candy

Easy to swallow

Awareness Message: Don't give large amounts randomly, and don't give anything to an unconscious person



What Do You Do After the Condition Improves?

Don't ignore the condition after improvement



Keep them seated until they stabilize



No driving immediately



Informing the family or the person in charge



Following their health plan



Watch for the symptoms returning



Request help if it recurs

Note: Improvement of symptoms doesn't mean ignoring the cause, especially if the condition is recurrent or severe



Handling an Unconscious Person

An emergency — call 999 immediately



Make sure the area is safe

1

Call 999 immediately

2

Check for normal breathing

3

No food or drink

4

Don't put anything in the mouth

5

Don't leave them alone

6

Follow the emergency operator's instructions

7

Important Note: Giving food or drink to an unconscious person can cause choking



If Seizures or Severe Weakness Appear

A quick, safe response



**Move away
dangerous
objects**



**Don't restrain
their
movement
forcefully**



**Don't put
anything in the
mouth**



**Call 999
immediately**



**Monitor
breathing and
responsiveness**



**Follow the
operator's
instructions**

Important Message: Seizures or loss of consciousness are signs that call for requesting help immediately



If the Person Is Known to Have Diabetes

They may carry a medical card or a treatment plan



**Ask them if
they're
conscious**



**Look for a
medical
bracelet**



**Don't use
medications
you don't know**



**Help them
reach their
medications**



**Call 999 in case
of loss of
consciousness**



**Inform the
specialists of
the
information**

Note: Don't give the person medication or an injection unless you're authorized, trained, and within clear instructions



Low Blood Sugar in Children

Watch for sudden changes in behavior



Unusual
fatigue or
crying



Sweating or
trembling



Severe hunger



Difficulty
concentrating



Drowsiness or
weakness



Unusual
behavior

Correct Action: Inform the guardian, help the child per their plan, and call 999 at signs of severity



Low Blood Sugar at School or Work

Organized procedures for health conditions



**Knowing the
disclosed
conditions**



**Contact
numbers kept
safely**



**Training the
supervisors**



**Don't leave
them alone**



**No crowding or
filming**



**Calling 999 in
case of danger**



**Documenting
the time and
observations**

Important Message: Advance preparation protects the person and reduces confusion during emergencies

What Not to Do with Low Blood Sugar

Actions that may increase the danger



**No food for
someone
unconscious**



**Don't force
them to
swallow**



**Don't leave
them alone
while confused**



**No driving
before
stabilizing**



**No injections
without
training**



**Don't ignore
recurrence**



**Don't delay
calling 999**

Awareness Message: Correct help begins with knowing when to help and when to call the specialists



When Do We Call the Emergency Number 999?

Don't wait at signs of severity

No improvement after help

5

Loss of consciousness

1

An injury or fall due to dizziness

6

Seizures

2

Not knowing the cause of the condition

7

Inability to swallow

3

A case in a child or an elderly person

8

Confusion or rapid deterioration

4

Important Message: When in doubt, request help immediately and don't wait for the condition to deteriorate



Preventing Low Blood Sugar

Following the health plan and awareness of the factors



**Adhering to
the doctor's
guidance**



**Not delaying
meals**



**Carrying a
suitable sugar
source**



**Attention
before
prolonged
activity**



**Informing the
family and
workplace**



**Monitoring
recurrent
symptoms**

Note: These are general guidelines and don't replace each person's individual medical plan



The Role of Those Present at the Scene

Organizing the response and reducing confusion



Someone stays with the person



Someone calls 999



Someone disperses the onlookers



Someone looks for the person in charge



Someone clears the way for the ambulance



Someone prepares the symptom timing

Important Message: Organizing roles helps the person and prevents chaos and delay



Do's and Don'ts with Low Blood Sugar

Correct Practices and Others to Avoid

Don't ×

No food for someone unconscious

1

Don't force them to swallow

2

No medications or injections without guidance

3

No driving before stabilizing

4

Don't crowd around them

5

Don't film the case

6

Don't delay calling emergency services

7

Do ✓

Sit them in a safe place

1

Help a conscious person per their plan

2

Monitor the condition's improvement

3

Call 999 in case of danger

4

Don't leave them alone while confused

5

Inform the specialists of the information

6

Respect the person's privacy

7



Interactive Scenarios

Practical Application of What You've Learned

Scenario 2

Signs of low blood sugar appeared, then they lost consciousness

Correct Action: Call 999 immediately, don't give them food or drink, and monitor breathing until the specialists arrive

Scenario 1

A diabetic person felt trembling, sweating, and dizziness while conscious

Correct Action: Sit them in a safe place, help them per their plan with a fast sugar source, and monitor for improvement

Scenario 4

After improving, they wanted to drive the vehicle right away

Correct Action: Don't allow driving until their condition stabilizes, and make sure someone is there to help them

Scenario 3

A student at school became confused and is sweating

Correct Action: Inform the supervisor, don't leave them alone, help them per their plan, and call 999 in case of danger



Test Your Knowledge

Quick Questions to Assess Your Understanding

Q1: What are the early signs of low blood sugar?

Answer: They may include trembling, sweating, severe hunger, dizziness, headache, and difficulty concentrating



Q2: Is it acceptable to give food to an unconscious person?

Answer: No, an unconscious person is not given any food or drink — call 999 immediately



Q3: What do we do if they're conscious and able to swallow?

Answer: We sit them in a safe place and help them per their plan or with a suitable sugar source while monitoring



Q4: When should you call 999?

Answer: In case of loss of consciousness, seizures, inability to swallow, deterioration, or no improvement





Thank You

Early recognition of the signs and providing safe help protect the person

Observe the signs • Help the conscious per their plan • No food for the unconscious • Call 999 in case of danger

"Quick awareness of low blood sugar and a safe response protect the person at the right moment"