



Managing Fainting and the Recovery Position

Calm, Securing the Area, and Monitoring Breathing

"Fainting is a condition requiring calm, securing the area, monitoring breathing and responsiveness, and requesting help when needed"



Why Do We Care About Fainting?

It can happen at home, school, work, or public places



Protecting the Injured Person
from Falling and Injury



Breathing Safety
and Monitoring
Responsiveness




Preventing Crowding
and Confusion



Requesting Help
at Signs of Danger



Reassuring the Injured Person
After Coming To



A Culture of
Safe First Aid

Remember: Recurrent fainting or that accompanied by severe pain or difficulty breathing needs an immediate call to 999



What Is Meant by Fainting?

A temporary loss of consciousness or a sudden weakening of responsiveness



**Temporary loss
of
consciousness**



**Severe
dizziness**



**Sudden
weakness**



**Sweating or
paleness**



**Inability to
stand**



**Confusion after
coming to**

Note: Not every fainting is minor — you must watch for the signs accompanying the condition

Common Causes of Fainting

Some of them need a medical assessment



Standing for a long time



Exhaustion or lack of sleep



Dehydration or low fluid intake



Severe heat or crowding



Strong emotion or fear



Low blood sugar



Health conditions or medications



An injury or severe pain

Awareness Message: Knowing the cause helps with the initial handling, but it doesn't replace requesting help in serious cases



Signs Before Fainting

Early intervention prevents a fall



Dizziness and loss of balance

1

Paleness in the face

2

Sudden sweating

3

Nausea and general weakness

4

Blurred vision

5

Ringing in the ears

6

Difficulty standing

7

Correct Action: Help the person sit or lie down in a safe place before they fall



Securing the Area Before Helping

Check for nearby hazards




**Vehicle or
bicycle traffic**




**Stairs or raised
edges**



**A slippery or
wet floor**



**Crowding or
pushing**



Heat or smoke



**Sharp objects
or obstacles**

Correct Action: Remove the hazards, provide space around the injured person, and don't put yourself in danger



What to Do When Fainting Feels Imminent

Act quickly and calmly



**Help them sit
or lie down**



**Move them
away from heat
or crowds**



**Adequate
ventilation in
the area**



**Encourage
them to
breathe calmly**



**Don't let them
stand quickly**



**Monitor their
condition and
request help**

Important Message: Intervening before a fall can prevent an additional injury



What to Do When Consciousness Is Lost

A quick assessment without confusion

No food or drink for someone unconscious

5

Don't move them if an injury is suspected

6

Follow the emergency operator's instructions

7

Make sure the area is safe

1

Check the injured person's responsiveness gently

2

Ask for 999 to be called

3

Check for normal breathing

4

Note: If they're unresponsive and not breathing normally, treat it as an emergency and call 999



When Do We Use the Recovery Position?

Unresponsive but breathing normally



**Loss of
consciousness
with normal
breathing**



**Protecting the
airway**



**Waiting for
help to arrive**



**No suspicion of
a serious injury**



**Following the
999 operator's
instructions**

Important Note: If a head, neck, or back injury is suspected, don't move the injured person



The Recovery Position in General

It helps keep the injured person in a relatively safe position



Make sure they're breathing normally

1

Don't use it with a back injury

2

Ask another person for help

3

Follow the 999 operator's instructions

4

Monitor breathing continuously

5

Don't leave the injured person alone

6

Awareness Message: The recovery position is useful in specific cases, but it doesn't mean the condition doesn't need monitoring



Monitoring Breathing and Responsiveness

Continue monitoring after fainting



Are they breathing normally?

1

Are they regaining consciousness?

2

Is there dizziness after coming to?

3

Is there pain in the chest or head?

4

Is there an injury from the fall?

5

Is the condition recurring or worsening?

6

Correct Action: If breathing changes, responsiveness weakens, or signs of severity appear, call 999 immediately



After Regaining Consciousness

Don't rush to get them up or leave them



**Speak calmly
and reassure
them**



**Don't let them
stand right
away**



**Ask them
about pain and
dizziness**



**Keep them
seated until
they stabilize**



**No food for
someone who
is confused**



**Request help if
it recurs**

Important Message: Quick improvement doesn't mean ignoring the cause of fainting, especially if it's unusual



When Is Fainting an Emergency?

Call 999 when these signs appear

A head or neck injury

5

Seizures or severe confusion

6

Fainting after a strong accident

7

Repeated fainting

8

Not regaining consciousness quickly

1

Difficulty breathing

2

Chest pain

3

A severe headache after a fall

4

Note: When in doubt about the severity of the case, call emergency services and don't wait



Fainting Due to Heat or Crowding

Exhaustion, dehydration, or poor ventilation



Move them to a ventilated place



Move them away from the sun and heat



Help them sit



Monitor their consciousness and breathing



No drink for someone who is unconscious



Call 999 if it persists

Awareness Message: Ventilation, rest, and drinking water regularly reduce dizziness and fainting in hot weather



Fainting After a Fall or Injury

Handle with greater caution



**Don't move
them with
head pain**



**Don't rush to
get them up**



**Check
breathing**



**Call 999 in case
of injury**



**Disperse the
onlookers**



**Wait for the
specialists**

Note: Fainting after an injury may indicate a condition needing urgent assessment

What Not to Do with Fainting

Actions that may increase the danger



**Don't splash
water
randomly**



**No food or
drink for
someone
unconscious**



**Don't force
them to stand
immediately**



**Don't move
them in case of
injury**



**Don't crowd
around them**



**Don't delay
calling 999**



**Don't film or
post the
injured person**

Awareness Message: Calmness and requesting help are more important than random actions



Preventing Recurrent Fainting

Attention to the contributing factors



**Drinking water
regularly**



**No prolonged
standing
without
movement**



**Sit down as
soon as you
feel dizzy**



**Avoid heat and
crowds**



Eat regularly



**Consult
specialists if it
recurs**



**Inform the
family or
workplace**

Note: Recurrent fainting needs assessment from healthcare specialists



The Role of Those Present at the Scene

Organizing roles reduces chaos



Someone calls
999



Someone
disperses the
onlookers



Someone
clears the way
for the
ambulance



Someone
monitors
breathing



Someone
gathers the
information



Someone
reassures the
injured person

Important Message: Cooperation and organization are better than crowding or filming

Do's and Don'ts with Fainting

Correct Practices and Others to Avoid

Don't ✗

No food or drink for someone unconscious

1

Don't force them to stand quickly

2

Don't move them if an injury is suspected

3

No recovery position with a back injury

4

Don't splash water randomly

5

Don't delay calling emergency services

6

Don't film or post the injured person

7

Do ✓

Make sure the area is safe

1

Help them sit when dizzy

2

Monitor breathing and responsiveness

3

Call 999 in case of danger

4

Don't leave them if they're unstable

5

Disperse the onlookers and provide ventilation

6

Follow the emergency operator's instructions

7



Interactive Scenarios

Practical Application of What You've Learned

Scenario 2

A person lost consciousness but is breathing normally with no signs of injury

Correct Action: Monitor their breathing, use the recovery position if you know it, and call 999 when needed

Scenario 1

A person felt dizzy in a crowded place and said they might pass out

Correct Action: Help them sit in a safe, ventilated place, monitor them, and request help when needed

Scenario 4

After coming to, the injured person tried to stand quickly

Correct Action: Stop them from standing, and help them stay seated until their condition stabilizes

Scenario 3

A person fainted after a fall and complains of neck pain after coming to

Correct Action: Don't move them, reassure them, keep the area safe, and call 999 immediately



Test Your Knowledge

Quick Questions to Assess Your Understanding

Q1: What is the first action when you see a person feeling severely dizzy?

Answer: Helping them sit or lie down in a safe place to reduce the risk of falling



Q2: Is it acceptable to give food or drink to an unconscious person?

Answer: No, food or drink is not given to someone unconscious or unable to swallow



Q3: When do you call 999 in fainting cases?

Answer: When consciousness isn't regained, or with difficulty breathing, chest pain, a head injury, or repeated fainting



Q4: When do we not use the recovery position?

Answer: When a head, neck, or back injury is suspected, except in case of an immediate danger and per the specialists





Thank You

Fainting requires calm, securing the area, monitoring breathing, and requesting help when needed

Help them sit • No food for the unconscious • Don't move them in case of injury • Call 999

"A mindful response to fainting begins with calm and ends with requesting help at the right time"