



Managing Fractures and Falls

Calm, Assessment, and Measured Movement

"Falls and fractures can be minor or serious, so they must be handled carefully, with help requested when there are signs of severity"



Why Do We Care About Fractures and Falls?

Some injuries don't show their severity right away



**Reducing
Worsening**
of the Injury



**Protecting the
Injured Person**
from Wrong Movement



**Preventing
Crowding**
and Confusion



Requesting Help
at the Right Time



**Emotional
Support**
Until the Specialists
Arrive



Fall Prevention
at Home and in
Facilities

Remember: Not every fall is minor — any severe pain or difficulty moving requires careful handling



Common Causes of Falls

Factors that increase the likelihood of accidents



Wet or slippery floors



Obstacles in walkways



Poor lighting



Stairs used unsafely



Running in enclosed spaces



Wires and unsecured rugs



Fatigue or dizziness



Unsuitable footwear

Awareness Message: Prevention starts with removing obstacles and being attentive to the area before moving

Who Is Most at Risk?

Groups that need follow-up and extra precautions



Children



The elderly



People of
Determination



Poor balance



Users of
mobility aids



Fatigue or
dizziness



Workers
constantly on
the move

Note: Protecting the most vulnerable groups requires a safe environment and appropriate supervision

Signs of a Suspected Fracture

Act carefully when there are strong signs



Severe pain at the injury site

1

Obvious swelling

2

Difficulty moving the limb

3

Inability to stand

4

A change in the shape or position of the limb

5

Pain that increases with movement

6

A sound or strong impact at the time of the fall

7

Correct Action: When a fracture is suspected, don't move the injured part, and request help when needed

Signs of a Serious Injury After a Fall

Especially in the head, neck, or back

Obvious bleeding or a major injury

5

Weakness or numbness in the limbs

6

Inability to move

7

Vomiting or a change in behavior

8

Loss of consciousness or weak responsiveness

1

Severe dizziness or confusion

2

Pain in the head or neck

3

Difficulty breathing

4

Urgent Action: Call 999 immediately and don't move the injured person except in case of an immediate danger



Securing the Area Before Helping

Check for hazards before approaching



A slippery or wet floor



Glass or sharp objects



Nearby vehicle traffic



A risk of an additional fall



Fire, smoke, or electricity



Crowding or pushing

Correct Action: Secure the area as much as possible, ask those present to move away, and don't put yourself in danger

Initial Response After a Fall

Act calmly and don't rush to move them

Call 999 when in doubt

5

Reassure the injured person and stay with them

6

Wait for the specialists

7

Make sure the area is safe

1

Ask the injured person if they can respond

2

Ask them not to move

3

Check for signs of severity

4

Important Message: Staying calm and not moving the injured person unnecessarily are among the most important safety steps



What Do You Do When a Fracture Is Suspected?

Minimize movement and wait for help



Don't move the injured part



Help them stay still



Don't reposition the limb



Simple support within your training



No strong pressure on the site



Call 999 in severe cases

Note: Attempting to reposition the injured limb or move it incorrectly can worsen the injury

Head, Neck, and Back Injuries

Moving the injured person can increase the danger



A fall from a height

1

Sustaining a strong blow

2

Pain in the neck or back

3

Dizziness or confusion

4

Loss of consciousness, even briefly

5

Weakness or numbness in the limbs

6

Inability to move

7

Correct Action: Don't move the injured person, and request help immediately via 999



Handling the Elderly After a Fall

Act carefully even with a minor fall



Don't rush to get them up



Ask them where the pain is



Watch for dizziness and confusion



Don't move them in case of severe pain



Call 999 in severe cases



Inform the specialists about medications

Awareness Message: A fall in the elderly requires careful assessment, even if it appears minor



Handling Children After a Fall

Children may not be able to describe the pain accurately



**Watch the
crying and
behavior**



**Difficulty
moving**



**Unusual
drowsiness**



**Don't force
them to stand**



**Call 999 in
severe cases**



**No
medications
without
medical
guidance**

Note: Any abnormal change in a child's consciousness or behavior after a fall needs immediate attention



Handling a Fall in Public Places

Organizing the area and preventing crowding



**Move away
those who
aren't needed**



**Provide
enough space**



**No filming or
posting**



**Someone calls
999**



**Someone
guides the
ambulance
teams**



**Don't move the
injured person
to clear the
area**

Important Message: Organizing the site helps the injured person and facilitates the arrival of the competent authorities



Using Ice or Cold Compresses

They may help with minor bruises — use with caution



**A compress
wrapped in
cloth**



**Don't apply ice
directly**



**Not for large
wounds**



**Don't delay
requesting
help**



**Monitor the
pain and
swelling**

Note: Cold compresses are not a substitute for assessing the injury when a fracture is suspected

Preventing Falls at Home

Simple modifications reduce the risk of falls



Drying floors immediately



Removing obstacles from walkways



Securing rugs



Good lighting for walkways



Support handles for the elderly



Storing children's toys



Wires away from walkways

Awareness Message: A safe home reduces the chance of falls, especially for children and the elderly




Preventing Falls at Schools and Work

Organizing educational and work environments



Report slippery floors



Warning signs



Walkways free of obstacles



No running in corridors



Using stairs properly



Storing tools in their places



Adequate lighting

Important Message: Reporting a hazard before an injury occurs is part of the safety culture



What Not to Do with Fractures and Falls

Actions that may increase pain or worsen the injury




Don't move the injured person unnecessarily



Don't reposition the injured limb



Don't force them to stand



No strong pressure on the injury



No food for an unstable injured person



Don't crowd around the injured person



Don't delay calling 999

Awareness Message: Correct help is sometimes preventing movement and requesting help



The Role of the First Aid Kit

Supplies for providing simple initial support



Medical gloves



Clean
dressings



Sterile gauze



An elastic
bandage



An instant cold
pack



Medical tape



A simplified
first aid guide

Note: Don't use the kit's supplies to perform a procedure you don't know or haven't been trained in

Do's and Don'ts with Fractures and Falls

Correct Practices and Others to Avoid

Don't ✗

Don't move the injured person unnecessarily

1

Don't force them to stand

2

Don't reposition the limb

3

Don't apply ice directly

4

Don't ignore head or neck pain

5

Don't delay requesting help

6

Don't film or post the injured person

7

Do ✓

Make sure the area is safe

1

Reassure them and ask them not to move

2

Call 999 in case of danger

3

Disperse the onlookers

4

Use the supplies within your knowledge

5

Wait for the specialists when in doubt

6

Report fall hazards

7

Interactive Scenarios

Practical Application of What You've Learned

Scenario 2 A child fell and can't move their arm

Correct Action: Don't force them to move, calm them, inform the guardian, and request help

Scenario 1 A person fell on the stairs and is complaining of back pain

Correct Action: Don't move them, secure the area, reassure them, and call 999 immediately

Scenario 4 A wet floor in a school corridor caused a student to slip

Correct Action: Help the student safely, report the hazard, and place a warning to prevent others from falling

Scenario 3 An elderly person fell and feels dizzy with difficulty standing

Correct Action: Don't rush to get them up, keep them safe, monitor them, and call 999 in severe cases



Test Your Knowledge

Quick Questions to Assess Your Understanding

Q1: What is the first step when a person falls in front of you?

Answer: Securing the area and calmly assessing the injured person's condition without moving them when a serious injury is suspected



Q2: When should you call 999 after a fall?

Answer: In case of loss of consciousness, head or neck pain, difficulty moving, or a suspected fracture



Q3: Is it acceptable to force the injured person to stand after a fall?

Answer: No, if they feel severe pain or have signs of a serious injury



Q4: Why don't we apply ice directly to the skin?

Answer: Because it may harm the skin — use a cold compress wrapped in a piece of cloth





Thank You

Calm, quick assessment, avoiding unnecessary movement, and requesting help in case of danger

Don't rush to move the injured person • Watch the head and neck • Prevention through a safe environment • Call 999

"Your calm action after a fall can prevent the injury from worsening and protect a person's life"