

Home Work #1

Ali BaniAsad 96108378

October 8, 2021

1 Question 1


2 Question 2

$$G(s) = \frac{K}{s(s^2 + s + a)}, \quad a > 0, \quad K > 0 \quad (1)$$

The transfer function in which s is replaced by $j\omega$ where ω is frequency.

Cause $a > 0$ we have just on pole on imaginary axis.

Figure 1: Contour near the origin of the s plane and closed contour in the s plane




On the semicircular path with radius ϵ (where $\epsilon \ll 1$), the complex variable s can be written

$$s = \epsilon \exp(j\theta)$$

where θ varies from $-\frac{\pi}{2}$ to $\frac{\pi}{2}$.

Figure 2: s-Plane



Now we calculate $j\omega$ in A, C, D and F.

A:

$$s = j\omega, \quad \omega = -\epsilon$$

$$\lim_{\omega \rightarrow -\epsilon} G(j\omega) = \frac{K}{-j\epsilon((j\epsilon)^2 - j\epsilon + a)} = \frac{K}{-j\epsilon(-\epsilon^2 - j\epsilon + a)} = \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{K}{-aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{Kj}{a\epsilon} = \infty \angle 90^\circ$$

C:

$$s = j\omega, \quad \omega = \epsilon$$

$$\lim_{\omega \rightarrow \epsilon} G(j\omega) = \frac{K}{j\epsilon((j\epsilon)^2 + j\epsilon + a)} = \frac{K}{j\epsilon(-\epsilon^2 + j\epsilon + a)} = \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{j\epsilon^3 - \epsilon^2 + aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{K}{-aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{-Kj}{a\epsilon} = \infty \angle -90^\circ$$

A to C:

$$s = j\omega, \quad \omega = \epsilon \exp(j\theta), \quad \theta : -\frac{\pi}{2} \rightarrow 0 \rightarrow \frac{\pi}{2} (CCW)$$

$$\lim_{\epsilon \rightarrow 0} G(\epsilon \exp(j\theta)) = \frac{K}{\epsilon \exp(j\theta)((\epsilon \exp(j\theta))^2 + \epsilon \exp(j\theta) + a)} = \frac{K}{\epsilon^3 \exp(3j\theta) + \epsilon^2 \exp(2j\theta) + a\epsilon \exp(j\theta)}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{a\epsilon \exp(j\theta)} = \lim_{\epsilon \rightarrow 0} \frac{K \exp(-j\theta)}{a\epsilon} = \infty \angle -\theta$$

D:

$$s = j\omega, \quad \omega = \infty$$

$$\lim_{\omega \rightarrow \infty} G(j\omega) = \lim_{\omega \rightarrow \infty} \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \lim_{\omega \rightarrow \infty} \frac{K}{j\omega(-\omega^2 + j\omega + a)} = \lim_{\omega \rightarrow \infty} \frac{K}{-j\omega^3 - \omega^2 + aj\omega}$$

$$\lim_{\omega \rightarrow \infty} \frac{K}{-j\omega^3} = \lim_{\omega \rightarrow \infty} \frac{jK}{\omega^3} = 0 \angle 90^\circ$$

F:

$$s = j\omega, \quad \omega = -\infty$$

$$\lim_{\omega \rightarrow -\infty} G(j\omega) = \lim_{\omega \rightarrow -\infty} \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \lim_{\omega \rightarrow -\infty} \frac{K}{j\omega(-\omega^2 + j\omega + a)} = \lim_{\omega \rightarrow -\infty} \frac{K}{-j\omega^3 - \omega^2 + aj\omega}$$

$$\lim_{\omega \rightarrow -\infty} \frac{K}{-j\omega^3} = \lim_{\omega \rightarrow -\infty} \frac{jK}{\omega^3} = 0 \angle -90^\circ$$

Figure 3: Nyquist plot using wolfram ($a = 1, K = 1$)





Figure 4: Nyquist plot using MATLAB ($a = 1, K = 1$)



Now we know shape of nyquist plot. we find out where it's equal to -1 .

$$G(j\omega) = -1 + 0j = \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \frac{K}{-j\omega^3 - \omega^2 + aj\omega} = -1$$

$$\rightarrow j\omega^3 + \omega^2 - aj\omega = K \rightarrow j\omega^3 + \omega^2 - aj\omega - K = 0$$

Two equation and two unknowns.

$$j\omega^3 = aj\omega \rightarrow \omega^2 = a$$


$$\omega^2 = K \rightarrow \omega^2 = K$$

When $a = K$ the nyquist plot cross from -1 point. When $a > K$ the nyquist plot is before -1 and when $a < K$ the nyquist plot cross -1 and system is unstable.

Now we check above using MATLAB.


- $a > K$
 $a = 10, K = 1$

Figure 5: Nyquist plot using MATLAB ($a = 10, K = 1$)




- $a = K$
 $a = 1, K = 1$

Figure 6: Nyquist plot using MATLAB ($a = 1, K = 1$)



- $a < K$
 $a = 1, K = 10$

Figure 7: Nyquist plot using MATLAB ($a = 1, K = 10$)



Contents

1	Question 1	2
2	Question 2	3

List of Figures

1	Contour near the origin of the s plane and closed contour in the s plane	3
2	s -Plane	4
3	Nyquist plot using wolfram ($a = 1, K = 1$)	6
4	Nyquist plot using MATLAB ($a = 1, K = 1$)	6
5	Nyquist plot using MATLAB ($a = 10, K = 1$)	7
6	Nyquist plot using MATLAB ($a = 1, K = 1$)	8
7	Nyquist plot using MATLAB ($a = 1, K = 10$)	8

List of Tables