

# Home Work #1

Ali BaniAsad 96108378

October 8, 2021


## 1 Question 1

$$G(s) = \exp(-\tau s), \quad \tau > 0$$

$$\angle G(j\omega) = -\tau\omega$$


$$|G(j\omega)| = 1 \rightarrow 20 \log(|G(j\omega)|) = 0$$

Figure 1: Bode digram using MATLAB ( $\tau = 1$ )




There is no pole in imaginary axis and nyquist path is from  $0 \rightarrow \infty \rightarrow -\infty \rightarrow 0$ .

Figure 2: Closed contour in the s plane



So the Nyquist plot is circle in orogin with radius 1 and clock wise.

Figure 3: Nyquist plot using MATLAB ( $\tau = 1$ )




## 2 Question 2

$$G(s) = \frac{K}{s(s^2 + s + a)}, \quad a > 0, \quad K > 0 \quad (1)$$

The transfer function in which  $s$  is replaced by  $j\omega$  where  $\omega$  is frequency.

Cause  $a > 0$  we have just on pole on imaginary axis.

Figure 4: Contour near the origin of the  $s$  plane and closed contour in the  $s$  plane




On the semicircular path with radius  $\epsilon$  (where  $\epsilon \ll 1$ ), the complex variable  $s$  can be written

$$s = \epsilon \exp(j\theta)$$

where  $\theta$  varies from  $-\frac{\pi}{2}$  to  $\frac{\pi}{2}$ .

Figure 5: s-Plane



Now we calculate  $j\omega$  in A, C, D and F.

A:

$$s = j\omega, \quad \omega = -\epsilon$$

$$\lim_{\omega \rightarrow -\epsilon} G(j\omega) = \frac{K}{-j\epsilon((j\epsilon)^2 - j\epsilon + a)} = \frac{K}{-j\epsilon(-\epsilon^2 - j\epsilon + a)} = \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{K}{-aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{Kj}{a\epsilon} = \infty \angle 90^\circ$$

C:

$$s = j\omega, \quad \omega = \epsilon$$

$$\lim_{\omega \rightarrow \epsilon} G(j\omega) = \frac{K}{j\epsilon((j\epsilon)^2 + j\epsilon + a)} = \frac{K}{j\epsilon(-\epsilon^2 + j\epsilon + a)} = \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{j\epsilon^3 - \epsilon^2 + aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{K}{-aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{-Kj}{a\epsilon} = \infty \angle -90^\circ$$

A to C:

$$s = j\omega, \quad \omega = \epsilon \exp(j\theta), \quad \theta : -\frac{\pi}{2} \rightarrow 0 \rightarrow \frac{\pi}{2} (CCW)$$

$$\lim_{\epsilon \rightarrow 0} G(\epsilon \exp(j\theta)) = \frac{K}{\epsilon \exp(j\theta)((\epsilon \exp(j\theta))^2 + \epsilon \exp(j\theta) + a)} = \frac{K}{\epsilon^3 \exp(3j\theta) + \epsilon^2 \exp(2j\theta) + a\epsilon \exp(j\theta)}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{a\epsilon \exp(j\theta)} = \lim_{\epsilon \rightarrow 0} \frac{K \exp(-j\theta)}{a\epsilon} = \infty \angle -\theta$$

D:

$$s = j\omega, \quad \omega = \infty$$

$$\lim_{\omega \rightarrow \infty} G(j\omega) = \lim_{\omega \rightarrow \infty} \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \lim_{\omega \rightarrow \infty} \frac{K}{j\omega(-\omega^2 + j\omega + a)} = \lim_{\omega \rightarrow \infty} \frac{K}{-j\omega^3 - \omega^2 + aj\omega}$$

$$\lim_{\omega \rightarrow \infty} \frac{K}{-j\omega^3} = \lim_{\omega \rightarrow \infty} \frac{jK}{\omega^3} = 0 \angle 90^\circ$$

F:

$$s = j\omega, \quad \omega = -\infty$$

$$\lim_{\omega \rightarrow -\infty} G(j\omega) = \lim_{\omega \rightarrow -\infty} \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \lim_{\omega \rightarrow -\infty} \frac{K}{j\omega(-\omega^2 + j\omega + a)} = \lim_{\omega \rightarrow -\infty} \frac{K}{-j\omega^3 - \omega^2 + aj\omega}$$

$$\lim_{\omega \rightarrow -\infty} \frac{K}{-j\omega^3} = \lim_{\omega \rightarrow -\infty} \frac{jK}{\omega^3} = 0 \angle -90^\circ$$

Figure 6: Nyquist plot using wolfram ( $a = 1, K = 1$ )





Figure 7: Nyquist plot using MATLAB ( $a = 1, K = 1$ )





Now we know shape of nyquist plot. we find out where it's equal to  $-1$ .

$$G(j\omega) = -1 + 0j = \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \frac{K}{-j\omega^3 - \omega^2 + aj\omega} = -1$$
$$\rightarrow j\omega^3 + \omega^2 - aj\omega = K \rightarrow j\omega^3 + \omega^2 - aj\omega - K = 0$$

Two equation and two unknowns.


$$j\omega^3 = aj\omega \rightarrow \omega^2 = a$$
$$\omega^2 = K \rightarrow \omega^2 = K$$

When  $a = K$  the nyquist plot cross from  $-1$  point. When  $a > K$  the nyquist plot is before  $-1$  and when  $a < K$  the nyquist plot cross  $-1$  and system is unstable.

Now we check above using MATLAB.


- $a > K$   
 $a = 10, K = 1$

Figure 8: Nyquist plot using MATLAB ( $a = 10, K = 1$ )




- $a = K$   
 $a = 1, K = 1$

Figure 9: Nyquist plot using MATLAB ( $a = 1, K = 1$ )



- $a < K$   
 $a = 1, K = 10$

Figure 10: Nyquist plot using MATLAB ( $a = 1, K = 10$ )




### 3 Question 3

$$G(s) = \frac{K}{s(s^2 + s + a)}, \quad a > 0, \quad K > 0 \quad (2)$$

The transfer function in which  $s$  is replaced by  $j\omega$  where  $\omega$  is frequency.

Cause  $a > 0$  we have just on pole on imaginary axis.

Figure 11: Contour near the origin of the  $s$  plane and closed contour in the  $s$  plane




On the semicircular path with radius  $\epsilon$  (where  $\epsilon \ll 1$ ), the complex variable  $s$  can be written

$$s = \epsilon \exp(j\theta)$$

where  $\theta$  varies from  $-\frac{\pi}{2}$  to  $\frac{\pi}{2}$ .

Figure 12: s-Plane



Now we calculate  $j\omega$  in A, C, D and F.

A:

$$s = j\omega, \quad \omega = -\epsilon$$

$$\lim_{\omega \rightarrow -\epsilon} G(j\omega) = \frac{K}{-j\epsilon((j\epsilon)^2 - j\epsilon + a)} = \frac{K}{-j\epsilon(-\epsilon^2 - j\epsilon + a)} = \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{K}{-aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{Kj}{a\epsilon} = \infty \angle 90^\circ$$

C:

$$s = j\omega, \quad \omega = \epsilon$$

$$\lim_{\omega \rightarrow \epsilon} G(j\omega) = \frac{K}{j\epsilon((j\epsilon)^2 + j\epsilon + a)} = \frac{K}{j\epsilon(-\epsilon^2 + j\epsilon + a)} = \frac{K}{j\epsilon^3 - \epsilon^2 - aj\epsilon}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{j\epsilon^3 - \epsilon^2 + aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{K}{-aj\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{-Kj}{a\epsilon} = \infty \angle -90^\circ$$

A to C:

$$s = j\omega, \quad \omega = \epsilon \exp(j\theta), \quad \theta : -\frac{\pi}{2} \rightarrow 0 \rightarrow \frac{\pi}{2} (CCW)$$

$$\lim_{\epsilon \rightarrow 0} G(\epsilon \exp(j\theta)) = \frac{K}{\epsilon \exp(j\theta)((\epsilon \exp(j\theta))^2 + \epsilon \exp(j\theta) + a)} = \frac{K}{\epsilon^3 \exp(3j\theta) + \epsilon^2 \exp(2j\theta) + a\epsilon \exp(j\theta)}$$

$$\lim_{\epsilon \rightarrow 0} \frac{K}{a\epsilon \exp(j\theta)} = \lim_{\epsilon \rightarrow 0} \frac{K \exp(-j\theta)}{a\epsilon} = \infty \angle -\theta$$

D:

$$s = j\omega, \quad \omega = \infty$$

$$\lim_{\omega \rightarrow \infty} G(j\omega) = \lim_{\omega \rightarrow \infty} \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \lim_{\omega \rightarrow \infty} \frac{K}{j\omega(-\omega^2 + j\omega + a)} = \lim_{\omega \rightarrow \infty} \frac{K}{-j\omega^3 - \omega^2 + aj\omega}$$

$$\lim_{\omega \rightarrow \infty} \frac{K}{-j\omega^3} = \lim_{\omega \rightarrow \infty} \frac{jK}{\omega^3} = 0 \angle 90^\circ$$

F:

$$s = j\omega, \quad \omega = -\infty$$

$$\lim_{\omega \rightarrow -\infty} G(j\omega) = \lim_{\omega \rightarrow -\infty} \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \lim_{\omega \rightarrow -\infty} \frac{K}{j\omega(-\omega^2 + j\omega + a)} = \lim_{\omega \rightarrow -\infty} \frac{K}{-j\omega^3 - \omega^2 + aj\omega}$$

$$\lim_{\omega \rightarrow -\infty} \frac{K}{-j\omega^3} = \lim_{\omega \rightarrow -\infty} \frac{jK}{\omega^3} = 0 \angle -90^\circ$$

Figure 13: Nyquist plot using wolfram ( $a = 1, K = 1$ )





Figure 14: Nyquist plot using MATLAB ( $a = 1, K = 1$ )



Now we know shape of nyquist plot. we find out where it's equal to  $-1$ .

$$G(j\omega) = -1 + 0j = \frac{K}{j\omega((j\omega)^2 + j\omega + a)} = \frac{K}{-j\omega^3 - \omega^2 + aj\omega} = -1$$

$$\rightarrow j\omega^3 + \omega^2 - aj\omega = K \rightarrow j\omega^3 + \omega^2 - aj\omega - K = 0$$

Two equation and two unknowns.

$$j\omega^3 = aj\omega \rightarrow \omega^2 = a$$


$$\omega^2 = K \rightarrow \omega^2 = K$$

When  $a = K$  the nyquist plot cross from  $-1$  point. When  $a > K$  the nyquist plot is before  $-1$  and when  $a < K$  the nyquist plot cross  $-1$  and system is unstable.

Now we check above using MATLAB.


- $a > K$   
 $a = 10, K = 1$

Figure 15: Nyquist plot using MATLAB ( $a = 10, K = 1$ )




- $a = K$   
 $a = 1, K = 1$

Figure 16: Nyquist plot using MATLAB ( $a = 1, K = 1$ )



- $a < K$   
 $a = 1, K = 10$

Figure 17: Nyquist plot using MATLAB ( $a = 1, K = 10$ )





## Contents

<b>1</b>	<b>Question 1</b>	<b>2</b>
<b>2</b>	<b>Question 2</b>	<b>5</b>
<b>3</b>	<b>Question 3</b>	<b>11</b>

## List of Figures

1	Bode digram using MATLAB ( $\tau = 1$ ) . . . . .	2
2	Closed contour in the s plane . . . . .	3
3	Nyquist plot using MATLAB ( $\tau = 1$ ) . . . . .	4
4	Contour near the origin of the s plane and closed contour in the s plane . . . . .	5
5	s-Plane . . . . .	6
6	Nyquist plot using wolfram ( $a = 1, K = 1$ ) . . . . .	8
7	Nyquist plot using MATLAB ( $a = 1, K = 1$ ) . . . . .	8
8	Nyquist plot using MATLAB ( $a = 10, K = 1$ ) . . . . .	9
9	Nyquist plot using MATLAB ( $a = 1, K = 1$ ) . . . . .	10
10	Nyquist plot using MATLAB ( $a = 1, K = 10$ ) . . . . .	10
11	Contour near the origin of the s plane and closed contour in the s plane . . . . .	11
12	s-Plane . . . . .	12
13	Nyquist plot using wolfram ( $a = 1, K = 1$ ) . . . . .	14
14	Nyquist plot using MATLAB ( $a = 1, K = 1$ ) . . . . .	14
15	Nyquist plot using MATLAB ( $a = 10, K = 1$ ) . . . . .	15
16	Nyquist plot using MATLAB ( $a = 1, K = 1$ ) . . . . .	16
17	Nyquist plot using MATLAB ( $a = 1, K = 10$ ) . . . . .	16

## List of Tables