


A


Hypervariability score
1.0
0.5
0.0
−0.5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 X Y
Chromosome


B


Distribution of CpG hypervariability across feature types
% of CpGs in feature
20
15
10
5
0
exon
intergenic
intron
promoter
Hypervariability score (percentile)
Bottom 10%
10−20th
20−30th
30−40th
40−50th
50−60th
60−70th
70−80th
80−90th
Top 10%

C


TE class enrichment of top 1% hvCpGs (0−1kb from centre)
(black box = mean element size)
RC
LINE
DNA
LTR
SINE
Retroposon
***
***
***
0−1kb
distance from TE centre
Mean Kimura divergence (CpG−corrected %) as proxy for age
30
25
20
15
log2 OR
0.2
0.0
−0.2

D


TE family enrichment of top 1% hvCpGs (0−1kb from centre)
(black box = mean element size)
CR1
tRNA−RTE
L2
tRNA
RTE−BovB
MIR
RTE−X
Gypsy
5S−Deu−L2
LTR
hAT−Ac
DNA
Helitron
Penelope
hAT−Tip100
ERVL
hAT−Charlie
TcMar−Tc2
hAT−Blackjack
L1
ERVL−MaLR
hAT
TcMar−Tigger
ERV1
TcMar−Mariner
MULE−MuDR
SVA
PiggyBac
ERVK
Alu
**
***
*
.
***
***
**
***
***
***
***
***
***
**
***
***
0−1kb
distance from TE centre
Mean Kimura divergence (CpG−corrected %) as proxy for age
30
20
10
log2 OR
0.4
0.2
0.0
−0.2