

# **Coupling between tolerance and resistance differs between related *Eimeria* parasite species: implications for coevolution with their mouse hosts**

## **Abstract**

Resistance (host capacity to reduce parasite burden) and tolerance (host capacity to reduce impact on its health for a given parasite burden) manifest two different lines of defence. Tolerance can be independent from resistance, traded-off against it, or the two can be positively correlated because of redundancy in underlying (immune) processes. We here tested whether closely related parasite species could show differences in this coupling between tolerance and resistance. We tested this in experimental infections with two parasite species of genus *Eimeria*. We measured proxies for resistance (the (inverse of) number of parasite transmission stages (oocysts) per gram of feces at the day of maximal shedding) and tolerance (the slope of maximum relative weight loss compared to day of infection on number of oocysts per gram of feces at the day of maximal shedding for each host strain) in four inbred mouse strains belonging to two mouse subspecies, *Mus musculus domesticus* and *M. m. musculus*. We found a negative correlation between resistance and tolerance against *E. falciformis*, while the two are uncoupled against *E. ferrisi*. We conclude that resistance and tolerance against the first parasite species might be traded off, but evolve more independently in different mouse genotypes against the latter. We argue that host evolution can be studied largely irrespective of parasite strains if coupling is absent or weak (*E. ferrisi*) but host-parasite coevolution is more likely observable and

23 best studied in a system with coupled tolerance and resistance (*E. falciformis*).

24 **Keywords:** Resistance, Tolerance, *Eimeria*, Coevolution

## 25 **Introduction**

26 Host defence mechanisms evolve to alleviate the detrimental effect of parasites. They  
27 can be categorised into two components: resistance and tolerance (Råberg et al.,  
28 2009). Resistance is the ability of a host to reduce parasite burden, resulting from  
29 defence against parasite infection or proliferation early after infection  
30 (Schmid-Hempel, 2013). The negative effect of resistance on parasite fitness can  
31 lead to antagonistic coevolution. According to theoretical models, fluctuating host and  
32 parasite genotypes arise, and balancing selection maintains resistance alleles  
33 polymorphic (Boots et al., 2008; Roy & Kirchner, 2000). Resistance has been the  
34 classical "catch all" measure for host-parasite systems, but recently it has been shown  
35 to be incomplete, especially with respect to potential fitness effects on the host  
36 (Kutzer & Armitage, 2016; Råberg et al., 2009).

37 Disease tolerance (not to be confused from "immunological tolerance",  
38 unresponsiveness to self antigens; Medzhitov et al., 2012) is the ability of the host to  
39 limit the impact of parasite on its fitness (Kutzer & Armitage, 2016; Råberg et al.,  
40 2009; Vale & Little, 2012). By potentially providing a longer-living niche, this defence  
41 mechanism improves, or at least does not deteriorate, the fitness of the parasite.  
42 Tolerance alleles are thus predicted by theoretical models to evolve to fixation due to  
43 positive feedback loops (Boots et al., 2008; Restif & Koella, 2004; Roy & Kirchner,  
44 2000). From a mechanistic perspective tolerance alleviates direct or indirect damage

45 (e.g. excessive immune response underlying resistance against parasites, called  
46 immunopathology; Graham et al., 2005) caused by parasites (Råberg et al., 2009).  
47 Tolerance mechanisms include modulation of inflammatory response (Ayres &  
48 Schneider, 2012), tissue repair (stress response, damage repair and cellular  
49 regeneration mechanisms; Soares et al., 2017), and compensation of  
50 parasite-induced damage by increase of reproductive effort (Baucom & de Roode,  
51 2011). The resulting metabolic costs of resistance and tolerance, with and without  
52 parasite infection, determine the optimal (steady state and infection inducible) extent  
53 and of both immune defences (Sheldon & Verhulst, 1996).

54 Resistance and tolerance can be positively associated if they involve the same  
55 metabolic pathway, as was shown in the plant model *Arabidopsis thaliana* in response  
56 against herbivory (Mesa et al., 2017). In animals, genetic association studies of  
57 resistance and tolerance of *Drosophila melanogaster* against the bacterium  
58 *Providencia rettgeri* have shown positively correlated genetic effects, as the same loci  
59 were associated with changes of both traits in the same direction (Howick & Lazzaro,  
60 2017).

61 Nevertheless, resistance and tolerance can also be genetically and physiologically  
62 independent, involving different proximate mechanisms. Lack of correlation between  
63 both defences was shown for example in monarch butterflies (*Danaus plexippus*)  
64 infected by the protozoan parasite *Ophryocystis elektroscirrha*. This study found  
65 genetic variation in resistance between butterflies families, but a fixed tolerance  
66 (Lefèvre et al., 2010). Similarly, no correlation could be found between resistance and  
67 tolerance for the fish *Leuciscus burdigalensis* in response to infection with its parasite

68 *Tracheliastes polycolpus*. The authors explain the decoupling of both defences by the  
69 fact that, in this system, tolerance likely involves wound repair rather than immune  
70 regulation, making resistance and tolerance mechanisms independent (Mazé-Guilmo  
71 et al., 2014).

72 Eventually, in other systems, resistance and tolerance have been found negatively  
73 correlated. For examples, inbred laboratory mouse strains lose weight upon infection  
74 with *Plasmodium chabaudi*. The extent of this impact on host health is negatively  
75 correlated with the peak number of parasites found in the blood (Råberg et al., 2007),  
76 meaning that mouse strains with higher resistance present lower tolerance. Similarly,  
77 infections of sea trout (*Salmo trutta trutta*) and Atlantic salmon (*Salmo salar*) with the  
78 trematode *Diplostomum pseudospathaceum* showed that resistance and tolerance  
79 were negatively correlated when assessing mean levels of both traits in different host  
80 populations (Klemme & Karvonen, 2016). This is interpreted as a result of trade-off  
81 between resistance and tolerance (Råberg et al., 2009; Restif & Koella, 2004;  
82 Sheldon & Verhulst, 1996).

83 We have seen that depending on the system studied resistance and tolerance can be  
84 (1) uncoupled (independent), (2) positively correlated (involving same genes and  
85 mechanisms), or (3) negatively correlated (traded-off). Theoretical models show that  
86 coupling between resistance and tolerance (or absence thereof) depends not only on  
87 the host but also on the parasite (Carval & Ferriere, 2010). This raises the following  
88 question: could there be differences in the resistance-tolerance coupling upon  
89 infection of one host type with two closely related parasite species? To answer this  
90 question, we infected four inbred mouse strains representative of two house mouse

91 subspecies, *M. m. domesticus* and *M. m. musculus*, with three parasite isolates  
92 representative of two naturally occurring parasite species, the protozoan parasite  
93 *Eimeria ferrisi* and *E. falciformis* (Jarquín-Díaz et al., 2019). *Eimeria* spp. are  
94 monoxenous parasites that expand asexually and reproduce sexually in intestinal  
95 epithelial cells, leading to malabsorption of nutrients, tissue damage and weight loss  
96 (Chapman et al., 2013). The evolutionary history of these different *Eimeria* species in  
97 the two house mouse subspecies is unknown and it is unclear whether  
98 subspecies-specific adaptation exists in one or the other.

99 We tested if coupling between resistance and tolerance differs between both parasite  
100 species and discussed the implication for parasite-host coevolution. As coevolving  
101 hosts and parasites can adapt to their local antagonist, we tested in parallel local  
102 adaptation (i.e. a higher parasite fitness in sympatric than in allopatric host (parasite  
103 local adaptation), or a higher host fitness when infected with sympatric than allopatric  
104 parasite (host local adaptation) (Schulte et al., 2011)) of *E. ferrisi* to *Mus musculus*,  
105 using a parasite isolated in a *M. m. domesticus* host and one in a *M. m. musculus*  
106 host. If found, local adaptation would be indirect evidence for coevolution of this  
107 parasite with *Mus musculus*.

## 108 **Material and methods**

### 109 **1. Parasite isolates**

110 The three parasite isolates used in this study were isolated from feces of three different  
111 *M. m. domesticus*/*M. m. musculus* hybrid mice captured in Brandenburg, Germany, in  
112 2016 (capture permit No. 2347/35/2014). The parasite isolates belong to both the most

113 prevalent *Eimeria* species in this area, namely *E. ferrisi* (isolates Brandenburg64 and  
114 Brandenburg139) and *E. falciformis* (isolate Brandenburg88)(Jarquín-Díaz et al., 2019).  
115 Isolate Brandenburg64 was isolated in a 92% *M. m. domesticus* individual (hybrid index  
116 (HI) = 0.08: Proportion of *M. m. musculus* alleles in a set of 14 diagnostic markers, see  
117 Balard et al. (2020)), isolate Brandenburg139 in a 85% *M. m. musculus* (HI=0.85) and  
118 isolate Brandenburg88 in a 80% *M. m. domesticus* (HI=0.2). Pre-patency and the peak  
119 day of parasite shedding for these isolates were estimated during infection in NMRI  
120 laboratory mice (Al-khlifeh et al., 2019) which were also used for serial passaging of  
121 the isolates. Parasite infective forms (oocysts) were recovered by flotation in saturated  
122 NaCl solution followed by washing and observation under light microscope (following  
123 the protocol described in Clerc et al. (2019)) and stored at room temperature in 1mL  
124 of 2% potassium dichromate for a maximum of 2 months before infection of the wild-  
125 derived mouse strains. Oocysts were allowed to sporulate 10 days before infection in  
126 a water bath at 30°C.

## 127 2. Mouse strains

128 We used four wild-derived inbred mouse strains from which we generated four groups  
129 of F1 hybrids. Two parental strains represented *M. m. domesticus*: **SCHUNT**  
130 (Locality: Schweben, Hessen, Germany [N: 50° 26', E: 9° 36'] (Martincová et al.,  
131 2019)) and **STRA** (Locality: Straas, Bavaria, Germany [N: 50° 11', E: 11° 46'] (Piálek  
132 et al., 2008)), and two derived from *M. m. musculus*: **BUSNA** (Locality: Buškovice,  
133 Bohemia, Czech Republic [N: 50° 14', E: 13° 22'] (Piálek et al., 2008)) and **PWD**  
134 (Locality: Kunratice, Bohemia, Czech Republic [N: 50° 01', E: 14° 29'] (Gregorová &  
135 Forejt, 2000)). The four groups of F1 hybrids consisted of two intrasubspecific hybrids

136 (SCHUNTxSTRA and PWDxBUSNA) and two intersubspecific hybrids  
137 (STRAxBUSNA and SCHUNTxPWD)(Figure 1). Age of the mice at the time of  
138 infection ranged between 5.6 and 21.4 weeks. All mouse strains were obtained from  
139 the Institute of Vertebrate Biology of the Czech Academy of Sciences in Studenec  
140 (licence number 61974/2017-MZE-17214; for further details on strains see  
141 <https://housemice.cz/en>).

142 Parasites of the *Eimeria* genus are known to induce host immune protection against  
143 reinfection (Rose et al., 1992; Smith & Hayday, 2000). To ensure that our mice were  
144 *Eimeria*-naive, mouse fecal samples were tested before infection for the presence of  
145 *Eimeria* spp. oocysts by flotation in saturated NaCl solution followed by washing and  
146 observation under light microscope.

### 147 3. Experimental infection

148 Mice were kept in individual cages during infection. Water and food (SNIFF,  
149 Rat/Mouse maintenance feed 10 mm) were provided *ad libitum* supplemented with 1  
150 g of sunflower and barley seeds per day. Mice were orally infected with 150  
151 sporulated oocysts of one *Eimeria* isolate suspended in 100 $\mu$ l phosphate-buffer saline  
152 (PBS) and monitored daily until their sacrifice by cervical dislocation at time of  
153 regression of infection (reduction of oocyst output). Individuals presenting severe  
154 health deficiency and/or a weight loss approaching 18% relative to their starting  
155 weight were sacrificed earlier at defined humane end points (experiment license Reg.  
156 0431/17). Weight was recorded and feces collected on a daily basis. Fecal pellets  
157 were collected every day from each individual cage and suspended in 2% potassium  
158 dichromate. Parasite oocysts were recovered using NaCl flotation (see above).

159 All individuals were negative for *Eimeria* at the beginning of our experiment (before  
160 infection of first batch, as described in the next paragraph). In total, 168 mice were  
161 infected. Mice were randomly allocated to experimental groups ensuring  
162 homogeneous distribution of ages and sexes between groups. Our experiments were  
163 conducted in four (partially overlapping) consecutive batches for logistical reasons.  
164 The first two batches were infected with the two *E. ferrisi* isolates (Brandenburg64 and  
165 Brandenburg139), the third and fourth by one *E. ferrisi* isolate (Brandenburg64) and  
166 one *E. falciformis* isolate (Brandenburg88). Our experimental design is summarized in  
167 **Table 1** (chronology of experimental batches can be scrutinized in **Supplementary**  
168 **Table S1**).

169 Nematode infection is common in breeding facilities (Baker, 1998) and could interact  
170 with *Eimeria* (Clerc et al., 2019). We surveyed for their presence and nematode eggs  
171 were observed in flotated feces of mice belonging to all genotypes before the  
172 experiment. Despite treatment of the first infection batch of mice (B1, 22 mice) with  
173 anthelmintics (Profender®, Bayer AG, Levekusen, Germany) following the protocole  
174 of Mehlhorn et al. (2005), nematodes were still detected with PCR (following the  
175 protocole of Floyd et al. (2005)) in randomly sampled fecal samples a week later. We  
176 therefore decided not to treat mice of the following infection batches. Moreover, we  
177 observed *Eimeria* oocysts in the feces of 28 mice belonging to the last experimental  
178 batch (batch B4) at the day of infection, likely due to cross-contamination between  
179 batches. For following statistical analyses, we considered along with the full data set  
180 (N=168) a conservative data set in which cross-contaminated animals and animals  
181 treated by anthelmintic were removed (N=118). Results obtained on the  
182 conservative data set can be found in **Supplementary Material S2**. Despite



183 differences in significance due to a lower statistical power, the main conclusions of our  
184 analyses were consistent with those obtained on the main data set.

## 185 **4. Statistical analyses**

### 186 **4.1. Choice of proxies for resistance, impact of parasite on host and tolerance**

187 As resistance is the capacity of a host to reduce its parasite burden, it is usually  
188 estimated by the inverse of infection intensity (Råberg et al., 2009). Pre-patency (the  
189 time to shedding of infectious stages, so called oocysts) is longer for *E. falciformis* (7  
190 days) than for *E. ferrisi* (5 days) (Al-khlifeh et al., 2019). Therefore, as a proxy of  
191 (inverse of) resistance we used the number of oocysts per gram of feces (OPG) at the  
192 day of maximal shedding. Using the Spearman's non-parametric rank correlation test,  
193 we found this measurement to be tightly correlated with the sum of oocysts shed  
194 throughout the experiment (Spearman's  $\rho=0.93$ ,  $N=168$ ,  $P<0.001$ ). Due to the  
195 aggregation characteristic of parasites (Shaw & Dobson, 1995), the appropriate  
196 distribution for maximum number of OPG was found to be the negative binomial  
197 distribution. This was confirmed based on log likelihood, AIC criteria and  
198 goodness-of-fits plots (density, CDF, Q-Q, P-P plots; R packages MASS (Venables &  
199 Ripley, 2002) and fitdistrplus (Delignette-Muller & Dutang, 2015)).

200 Both parasite species provoke inflammation, cellular infiltration, enteric lesions,  
201 diarrhea, and ultimately weight loss (Al-khlifeh et al., 2019; Ankrom et al., 1975; Ehret  
202 et al., 2017; Schito et al., 1996). Therefore, the impact of parasites on host health was  
203 measured as the maximum relative weight loss compared to day 0 (body weight  
204 measured at the start of the experimental infection). For mice sacrificed at humane

205 end points before the end of the experiment, last weight of the living animal was used.  
206 This weight (loss) can be expected to be a very conservative estimate for our  
207 analyses (rendering tolerance conservatively low for these animals, which might have  
208 lost more weight if not sacrificed).

209 Tolerance is usually defined as a reaction norm, i.e. the regression slope of host fitness  
210 (or health condition if that is the parameter of interest) on infection intensity per host  
211 genotype (Råberg et al., 2009; Simms, 2000). Thus tolerance was assessed as the  
212 slope of maximum relative weight loss compared to day 0 on number of OPG at the  
213 day of maximal shedding, within each mouse strain and for each parasite isolate. A  
214 steep slope indicates a low tolerance (high weight lost for a given parasite burden).

## 215 **4.2. Statistical modelling**

216 Maximum OPG and relative weight loss were modelled separately as a response of  
217 either mouse strain, parasite isolate and their interaction. We used a negative binomial  
218 generalised linear model for maximum OPG, and a linear model for relative weight loss.  
219 For tolerance, we performed a linear regression with null intercept (as each mouse was  
220 controlled against itself at start of the experiment, before losing weight or shedding  
221 parasite), modelling relative weight loss as a response of maximum OPG interacting  
222 either mouse strain, parasite isolate and their interaction. To test the significance of the  
223 marginal contribution of each parameter to the full model, each parameter was removed  
224 from the full model, and the difference between full and reduced model was assessed  
225 using likelihood ratio tests (G).

226 For each of our model, we also asked within each **parasite isolate** if the response

227 differed between mouse strains using likelihood ratio tests (G) as described above. Of  
228 note, four mice infected by *E. falciformis* isolate Brandenburg88 did not shed any  
229 oocysts as death occurred at or one day before the peak of oocysts shedding in other  
230 mice. For this reason, we modelled maximum OPG for mice infected with this parasite  
231 using a zero-inflated negative binomial (ZINB) generalised linear model, after verifying  
232 that it provided a better fit than the simple negative binomial based on log likelihood  
233 and AIC criteria.

#### 234 4.3. Test of local adaptation

235 Local adaptation of *E. ferrisi* was tested using two isolates (the "Western"  
236 Brandenburg64 and "Eastern" Brandenburg139) and our four parental mouse strains  
237 (the two *M. m. domesticus* Western SCHUNT and STRA, and the two *M. m. musculus*  
238 Eastern BUSNA and PWD). We hypothesised a possible local adaptation of *E. ferrisi*,  
239 i.e. (1) a higher parasite fitness in sympatric than in allopatric host, or (2) a higher host  
240 fitness when infected with sympatric than allopatric parasite. The prediction drawn  
241 from (1) would be that the Eastern parasite (*E. ferrisi* isolate Brandenburg139)  
242 reproduces better in the matching Eastern mouse subspecies (*M. m. musculus*) than  
243 in the allopatric one (*M. m. musculus*), and similarly the Western parasite (*E. ferrisi*  
244 isolate Brandenburg64) reproduce better in *M. m. domesticus* than in *M. m. musculus*.  
245 According to hypothesis (2), a higher tolerance of each host infected by its matching  
246 parasite despite similar parasite reproductive output could indicate increased host  
247 fitness, and host local adaptation.

#### 248 4.4. Test of coupling between resistance and tolerance

249 We tested coupling between resistance and tolerance for *E. ferrisi* and *E. falciformis*  
250 using the isolates Brandenburg64 and Brandenburg88 and our eight mouse strains.  
251 To test such coupling, one can assess the strength of correlation between measure of  
252 resistance and measure of tolerance (Råberg et al., 2007). Of note, tolerance (in  
253 absolute value) is measured as the slope  $\alpha$  of the linear regression of parasite load (x)  
254 on maximum relative weight loss (y) of equation  $y = \alpha x + \beta$  ( $\alpha$  being the slope and  $\beta$   
255 the intercept, 0 in our case). Therefore, tolerance is expressed as  $\alpha = y/x - \beta/x$ . As x  
256 and y/x are by definition not independent, testing the correlation between resistance  
257 and tolerance can lead to spurious correlation (Brett, 2004). To alleviate the dangers  
258 of this statistical artifact, we additionally tested differences in resistance, impact on  
259 health and tolerance between mouse strains separately and also the underlying  
260 correlation between mean parasite load (x) and mean relative weight loss (y). We use  
261 the terminology "coupling" (between resistance and tolerance) to describe  
262 genotype-level correlation between tolerance and resistance additionally supported by  
263 the absence of positive correlation between health-effect and resistance. Correlations  
264 were tested using Spearman's rank correlation.

265 All analyses were performed using R version 3.5.2 (R Development Core Team,  
266 2013)(negative binomial: function glm.nb from R package MASS (Venables & Ripley,  
267 2002); ZIBN: function zeroinfl from R package pscl (Jackman, 2020; Zeileis et al.,  
268 2008); linear model: function lm from R core package stats; mean and 95%  
269 confidence intervals: function ggpredict from R package ggeffect (Lüdecke, 2018)).  
270 Graphics were produced using the R package ggplot2 (Wickham, 2016) and compiled

271 using the free software inkscape (<https://inkscape.org>). Code and data used for this  
272 article can be found at: [https://github.com/alicebalard/Article\\_RelatedParasitesResTol](https://github.com/alicebalard/Article_RelatedParasitesResTol)

## 273 Results

### 274 1. General

275 Parasites of all isolates successfully infected all mouse strains (at the exception of 5  
276 individuals infected by *E. falciformis* isolate Brandenburg88 that died or had to be  
277 sacrificed due to a strong weight loss before the peak of shedding for this parasite),  
278 meaning that no "qualitative infection resistance" (*sensu* Gandon and Michalakis  
279 (2000)) was detected. For *E. ferrisi* (both isolates Brandenburg139 and  
280 Brandenburg64), the pre-patent period was 5 days post infection (dpi) and the median  
281 day of maximal oocyst shedding was 6 dpi (standard deviation sd=0.7 and 0.9,  
282 respectively). The median day of maximum weight loss was 5 dpi for both isolates  
283 (sd=2.1 and 1.7 respectively). For *E. falciformis* (isolate Brandenburg88) pre-patency  
284 was 7 dpi, median day of maximal shedding was 8 dpi (sd=1.3) and median day of  
285 maximal weight loss 9 dpi (sd=1.6)(**Figure 2**). Of note a considerable number of mice  
286 infected with this isolate (13 out of 56 = 23% ) died or had to be sacrificed at humane  
287 end points less than 3 days after the oocysts shedding peak for the group, all  
288 belonging to *M. m. musculus* subspecies (PWD, BUSNA, or their F1 PWDxBUSNA; 5  
289 died at dpi 8, 5 at dpi 9, 3 at dpi 10). *E. falciformis* isolate Brandenburg88 was more  
290 lethal for the *M. m. musculus* mice strains than for the other strains ( $\chi^2_7 = 31.96$ ,  
291  $P < 0.001$ ; **Table 2**).

## 2. No indication of local adaptation of *E. ferrisi*

We tested if our proxies for resistance, impact on weight and tolerance were different between the four parental mouse strains and between both *E. ferrisi* infection isolates (isolate Brandenburg64 and Brandenburg139). Maximum parasite load differed between mouse strains (LRT:  $G=25.5$ ,  $df=6$ ,  $P<0.001$ ), but the interaction term mouse strain-parasite isolate was non significant (LRT:  $G=4.1$ ,  $df=3$ ,  $P=0.25$ ). A similar result was found for maximum relative weight loss (LRT: mouse strain:  $G=16.8$ ,  $df=6$ ,  $P=0.01$ ; interaction mouse strain-parasite isolate:  $G=4.1$ ,  $df=3$ ,  $P=0.25$ ). This indicates that when resistance and impact on weight vary between host strains, they do so independently of the parasite isolate. Eventually, the variables mouse strain, parasite isolate and their interaction were found non significant at the 0.05 threshold for the slope of the linear regression between the two, indicating that differences of tolerance could not be detected between mouse strains or parasite isolates (**Figure 3**). Our results do not indicate either (1) an increased reproduction of each parasite in its matching host or (2) a higher tolerance of host infected by its matching parasite despite similar parasite reproductive output. Thus they do not support the hypothesis of local adaptation between *E. ferrisi* and its host.

## 3. Resistance and tolerance to *E. ferrisi* isolate Brandenburg64 are uncoupled

We tested coupling between resistance and tolerance for *E. ferrisi* isolate Brandenburg64 in our eight mouse strains. First, we tested whether our proxies for resistance, impact on weight and tolerance were different between the four mouse

314 strains. We found the maximum number of OPG and relative weight loss to be  
315 statistically different between mouse strains (LRT: maximum number of OPG:  $G=26.6$ ,  
316  $df=7$ ,  $P<0.001$ ; **Figure 4A**; maximum relative weight loss:  $G=21.5$ ,  $df=7$ ,  $P<0.01$ ;  
317 **Figure 4B**). Tolerance was not found to significantly differ between mouse strains for  
318 this parasite isolate (LRT:  $G=6.8$ ,  $df=7$ ,  $P=0.45$ ; **Figure 4C**).

319 We found a non significant positive correlation between resistance (inverse of maximum  
320 number of OPG) and impact on health (maximum weight loss) (Spearman's  $\rho=0.69$ ,  
321  $P=0.07$ ,  $N=8$ ; **Figure 4D**). Eventually, we did not find a correlation between resistance  
322 (inverse of maximum number of OPG) and tolerance (inverse of slope of maximum  
323 weight loss on maximum OPG) (Spearman's  $\rho=0$ ,  $P=1$ ,  $N=8$ ; **Figure 4E**).

324 In conclusion, we did not find indications of resistance-tolerance coupling for *E. ferrisi*  
325 isolate Brandenburg64, the different mouse strains infected by this parasite presenting  
326 a similar level of tolerance while showing an effect of quantitative resistance on health.

#### 327 **4. Coupling between resistance and tolerance to *E. falciformis***

328 We then tested coupling between resistance and tolerance for *E. falciformis* isolate  
329 Brandenburg88 in our eight mouse strains. First, we tested if our proxies for  
330 resistance, impact on weight and tolerance were different between the four mouse  
331 strains. We found the maximum number of OPG and relative weight loss to be  
332 statistically different between mouse strains (LRT: maximum number of OPG:  $G=28.6$ ,  
333  $df=14$ ,  $P=0.012$ ; **Figure 5A**; maximum relative weight loss:  $G=21$ ,  $df=7$ ,  $P<0.01$ ;  
334 **Figure 5B**). Finally, contrary to our results on *E. ferrisi* isolate Brandenburg64, the  
335 tolerance slopes for *E. falciformis* isolate Brandenburg88 were different between

336 mouse strains (LRT:  $G=13.9$ ,  $df=7$ ,  $P=0.05$ ; **Figure 5C**).

337 We detected a strong negative correlation between (inverse of) resistance (maximum  
338 number of OPG) and tolerance (inverse of slope of maximum weight loss on  
339 maximum OPG) (Spearman's  $\rho=-0.95$ ,  $P=0.001$ ; **Figure 5E**). We conclude that this  
340 correlation is unlikely a statistical artifact, as (1) mouse strains present statistically  
341 different values of resistance and tolerance and (2) we found a (non significant)  
342 negative correlation between resistance (inverse of maximum number of OPG) and  
343 impact on health (maximum weight loss) (Spearman's  $\rho=-0.5$ ,  $P=0.22$ ; **Figure 5D**),  
344 indicating that mouse strains losing more weight also shed less parasites.

345 We conclude that our results indicate the presence of negative resistance-tolerance  
346 coupling for *E. falciformis* isolate Brandenburg88.

## 347 **Discussion**

348 In this study, we assessed resistance and tolerance to two closely related parasites,  
349 *E. ferrisi* (two isolates) and *E. falciformis* (one isolate), in four mouse strains and their  
350 intra-and intersubspecific hybrids. Understanding this coupling has two major  
351 implications.

352 From a practical "measurement" perspective we can ask whether tolerance can be  
353 predicted from resistance, as the latter is easier to measure (e.g. in field sampling).  
354 Many studies assess the impact of parasites on host fitness based on resistance. If,  
355 as we found in the present study, resistance and tolerance are decoupled this can be  
356 misleading. In our host system, the house mice, for example, it has been shown that  
357 hybrids between *M. m. domesticus* and *M. m. musculus* are more resistant to



358 parasites (Baird et al., 2012), including *Eimeria*, but tolerance could not be measured  
359 under natural conditions (Balard et al., 2020). The effect of parasites on host fitness in  
360 the evolution of the house mouse hybrid zone is thus still rather ambiguous (Baird &  
361 Goüy de Bellocq, 2019). We show that careful distinction between parasite species is  
362 necessary when analysing parasite host interaction (see also Jarquín-Díaz et al.,  
363 2019) and that it is indispensable to measure both resistance and tolerance in *Eimeria*  
364 infections of house mice.

365 More generally, in a evolutionary perspective, coupling between resistance and  
366 tolerance might determine whether coevolution between host and parasite can be  
367 expected. As such, coevolution in host-parasite systems is often assumed but rarely  
368 proven (Woolhouse et al., 2002). Janzen (1980) notes that not all parasite-host  
369 systems are coevolving. The presence of efficient host defences against a given  
370 parasite is not necessarily produced in response to this parasite specifically and the  
371 parasite does not necessarily respond specifically. In the mouse-*E. ferrisi* system,  
372 where resistance and tolerance are decoupled, host and parasite fitness might be  
373 decoupled as a result, making host-parasite coevolution less likely. In the  
374 mouse-*E. falciformis* system we found a negative coupling between tolerance and  
375 resistance, making coevolution between host and parasite more likely.

376 Differences between parasite species could explain the evolution of different  
377 strategies: *E. ferrisi* commits to sexual reproduction after a relatively short time with  
378 few cycles of asexual expansion (Al-khlifeh et al., 2019; Ankrom et al., 1975), while  
379 *E. falciformis* has a relatively longer life cycle (Al-khlifeh et al., 2019; Haberkorn,  
380 1970). As *E. ferrisi* infections do not reach extremely high intensities, high tolerance

381 might be the optimal strategy for both house mouse subspecies. Resistance could  
382 then evolve relatively freely without any major impact of the parasite on the hosts'  
383 health. Moreover, our results did not support local adaptation of *E. ferrisi*, which might  
384 be explained by the absence of host-parasite coevolution caused by uncoupling of  
385 parasite and host fitness. In the case of *E. falciformis*, the long life cycle might lead to  
386 high tissue load. Tissue damage is observed during sexual reproduction for this  
387 parasite (Ehret et al., 2017) and might mean that a certain level of resistance is  
388 required. On the other hand, immunopathology has been observed in advanced  
389 *E. falciformis* infections (Stange et al., 2012). These intrinsic characteristics of  
390 *E. falciformis* might lead to multiple different optima for resistance and tolerance,  
391 leading to a trade-off.

392 In addition, we could speculate on two related alternative explanations. Firstly,  
393 *E. falciformis* could originally be a *M. m. domesticus* parasite dissipated into  
394 *M. m. musculus* territory by a spillover through the hybrid zone. Secondly, the  
395 particular *E. falciformis* isolate employed here was collected from a predominantly  
396 *M. m. domesticus* mouse (hybrid index 0.2). The isolate could hence be locally  
397 adapted to *M. m. domesticus*. Experiments with additional *E. falciformis* isolates from  
398 *M. m. musculus* are needed to test whether host subspecies adaptation can lead to  
399 high tolerance and low resistance in matching pairs of *E. falciformis* isolates and  
400 mouse subspecies. This seems plausible, as the coupling between resistance and  
401 tolerance links host and parasite fitness, making coevolution and hence local  
402 adaptation more likely. Interestingly, this parasite-host coevolution wouldn't be  
403 antagonistic but rather mutualistic with regards to tolerance and parasite reproduction  
404 (that is, the inverse of resistance) (Little et al., 2010; Råberg et al., 2009).

405 Alternatively, though descriptive, molecular-genetic analyses of diagnostic marker can  
406 be used to infer coevolutionary pathways between host and their parasites (e.g.  
407 Goüy de Bellocq et al., 2018; Kváč et al., 2013).

408 In conclusion, we argue that the difference between resistance and tolerance coupling  
409 in two different parasites can guide research in the house mouse system: if the effects  
410 of host hybridisation should be studied independently of potential host-parasite  
411 coadaptation, the prevalent *E. ferrisi* might be the most suitable parasite. If  
412 coevolution between hosts and parasites should be studied, the pathogenic  
413 *E. falciformis* is a more plausible target. Generally, the coupling between resistance  
414 and tolerance can differ between closely related parasite species and we argue that  
415 this trait of a host-parasite system determines the questions to be best approached  
416 with a particular parasite.

## 417 References

- 418 Al-khlifeh, E., Balard, A., Jarquín-Díaz, V. H., Weyrich, A., Wibbelt, G. & Heitlinger, E. (2019). *Eimeria*  
419 *falciformis* BayerHaberKorn1970 and novel wild derived isolates from house mice: Differences in  
420 parasite lifecycle, pathogenicity and host immune reactions. *bioRxiv*, 611277. doi:[10.1101/611277](https://doi.org/10.1101/611277)  
421 Ankrom, S. L., Chobotar, B. & Ernst, J. V. (1975). Life cycle of *Eimeria ferrisi* Levine & Ivens, 1965  
422 in the mouse, *Mus musculus*. *The Journal of Protozoology*, 22, 317–323. doi:[10.1111/j.1550-](https://doi.org/10.1111/j.1550-7408.1975.tb05177.x)  
423 [7408.1975.tb05177.x](https://doi.org/10.1111/j.1550-7408.1975.tb05177.x)  
424 Ayres, J. S. & Schneider, D. S. (2012). Tolerance of infections. *Annual Review of Immunology*, 30, 271–  
425 294. doi:[10.1146/annurev-immunol-020711-075030](https://doi.org/10.1146/annurev-immunol-020711-075030)  
426 Baird, S. J. E. & Goüy de Bellocq, J. (2019). Shifting paradigms for studying parasitism in hybridising  
427 hosts: Response to Theodosopoulos, Hund, and Taylor. *Trends in ecology & evolution*, 34, 387–  
428 389. doi:[doi.org/10.1016/j.tree.2019.01.011](https://doi.org/10.1016/j.tree.2019.01.011)

429 Baird, S. J. E., Ribas, A., Macholán, M., Albrecht, T., Piálek, J. & Goüy de Bellocq, J. (2012). Where  
 430 are the wormy mice? A reexamination of hybrid parasitism in the European house mouse hybrid  
 431 zone. *Evolution*, 66, 2757–2772. doi:[10.1111/j.1558-5646.2012.01633.x](https://doi.org/10.1111/j.1558-5646.2012.01633.x)

432 Balard, A., Jarquín-Díaz, V. H., Jost, J., Martincová, I., Ďureje, L., Piálek, J., Macholán, M., de Bellocq,  
 433 J. G., Baird, S. J. E. & Heitlinger, E. (2020). Intensity of infection with intracellular *Eimeria* spp.  
 434 and pinworms is reduced in hybrid mice compared to parental subspecies. *Journal of Evolutionary*  
 435 *Biology*, 33, 435–448. doi:[10.1111/jeb.13578](https://doi.org/10.1111/jeb.13578)

436 Baucom, R. S. & de Roode, J. C. (2011). Ecological immunology and tolerance in plants and animals.  
 437 *Functional Ecology*, 25, 18–28. doi:[10.1111/j.1365-2435.2010.01742.x](https://doi.org/10.1111/j.1365-2435.2010.01742.x)

438 Boots, M., Best, A., Miller, M. R. & White, A. (2008). The role of ecological feedbacks in the evolution  
 439 of host defence: What does theory tell us? *Philosophical Transactions of the Royal Society B:*  
 440 *Biological Sciences*, 364, 27–36. doi:[10.1098/rstb.2008.0160](https://doi.org/10.1098/rstb.2008.0160)

441 Brett, M. T. (2004). When is a correlation between non-independent variables “spurious”? *Oikos*, 105,  
 442 647–656. doi:[10.1111/j.0030-1299.2004.12777.x](https://doi.org/10.1111/j.0030-1299.2004.12777.x)

443 Carval, D. & Ferriere, R. (2010). A unified model for the coevolution of resistance, tolerance, and  
 444 virulence. *Evolution*, 64, 2988–3009. doi:[10.1111/j.1558-5646.2010.01035.x](https://doi.org/10.1111/j.1558-5646.2010.01035.x)

445 Chapman, H. D., Barta, J. R., Blake, D., Gruber, A., Jenkins, M., Smith, N. C., Suo, X. & Tomley, F. M.  
 446 (2013). Chapter two - a selective review of advances in coccidiosis research. 83, 93–171. doi:[10.1016/B978-0-12-407705-8.00002-1](https://doi.org/10.1016/B978-0-12-407705-8.00002-1)

447 [1016/B978-0-12-407705-8.00002-1](https://doi.org/10.1016/B978-0-12-407705-8.00002-1)

448 Clerc, M., Fenton, A., Babayan, S. A. & Pedersen, A. B. (2019). Parasitic nematodes simultaneously  
 449 suppress and benefit from coccidian coinfection in their natural mouse host. *Parasitology*, 146,  
 450 1096–1106. doi:[10.1017/S0031182019000192](https://doi.org/10.1017/S0031182019000192)

451 Delignette-Muller, M. L. & Dutang, C. (2015). Fitdistrplus: An r package for fitting distributions. *Journal of*  
 452 *Statistical Software*, 64, 1–34. doi:[10.18637/jss.v064.i04](https://doi.org/10.18637/jss.v064.i04)

453 Ďureje, L., Macholán, M., Baird, S. J. E. & Piálek, J. (2012). The mouse hybrid zone in central europe:  
 454 From morphology to molecules. *Journal of Vertebrate Biology*, 61, 308–318. doi:[10.25225/fozo.](https://doi.org/10.25225/fozo.v61.i3.a13.2012)  
 455 [v61.i3.a13.2012](https://doi.org/10.25225/fozo.v61.i3.a13.2012)

456 Ehret, T., Spork, S., Dieterich, C., Lucius, R. & Heitlinger, E. (2017). Dual RNA-seq reveals no plastic  
 457 transcriptional response of the coccidian parasite *Eimeria falciformis* to host immune defenses.  
 458 *BMC Genomics*, 18, 686. doi:[10.1186/s12864-017-4095-6](https://doi.org/10.1186/s12864-017-4095-6)  
 459 Floyd, R. M., Rogers, A. D., Lamshead, P. J. D. & Smith, C. R. (2005). Nematode-specific PCR primers  
 460 for the 18S small subunit rRNA gene. *Molecular Ecology Notes*, 5, 611–612. doi:[10.1111/j.1471-8286.2005.01009.x](https://doi.org/10.1111/j.1471-8286.2005.01009.x)  
 461  
 462 Gandon, S. & Michalakis, Y. (2000). Evolution of parasite virulence against qualitative or quantitative  
 463 host resistance. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 267,  
 464 985–990. doi:[10.1098/rspb.2000.1100](https://doi.org/10.1098/rspb.2000.1100)  
 465 Goüy de Bellocq, J., Wasimuddin, Ribas, A., Bryja, J., Piálek, J. & Baird, S. J. E. (2018). Holobiont suture  
 466 zones: Parasite evidence across the European house mouse hybrid zone. *Molecular Ecology*.  
 467 doi:[10.1111/mec.14938](https://doi.org/10.1111/mec.14938)  
 468 Graham, A. L., Allen, J. E. & Read, A. F. (2005). Evolutionary causes and consequences of  
 469 immunopathology. *Annual Review of Ecology, Evolution, and Systematics*, 36, 373–397.  
 470 doi:[10.1146/annurev.ecolsys.36.102003.152622](https://doi.org/10.1146/annurev.ecolsys.36.102003.152622)  
 471 Gregorová, S. & Forejt, J. (2000). PWD/Ph and PWK/Ph inbred mouse strains of *Mus m. musculus*  
 472 subspecies—a valuable resource of phenotypic variations and genomic polymorphisms. *Folia*  
 473 *Biologica*, 46, 31–41.  
 474 Haberkorn, A. (1970). Die Entwicklung von *Eimeria falciformis* (Eimer 1870) in der weißen Maus  
 475 (*Mus musculus*). *Zeitschrift für Parasitenkunde*, 34, 49–67. doi:[10.1007/BF00629179](https://doi.org/10.1007/BF00629179)  
 476 Howick, V. M. & Lazzaro, B. P. (2017). The genetic architecture of defence as resistance to and tolerance  
 477 of bacterial infection in *Drosophila melanogaster*. *Molecular Ecology*, 26, 1533–1546. doi:[10.1111/mec.14017](https://doi.org/10.1111/mec.14017)  
 478  
 479 Jackman, S. (2020). *pscl: Classes and methods for R developed in the political science computational*  
 480 *laboratory*. United States Studies Centre, University of Sydney. Sydney, New South Wales,  
 481 Australia.  
 482 Janzen, D. H. (1980). When is it coevolution? *Evolution*, 34, 611–612. doi:[10.1111/j.1558-5646.1980.tb04849.x](https://doi.org/10.1111/j.1558-5646.1980.tb04849.x)  
 483

484 Jarquín-Díaz, V. H., Balard, A., Jost, J., Kraft, J., Dikmen, M. N., Kvičerová, J. & Heitlinger, E. (2019).  
 485 Detection and quantification of house mouse *Eimeria* at the species level – Challenges and  
 486 solutions for the assessment of coccidia in wildlife. *International Journal for Parasitology:  
 487 Parasites and Wildlife*, 10, 29–40. doi:[10.1016/j.ijppaw.2019.07.004](https://doi.org/10.1016/j.ijppaw.2019.07.004)  
 488 Klemme, I. & Karvonen, A. (2016). Vertebrate defense against parasites: Interactions between  
 489 avoidance, resistance, and tolerance. *Ecology and Evolution*, 7, 561–571.  
 490 doi:[10.1002/ece3.2645](https://doi.org/10.1002/ece3.2645)  
 491 Kutzer, M. A. M. & Armitage, S. A. O. (2016). Maximising fitness in the face of parasites: A review of host  
 492 tolerance. *Zoology*, 119, 281–289. doi:[10.1016/j.zool.2016.05.011](https://doi.org/10.1016/j.zool.2016.05.011)  
 493 Kváč, M., McEvoy, J., Loudová, M., Stenger, B., Sak, B., Květoňová, D., Ditrich, O., Rašková, V., Moriarty,  
 494 E., Rost, M., Macholán, M. & Piálek, J. (2013). Coevolution of *Cryptosporidium tyzzeri* and the  
 495 house mouse (*Mus musculus*). *International Journal for Parasitology*, 43, 805–817. doi:[10.1016/  
 496 j.ijpara.2013.04.007](https://doi.org/10.1016/j.ijpara.2013.04.007)  
 497 Lefèvre, T., Williams, A. J. & de Roode, J. C. (2010). Genetic variation in resistance, but not tolerance,  
 498 to a protozoan parasite in the monarch butterfly. *Proceedings of the Royal Society B: Biological  
 499 Sciences*, 278, 751–759. doi:[10.1098/rspb.2010.1479](https://doi.org/10.1098/rspb.2010.1479)  
 500 Little, T. J., Shuker, D. M., Colegrave, N., Day, T. & Graham, A. L. (2010). The coevolution of virulence:  
 501 Tolerance in perspective. *PLoS pathogens*, 6. doi:[10.1371/journal.ppat.1001006](https://doi.org/10.1371/journal.ppat.1001006)  
 502 Lüdtke, D. (2018). Ggeffects: Tidy data frames of marginal effects from regression models. *Journal of  
 503 Open Source Software*, 3, 772. doi:[10.21105/joss.00772](https://doi.org/10.21105/joss.00772)  
 504 Macholán, M., Baird, S. J. E., Fornůsková, A., Martincová, I., Rubík, P., Ďureje, L., Heitlinger, E. & Piálek,  
 505 J. (2019). Widespread introgression of the *Mus musculus musculus* Y chromosome in Central  
 506 Europe. *bioRxiv*. doi:[10.1101/2019.12.23.887471](https://doi.org/10.1101/2019.12.23.887471)  
 507 Martincová, I., Ďureje, L., Kreisinger, J., Macholán, M. & Piálek, J. (2019). Phenotypic effects of the  
 508 Y chromosome are variable and structured in hybrids among house mouse recombinant lines.  
 509 *Ecology and Evolution*, 9, 6124–6137. doi:[10.1002/ece3.5196](https://doi.org/10.1002/ece3.5196)  
 510 Mazé-Guilmo, E., Loot, G., Páez, D. J., Lefèvre, T. & Blanchet, S. (2014). Heritable variation in host  
 511 tolerance and resistance inferred from a wild host–parasite system. *Proceedings of the Royal  
 512 Society B: Biological Sciences*, 281, 20132567. doi:[10.1098/rspb.2013.2567](https://doi.org/10.1098/rspb.2013.2567)

513 Medzhitov, R., Schneider, D. S. & Soares, M. P. (2012). Disease tolerance as a defense strategy. *Science*,  
514 335, 936–941. doi:[10.1126/science.1214935](https://doi.org/10.1126/science.1214935)

515 Mesa, J. M., Scholes, D. R., Juvik, J. A. & Paige, K. N. (2017). Molecular constraints on resistance–  
516 tolerance trade-offs. *Ecology*, 98, 2528–2537. doi:[10.1002/ecy.1948](https://doi.org/10.1002/ecy.1948)

517 Piálek, J., Vyskočilová, M., Bímová, B., Havelková, D., Piálková, J., Dufková, P., Bencová, V., Ďureje, L.,  
518 Albrecht, T., Hauße, H. C., Macholán, M., Munclinger, P., Storchová, R., Zajíčková, A., Holáň, V.,  
519 Gregorová, S. & Forejt, J. (2008). Development of unique house mouse resources suitable for  
520 evolutionary studies of speciation. *Journal of Heredity*, 99, 34–44. doi:[10.1093/jhered/esm083](https://doi.org/10.1093/jhered/esm083)

521 R Development Core Team. (2013). *R: A language and environment for statistical computing*.  
522 <http://www.R-project.org/>. R Foundation for Statistical Computing. Vienna, Austria.

523 Råberg, L., Graham, A. L. & Read, A. F. (2009). Decomposing health: Tolerance and resistance to  
524 parasites in animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*,  
525 364, 37–49. doi:[10.1098/rstb.2008.0184](https://doi.org/10.1098/rstb.2008.0184)

526 Råberg, L., Sim, D. & Read, A. F. (2007). Disentangling genetic variation for resistance and tolerance to  
527 infectious diseases in animals. *Science*, 318, 812–814. doi:[10.1126/science.1148526](https://doi.org/10.1126/science.1148526)

528 Restif, O. & Koella, J. C. (2004). Concurrent evolution of resistance and tolerance to pathogens. *The*  
529 *American Naturalist*, 164, E90–E102. doi:[10.1086/423713](https://doi.org/10.1086/423713)

530 Rose, M. E., Hesketh, P. & Wakelin, D. (1992). Immune control of murine coccidiosis: CD4+ and CD8+  
531 T lymphocytes contribute differentially in resistance to primary and secondary infections.  
532 *Parasitology*, 105, 349–354. doi:[10.1017/S0031182000074515](https://doi.org/10.1017/S0031182000074515)

533 Roy, B. A. & Kirchner, J. W. (2000). Evolutionary dynamics of pathogen resistance and tolerance.  
534 *Evolution*, 54, 51–63. doi:[10.1111/j.0014-3820.2000.tb00007.x](https://doi.org/10.1111/j.0014-3820.2000.tb00007.x)

535 Schito, M. L., Barta, J. R. & Chobotar, B. (1996). Comparison of four murine *Eimeria* species in  
536 immunocompetent and immunodeficient mice. *The Journal of Parasitology*, 82, 255–262.  
537 doi:[10.2307/3284157](https://doi.org/10.2307/3284157)

538 Schmid-Hempel, P. (2013). *Evolutionary parasitology: The integrated study of infections, immunology,*  
539 *ecology, and genetics*. Oxford University Press. doi:[10.1093/acprof:oso/9780199229482.001.](https://doi.org/10.1093/acprof:oso/9780199229482.001.0001)  
540 [0001](https://doi.org/10.1093/acprof:oso/9780199229482.001.0001)



541 Schulte, R. D., Makus, C., Hasert, B., Michiels, N. K. & Schulenburg, H. (2011). Host–parasite local  
 542 adaptation after experimental coevolution of *Caenorhabditis elegans* and its microparasite *Bacillus*  
 543 *thuringiensis*. *Proceedings of the Royal Society B: Biological Sciences*, 278, 2832–2839. doi:[10.](https://doi.org/10.1098/rspb.2011.0019)  
 544 [1098/rspb.2011.0019](https://doi.org/10.1098/rspb.2011.0019)  
 545 Shaw, D. J. & Dobson, A. P. (1995). Patterns of macroparasite abundance and aggregation in wildlife  
 546 populations: A quantitative review. *Parasitology*, 111, S111–S133.  
 547 doi:[10.1017/S0031182000075855](https://doi.org/10.1017/S0031182000075855)  
 548 Sheldon, B. C. & Verhulst, S. (1996). Ecological immunology: Costly parasite defences and trade-offs in  
 549 evolutionary ecology. *Trends in ecology & evolution*, 11, 317–321.  
 550 Simms, E. L. (2000). Defining tolerance as a norm of reaction. *Evolutionary Ecology*, 14, 563–570. doi:[10.](https://doi.org/10.1023/a:1010956716539)  
 551 [1023/a:1010956716539](https://doi.org/10.1023/a:1010956716539)  
 552 Smith, A. L. & Hayday, A. C. (2000). Genetic Dissection of primary and secondary responses to a  
 553 widespread natural pathogen of the gut, *Eimeria vermiformis*. *Infection and Immunity*, 68,  
 554 6273–6280. doi:[10.1128/IAI.68.11.6273-6280.2000](https://doi.org/10.1128/IAI.68.11.6273-6280.2000)  
 555 Soares, M. P., Teixeira, L. & Moita, L. F. (2017). Disease tolerance and immunity in host protection against  
 556 infection. *Nature Reviews Immunology*, 17, 83–96. doi:[10.1038/nri.2016.136](https://doi.org/10.1038/nri.2016.136)  
 557 Stange, J., Hepworth, M. R., Rausch, S., Zajic, L., Kühl, A. A., Uyttenhove, C., Renauld, J.-C., Hartmann,  
 558 S. & Lucius, R. (2012). IL-22 mediates host defense against an intestinal intracellular parasite in  
 559 the absence of IFN- $\gamma$  at the cost of Th17-driven immunopathology. *Journal of Immunology*, 188,  
 560 2410–2418. doi:[10.4049/jimmunol.1102062](https://doi.org/10.4049/jimmunol.1102062)  
 561 Vale, P. F. & Little, T. J. (2012). Fecundity compensation and tolerance to a sterilizing pathogen in daphnia.  
 562 *Journal of Evolutionary Biology*, 25, 1888–1896. doi:[10.1111/j.1420-9101.2012.02579.x](https://doi.org/10.1111/j.1420-9101.2012.02579.x)  
 563 Venables, W. N. & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.). New York, NY: Springer.  
 564 doi:[10.1007/978-0-387-21706-2](https://doi.org/10.1007/978-0-387-21706-2)  
 565 Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis (second edition)*. New York, NY:  
 566 Springer. doi:[10.1007/978-0-387-98141-3](https://doi.org/10.1007/978-0-387-98141-3)  
 567 Woolhouse, M. E. J., Webster, J. P., Domingo, E., Charlesworth, B. & Levin, B. R. (2002). Biological and  
 568 biomedical implications of the co-evolution of pathogens and their hosts. *Nature Genetics*, 32,  
 569 569–577. doi:[10.1038/ng1202-569](https://doi.org/10.1038/ng1202-569)



570 Zeileis, A., Kleiber, C. & Jackman, S. (2008). Regression models for count data in R. *Journal of Statistical*  
 571 *Software*, 27. doi:[10.18637/jss.v027.i08](https://doi.org/10.18637/jss.v027.i08)

## 572 Tables


Mouse		<i>Eimeria</i>		
strains	subspecies	<i>E. ferrisi</i> Brandenburg139	<i>E. ferrisi</i> Brandenburg64	<i>E. falciformis</i> Brandenburg88
SCHUNT	F0 <i>M.m.domesticus</i>	7 (5M / 2F)	14 (6M / 8F)	6 (3M / 3F)
STRA	F0 <i>M.m.domesticus</i>	6 (2M / 4F)	15 (8M / 7F)	7 (4M / 3F)
SCHUNTxSTRA	F1 <i>M.m.domesticus</i>		6 (2M / 4F)	8 (5M / 3F)
STRAxBUSNA	F1 Hybrid		8 (5M / 3F)	8 (3M / 5F)
SCHUNTxPWD	F1 Hybrid		8 (3M / 5F)	6 (4M / 2F)
PWDxBUSNA	F1 <i>M.m.musculus</i>		9 (4M / 5F)	7 (4M / 3F)
BUSNA	F0 <i>M.m.musculus</i>	6 (2M / 4F)	14 (8M / 6F)	7 (3M / 4F)
PWD	F0 <i>M.m.musculus</i>	6 (3M / 3F)	13 (10M / 3F)	7 (1M / 6F)

**Table 1.** Infection experiment design.


Mouse			
subspecies	strains	status at dpi 11	
		alive	dead
Mmd	SCHUNT	6	0
Mmd	STRA	7	0
Mmd	SCHUNTxSTRA	8	0
Mmd-Mmm	STRAxBUSNA	8	0
Mmd-Mmm	SCHUNTxPWD	6	0
Mmm	PWDxBUSNA	4	3
Mmm	BUSNA	3	4
Mmm	PWD	1	6
	<b>total</b>	<b>43</b>	<b>13</b>

**Table 2.** Contingency table: number of mice and status at dpi 11 for each mouse strain upon infection with *E. falciformis* isolate Brandenburg88.


## 573 Figures




**Figure 1: Parasite isolates and mouse strains.** (A) Map showing locations at which mice were collected for breeding of mouse strains and isolation of parasites. The purple line is an estimation of the center of the house mouse hybrid zone between *M. m. domesticus* and *M. m. musculus* based on sampling and genotyping of mice in this area (Balard et al., 2020; Ďureje et al., 2012; Macholán et al., 2019). (B) The eight mouse groups (parents and F1s) used in our experimental infections.




**Figure 2: Parasite density (A) and relative weight (B) during *Eimeria* infection.** Parasite density is calculated as number of oocysts detected (in millions) per gram of feces, relative weight is calculated as the percentage of weight compared to day 0. Mean and 95% CI are plotted for each parasite isolate. All hosts strains are pooled together.



**Figure 3: Comparison of resistance, impact on weight and tolerance between mouse strain for both *Eimeria ferrisi* isolates.** (A) Maximum oocysts per gram of feces used as a proxy for (inverse of) resistance; (B) Impact on host health measured as the maximum weight loss during patent period relative to starting weight (%); (C) Tolerance estimated by the slope of the linear regression with null intercept modelling maximum relative weight loss as a response of maximum oocysts per gram of feces. A steep slope corresponds to a low tolerance. We did not detect (A) either higher parasite shedding of the Eastern parasite isolate in Eastern mouse strains and vice versa or (C) higher tolerance of Eastern hosts infected by Eastern parasite isolate and vice versa, thus our results do not support the hypothesis of local adaptation between *E. ferrisi* and its host.



**Figure 4: No indication of resistance-tolerance coupling for *E. ferrisi* isolate Brandenburg64.** Colors represent mouse subspecies (blue: *M. m. domesticus*, red: *M. m. musculus*, purple: Mmd-Mmm). Left side: comparison of maximum oocysts per gram of feces used as a proxy for (inverse of) resistance (A), impact on weight measured as the maximum weight loss during patent period relative to starting weight (B) and tolerance between mouse groups estimated by the slope of the linear regression with null intercept modelling maximum relative weight loss as a response of maximum oocysts per gram of feces, a steep slope corresponding to a low tolerance (C). Maximum number of OPG and relative weight loss differ between mouse groups, but tolerance is similar. Right side: non significant positive correlation between mean maximum oocysts per gram of feces and mean relative weight loss (D) and absence of correlation between maximum oocysts per gram of feces used as a proxy for (inverse of) resistance and tolerance (E); Grey error bars represent 95% confidence intervals. Our results do not support coupling between resistance and tolerance *E. ferrisi* isolate Brandenburg64.



**Figure 5: Coupling between resistance and tolerance for *E. falciformis* isolate Brandenburg88.** Colors represent mouse subspecies (blue: *M. m. domesticus*, red: *M. m. musculus*, purple: Mmd-Mmm). Left side: comparison of maximum oocysts per gram of feces used as a proxy for (inverse of) resistance (A), impact on weight measured as the maximum weight loss during patent period relative to starting weight (B) and tolerance between mouse groups estimated by the slope of the linear regression with null intercept modelling maximum relative weight loss as a response of maximum oocysts per gram of feces, a steep slope corresponding to a low tolerance (C). Maximum number of OPG, relative weight loss and tolerance differ between mouse groups. Right side: non significant negative correlation between mean maximum oocysts per gram of feces and mean relative weight loss (D) and strong negative correlation between maximum oocysts per gram of feces used as a proxy for (inverse of) resistance and tolerance (E); Grey error bars represent 95% confidence intervals. Our results support coupling between resistance and tolerance *E. falciformis* isolate Brandenburg88.