

Contents

| | |
|--|-----------|
| Abstract | 2 |
| Introduction | 3 |
| Functional traits and niche differentiation | 5 |
| Range shifts and response to climate change | 6 |
| Latitudinal gradients as field laboratories | 8 |
| Predictions and hypotheses | 9 |
| Methods | 12 |
| Sites | 12 |
| Species Selection | 13 |
| Competition and community data | 13 |
| Climate data | 14 |
| Functional Traits | 15 |
| Statistical analyses | 16 |
| Results | 18 |
| Figures | 21 |
| Range Maps | 23 |
| Discussion | 37 |
| Competition across latitudinal and climatic range limits | 37 |
| Suites of functional traits across ranges | 40 |
| Areas of future research | 41 |
| Reference | 44 |

Abstract

Plant species' ranges are determined through biotic and abiotic factors such as competition and the level of environmental stress (e.g., from drought or cold). Tree species in broadleaf temperate forests in the Northeast face changes to these ecological limits due to shifting temperatures across their range. The effect of this shift is predicted to be most apparent at species' range limits, where such a change in conditions—that previously halted population expansion—can cause a range shift. By examining species at their range limits, we can predict their future responses to climate change and the implications for their distribution and success. The goal of this study was to determine how species richness and composition of competitive neighborhoods in broadleaf temperate forests in the Northeast change across a latitudinal and climatic gradient. I examined whether focal deciduous tree species face a competitive disadvantage at their range limit, represented by the competitive index of relative basal area of trees larger than the focal individual (a distance-independent strategy). To accomplish this, I recorded the size and community composition of seven deciduous tree species at their range limits, varying from the northeastern United States to southeastern Quebec. Additionally, I examined the relationship between the competitiveness of a species at its limits and the variation in its functional traits, using previously-collected data on the functional traits of the same seven species. Using convex hull values for each species at each site, I examined whether the variation in the suites of functional traits for each species points to evidence of habitat filtering or niche differentiation. This experiment can thus provide valuable insight into how the species will respond to changing temperatures in the future by determining the relative influence of biotic and abiotic factors on each species at their range extremes.

Introduction

A fundamental aim of community ecology is to determine the rules by which a community assembles. Community assembly is an ecological sorting process that filters out some species, but allows others to survive (Ackerly 2003). Studies into species distributions shed light on what factors specifically cause the limitations of particular species, with the larger goal of understanding the overarching rules dictating community assembly. Two general processes have been proposed: (1) habitat filtering that limits species' life strategies, and (2) resource partitioning that limits similarity in co-occurring species (Begon et. al 1986; Cornwell and Ackerly 2009). Fundamentally, the rules of community assembly can be described by examining the species that appear in a given community, and their interactions with their environment.

The concept of a species' niche is useful in describing both intra- and inter-specific interactions, as well as in illustrating the effect the environment may have on a species. In Hutchinsonian niche space, there are two types of niche. The fundamental niche acknowledges the environment a species may tolerate, while the realized niche describes the performance of a species in the presence of competitors. The final distribution of a species and its community composition results from the interplay of many abiotic (i.e. environmental) and biotic (e.g. competition) factors (Ackerly 2003).

Species distribution and ranges serve as an actualization of the concept of a niche. Indeed, understanding the sorting processes that prevent a species from inhabiting space across continuous landscapes has long been the subject of research. Seminal works have theorized about the different causes underlying the filtering out of a species at its range limit (Darwin 1859). More specifically, the two limitations imposed on species in determining their ranges are theorized as (1) abiotically stressful environments that limit species at high latitudes and altitudes and (2) species interactions that further impose restrictions on the species through competition (Louhan et. al, 2015; Pigot & Tobias, 2013). On the edges of

its range, a species sees either biotic and/or abiotic factors forcing it out in a process called habitat filtering (Weiher & Keddy 1998). Range limits are then predetermined by a species' environment and the strength of its surrounding biotic factors (Pigot & Tobias 2013; Sexton et. al 2009).

It is currently uncertain to what extent range limits are caused by biotic versus abiotic factors (Sexton et. al 2009; Hargreaves et. al 2014). Darwin long ago theorized that abiotic factors constrain species at high latitudes, high elevation, and likewise environmentally stressful areas. Meanwhile, community interactions limit species in less stressful environments (Louthan et. al 2015). In a meta-analysis of both elevational and latitudinal range studies, Hargreaves et. al (2004) found that studies conducted with latitudinal range limits showed a pattern that upper range limits were governed solely by abiotic factors. However, there were too few studies conducted at lower latitudes to ascertain a concrete alternative behavior at the lower limits. Thus, it is likely that biotic interactions will increase in importance in governing range limits as the climate warms (Hargreaves et. al 2014).

The relative importance of biotic factors and abiotic factors varies greatly among species, complicating our ability to predict how a species within a given community will respond to a warming environment. Additionally, there is a dearth of research looking at both biotic and abiotic factors for the same species, thus controlling for variation among species (Sexton et. al 2009). Examining a species at its range edge compared to its range interior, taking note of both the competitive niche it occupies and the environment the species lives in, thus provides important information on the interplay of factors limiting it. By understanding the factors limiting a species, we can predict more accurately how well a species will be able to extend its realized niche in a changing environment.

Functional traits and niche differentiation

Certain traits in plants have long been theorized as a major determining factor in a plant species' success in different landscapes, as well as in the ability of species to co-exist (Schimper 1898). In Schimper's theory, the success of different plant species across landscapes is due to different quantitative traits, such as leaf nitrogen and carbon concentrations, wood densities, leaf sizes, and potential canopy heights. Such traits are dubbed plant functional traits. Although they are far from a recent concept, functional traits have seen a resurgence in focus among community ecologists (Pigot and Tobias 2013), so much so that McGill et. al (2006) suggests rebuilding the entire study of community ecology with an emphasis on functional traits as a core concept.

Phenotypic differences among species impact diversity in communities (MacArthur and Levins 1967). In the rules of community assembly, functional traits provide insight into co-occurring species' ecological strategies. McGill et. al (2006) posit that functional traits are one of four major themes governing the composition of a species' fundamental and realized niches (the others being environmental gradients, interaction milieu, and performance currency). Functional traits have become a useful tool in understanding the nature of the relationship between plants within their communities, as well as understanding the relationship between environment and community assembly. By examining functional traits, we can more fully understand the resource partitioning within niches.

Functional traits also serve as useful predictors of other traits that are more difficult to measure, but that may provide better projections of success. For example, certain seed sizes and shapes strongly correlate with persistence of the seed in soil (Thompson et. al 1993). These studies have seen subtle differences in traits in co-occurring species, theorizing that these differences contribute to species coexistence. Additionally, a large amount of intra-specific variation in functional traits may reduce the effect of phylogenetic relatedness in nearby species (i.e. limiting similarity) (Burns & Strauss 2012). Kraft et. al (2008) showed

that species with a broad distribution of trait values co-occur more often than expected by a null model, indicating that variable trait patterns enable more co-occurrence of species within a community. Similarly, certain suites of functional traits can enable individuals to be more successful than others of the same species under environmental conditions that favor those traits (Jung et. al 2010). It follows that in a changing environment, certain functional traits may prove to be better suited to the new conditions.

Functional traits have also been shown to be accurate predictors of species' geographic ranges (Stahl & Wirth 2014), lending credence to the theory of niche differentiation and resource partitioning as a primary factor in determining a species' geographic range. Intra-specific variation in physiological traits may give species the chance to survive in the face of rapid climatic changes within their range (Davis & Shaw, 2001; Nussey et al., 2005; Skelly et al., 2007). The functional traits of a species across its range can thus provide valuable information about the biotic and abiotic factors that limit it beyond its range, and how it is responding to those limiting factors. By using functional traits to describe a plant's resource partitioning across its geographic range, I can more accurately predict how a species could respond to shifts in climate.

Range shifts and response to climate change

In particular, studies into range limits have come to the foreground in attempts to understand how these ranges might shift with changes in climate. Observing a species at its range edge can help test the conditions by which some populations adapt successfully (or not) to novel conditions. Based on the niche conservatism hypothesis, when environmental conditions stretch beyond a species' physiological tolerances, that species will either shift its distribution or face extinction (Peterson, Soberón, & Sánchez-Cordero 1999). More specifically, in response to a changing environment, a population may respond one of a few ways: 1) small-scale changes in distributions of local populations; 2) large-scale shifts in geographic distribution,

consisting of range expansions and range contraction in the former range; 3) adaptive evolution of the niche in response to the new conditions; and 4) extinction (Ackerly 2003; Thomas, Franco & Hill 2006).

Species ranges are very mobile, often contracting or expanding over time (Brown, Stevens & Kaufman 1996; Davis & Shaw 2001). Indeed, range shifts for many species have been observed in response to increasing temperature over time (Chen et. al 2011). The observed pattern is upward, to higher latitudes and elevations as populations track their preferred environments (Fig. 2, Sexton et. al 2009; Ackerly 2003; Parmesan 2006). Past studies have observed a noticeable expansion of woody plants north past their original northern range limits in response to rising mean annual temperatures (Matías & Jump 2015).

Yet on their southern range edges, species are faced with possible extinction. In a study observing the distributional shifts of *Silene acaulis* (moss campion) and the geophyte *Polygonum viviparum* (alpine bistort) across six years, both populations saw die-out at their southern range edge (Doak & Morris 2010). The same effect has been seen earlier in history. As the temperature increased late in the last glacial interval, woody species populations established themselves at increasingly higher latitudes in so-called “migrations” (David & Shaw 2001).

To compensate for population loss at their southern edge due to the loss of their preferred environment, a species must shift northwards. Evidence shows that this shift may not occur quickly enough to account for the rate of changing temperatures and population loss. In a meta-analysis of plant distributions spanning 40 years, Bertrand et. al (2011) found a lag in the responses of herbaceous forest species to climate change. Since species often track their optimal environment (Ackerly 2003), strong limiting factors due to intra-specific competition may signal that a species will not be able to migrate at a rate sufficient to account for population loss at their southern range edge.

In addition to shifts in distribution, there is evidence that plant species underwent

local (genetic) adaptation in response to changing climates over the last 10,000 years (Davis & Shaw 2001). Yet populations are unlikely to undergo local adaptation at the pace necessary to match many current changes. Such changes include the invasion of competitors from adjacent positions and climate change (Ackerly 2003). Furthermore, studies show that competition hinders the advancement of colonists into new environments during shifts due to changing environments (Suttle, Thomsen & Power 2001; Urban et. al 2012), which limits a species' options in terms of biotic factors. Simultaneously, the speed of climate change affects how prevalent adaptive responses are (Ackerly 2003). Species are left with only one option: to migrate.

Work across species range extremes and range interiors such as the work that I have conducted allows us to see whether a species is primed to migrate. Additionally, such experiments have the potential to reveal how many limiting factors a species might face during migration, given its decreasing preferred climatic envelope. Because edge populations may see more variability in survival and reproduction due to the frequency at which they must respond to limiting factors (Reich & Oleksyn 2008; Sexton et. al 2011), range extremes are a unique opportunity to understand current modes of community assembly, as well as to predict species' future responses to climate change.

Latitudinal gradients as field laboratories

The study of range expansions and retractions is best accomplished through studies across latitudinal gradients. Using a latitudinal gradient as a study system allows for observations of a species across a larger spatial scale than in warming facilities, and provides a much more accurate representation of the habitat that either allows or prevents a species from establishing itself (De Frenne et. al 2013). Generally, there is a reliable poleward decrease in temperature across latitudes. On average, temperature decreases approaching the poles on the order of -0.73 C per degree latitude in the Northern hemisphere. Other climatic

variables, such as precipitation, also change across latitudes (De Frenne et. al 2013). Looking at species' traits and community composition across latitudes can provide insight into the effect of environmental variables on range limits and species' future success.

There is a dearth of research using latitudinal gradients, with many recent studies using altitudinal gradients as a predictor of species' responses to climate change (Walther 2003). This is likely due to the relative ease of collection relative to sampling across latitudes (Matías & Jump 2015). Altitudinal gradients have a much smaller rate of spatial change in temperature as compared to latitudinal gradients (Jump et. al 2009). This further increases the chances that local adaptation will successfully occur due to genetic drift (Davis & Shaw 2001).

Recording a species' functional traits and community composition across a latitudinal gradient provides a proxy for understanding how that species will respond to warming temperatures in its range exterior. Because the observed pattern of range shifts is up and poleward, studying a species across its latitudinal range provides important information about the effect of biotic and abiotic interactions that might decrease the possible rate of migration for a species. Latitudinal gradients are thus an apt study system with which to integrate species interactions and environmental stressors as predictors of a species' future distributions with regards to climate change.

Predictions and hypotheses

To determine the nature of woody species' future responses to climate change, I sampled six species of woody deciduous trees across four different sites on a latitudinal gradient in the northeastern United States and into Quebec. I predicted that:

H1: a species would become less competitive as they approached its latitudinal and climatic range limits, since populations at geographic range edges more frequently experience strong, limiting factors (Sexton et. al 2009).

A1: Alternatively, a species could grow more competitive as it approaches its range limit due to a mix of possible factors (e.g., the presence of fewer competitors, or increasing competitive ability of species at the abiotic environment of the edge etc.), thus leading to a greater realized niche.

A2: Finally, it is a possibility that competition is not predicted by position in its climatic or latitudinal range, indicating that local factors, such as habitat or resource availability, might matter more within each population. Similarly, other factors could determine competitiveness, such as community composition and local adaptation, rather than a species' position within its range limit. It is possible that local community composition around each tree is most predictive of its competitive success, with little overall pattern across the latitudinal gradient.

H2: I also predicted that trees at their range extremes would exhibit altered suites of functional traits from trees in the range interior (Weiher & Keddy 1998). Similar to the prediction (H1) above, because populations at their range limits often see more extreme environments, I predicted that their functional traits would differ from functional traits of individuals at their range interiors. Since species' functional traits that are linked to life strategies have been shown to vary based on the latitude of their origin (Cavender-Bares 2007), it seems likely that functional traits will exhibit variation across a species' range.

A1: Increases in intra-specific trait variation across a latitudinal gradient at the range limit could indicate increased genetic adaptation and/or increased plasticity. Since populations at their range extremes (and at more northern latitudes) face harsher environments than populations in their range interior, this could be due to resource partitioning in the presence of competitors in a sub-optimal environment, or simply because it is exhibiting greater plasticity in response to the more extreme environment.

A2: Alternatively, decreased intra-specific trait variation approaching a range limit would indicate that only one combination of traits is successful in that particular limiting environment (Burns & Straus 2012).

A3: Finally, no clear pattern in intra-specific variation in functional traits approaching the range limit of a species would indicate that other rules of community assembly are playing a larger part in the determining of range limits, such as competition or other biotic factors.

By examining community composition of certain woody species in conjunction with their corresponding functional traits across this latitudinal gradient, we can better predict how species composition will change at the forest level. In looking at both the community composition of species at these sites and their functional traits specifically at the range limits, we can understand which species are most at risk with the changing environment, and why they might not survive.

Here I report on a study to test these hypotheses (H1 and H2) by examining six woody species with individuals across their latitudinal ranges. I will evaluate (a) the effect of latitude on six woody species' competitiveness indices and functional traits and (b) the effect of distance from the environment at the center of a species' distribution (taken as its preferred environment) on the species' competitiveness indices and functional traits.

Methods

To understand community composition and functional traits in response to range limits, I examined the immediate vicinity of six deciduous tree species. I collected data on the community composition of the understory species, as well as current competitive environment for each focal tree. Data on functional traits for some of the same individuals were collected the year prior across all four sites through the summer.

Sites

I collected data at four sites, all spaced approximately one latitudinal degree apart from each other. Individuals of the six species had previously been tagged for past lab work. All sites were temperate deciduous forests with a small amount of human disturbance. The southernmost site was Harvard Forest, an LTER in Petersham, Massachusetts. The Harvard Forest spans approximately 3000 acres, and lies 110 kilometers west of Boston. The second site was located in the southern White Mountains in New Hampshire, off the Kancamagus Highway. As the southernmost part of the White Mountains, the site does not lie within a protected research area. However, as part of a national forest, it is fairly protected from human intrusion. The third site was in the Dartmouth College Second Grant, a site maintained by Dartmouth College for logging, recreational use, and scientific research. The Grant in northern New Hampshire, near the border of Maine. Finally, the northernmost site was located at the Station de biologie des Laurentides in Quebec. The Station de biologie des Laurentides is slightly larger than Harvard Forest, spanning 4000 acres, and is 75 kilometers northeast of Montreal. Due to the varying latitudes, the northern-most sites see a climate of colder winters and shorter growing seasons. All follow mild summers.

All four sites had previously been used by the Wolkovich lab for research in temporal ecology into the species used for this experiment, thus ensuring the presence of suitable

individuals at each site. Nearly all sampling sites occurred in closed canopy forests, except for some sampling plots of *Cornus alternifolia* and *Sorbus americana* which frequently appear on forest edges.

Species Selection

I sampled six different species of deciduous woody plants that are abundant across the latitudinal range of our sites in the Eastern US. The species' ranges vary, though all reach a range extreme near one of our four sites. The species in question are: *Acer pensylvanicum*, *Betula papyrifera*, *Cornus alternifolia*, *Fagus grandifolia*, *Hamamelis virginiana*, and *Sorbus americana*. Four of the six species reach their northern range extremes within or just outside of the latitudinal range covered by the four study sites. Two species, *Betula papyrifera* and *Sorbus americana*, reach their southern limits just below the southern-most site at Harvard Forest. *Hamamelis virginiana*, on the other hand, reaches its northern limit near our second southern-most site in the White Mountains.

Each species was selected based on (a) which species had been included in past studies at the four sites as part of the Wolkovich lab's work, and (b) the location of its range limits with relation to the four sites. I found each species' latitudinal range limits through the websites bonap.org, plants.usda.gov, and gbif.org. This was to account for records of tree species that were recorded outside their latitudinal limits (as evidenced by highly decreased abundance), but had still been recorded as appearing in different counties.

Competition and community data

For each of the six species, I tagged six (or found some previously tagged) individuals at each site that the individuals were present in. I compared total basal area of the focal individual with trees in the competitive environment around each focal individual. The direct

competitive environment was estimated as being within a 5-m radius around the focal tree. I recorded the DBH of all trees falling within this circle. I took note of the presence and absence of species in the understory, including species of saplings appearing under 1.3m. To determine competitiveness of our focal individuals, I used the percentage of total basal area attributable to the focal individuals as a proxy for competitive advantage in each plot.

Competition index was calculated according to methods outlined in Biging and Dobertin (1995). I calculated the sum of the trees larger than the focal tree, and took the ratio of the basal area of the focal tree to this sum. By using relative basal area, the trend of decreasing DBH as species increase in latitude would not show in the model.

Species composition in each site was recorded in both the overstory (in the form of recording the DBH of the surrounding trees over 1.3m tall) and the understory, including saplings. I used this data to generate a species presence/absence matrix for both the understory and overstory. This data was then used in the multi-dimensional scalar plots (Figs. 1 and 2). The multi-dimensional scalar plots were generated using the *vegan* package in *R*.

Climate data

Climatic centroid data was derived from data from the 30 arc-second WorldClim database (the highest resolution, available at <http://www.worldclim.org/>) across each species' recorded distributions. Once the data was downloaded, it was then averaged within an equal area grid of 50km^2 . The variables used to compute the climatic envelope in multi-dimensional space were average temperature (BIO1 Raster), mean diurnal range (BIO2 Raster), temperature seasonality (BIO4 Raster, akin to standard deviation*100), and annual precipitation (BIO12 Raster). This data was then used to calculate the climatic envelope by conducting a principal components analysis on the bioclimatic variables used. These two axes were then averaged for each species to find the coordinates of the climatic centroid. I then quantified the distance

from the climatic centroid and each of the focal individuals used in the study based on a climatic envelope specific to their species and their species' distribution. This was done through the formula for euclidean distance, putting in the coordinates of the climatic centroid and the coordinates of the location of each individual at its point in climatic space.

Functional Traits

For functional trait data, I used the data collected the previous summer by Harry Stone during the Harvard Forest REU Summer Program in 2015. The height of individuals not measured summer 2015 was measured through a clinometer. Additionally, I measured the Diameter at Breast Height (DBH) of any newly-tagged individuals (in cases where an individual is too close to another focal plant). Functional traits for newly-tagged individuals were not measured, but instead the traits measured in 2015 served as a representation of the traits of all species within that site.

Data on the functional traits Specific Leaf Area (SLA), Leaf N%, leaf dry matter content, and height were all collected for at least six individuals of the six species across the four sites (when applicable). Samples were processed in the John Torrey laboratory at the Harvard Forest. When possible, leaf samples were collected near the center of the tree canopy. Leaf size was measured using the LI-COR 3100 leaf area meter. To account for variation within an individual, a wide range of leaf sizes were measured to achieve a representative average size. Specific leaf area (SLA) was determined by the leaf size divided by dry mass of the leaves (Cornelissen 2003). The leaves were weighed after drying for than 24 hours at 60 OC.

Harry Stone and colleagues measured leaf dry matter content by comparing the fresh mass of the leaves, determined immediately after removal from the individual and the dried weight. Height of the individual was measured using a clinometer from 15 meters away from the tree. Stem specific density was measured by comparing the volume of the stem sample

to the dried mass. The volume of the stem was calculated by submersion in a container of water and the volume of the water displaced is equivalent to the volume of the stem. Then the stems were dried in excess of 24 hours in the drying oven at 60 OC.

Functional traits were analyzed using the convex analysis framework proposed in Cornwell et. al (2006). The convex hull is defined as the smallest convex shape enclosing a set of points. While most functional trait analyses take the mean of each trait for each species at a given site, using the convex hull method allows me to account for intraspecific variation in traits due to plasticity. This is particularly relevant given my hypothesis that the variation in traits will vary across a species' latitudinal distribution, indicating varying levels of effectiveness in different suites of traits for the plants' survival.

Convex hull volumes were calculated using the three traits specific leaf area (SLA), stem density, DBH, and ratio of percent carbon to percent nitrogen content. The convex hull volumes were calculated at both the species level for each site, as well as the community level for each site. For the species level at each site, I used the *R* package *geometry* to find the convex hull volume for all the species. For the community level, I used the *R* package *FD*. The package requires a presence/absence matrix for species composition to calculate the Functional Richness value (the same as the convex hull volume). I used the mean trait values for each species at each site and this presence and absence matrix to calculate the functional richness at each of the four sites. I then used this data to calculate the relative convex hull volume of my focal species to the community functional richness (or convex hull volume). This then quantifies the relationship between the focal species' convex hull volumes to the overall community's functional richness.

Statistical analyses

I used both linear mixed effects models and simple linear models to evaluate the relationship between competition and latitude for each species at each sampling location. Using the simple

linear models allowed me to quantify the basic relationship between distance from minimum latitude and the competition index for each species (Fig. 3). The linear mixed effects model allows me to quantify the general behavior of a species' competition index in relation to its distance from its minimum latitude for all species across their latitudinal distributions. I did the same for quantifying the relationship between the distance from the climatic centroid and the relative basal area (Fig. 4). I also used a simple linear model to quantify the correlation between the relative convex hull trait volume for each focal species and the sites' latitude, as well as for the species' relative convex hull trait volume in relation to their relative basal area (Figs. 5 and 6, respectively).

Results

The species composition across the four sites, as evidenced by the multi-dimensional scalar plots (Fig. 1), shows a marked difference between the species compositions of Harvard Forest and the other three sites. This is mirrored in both the overstory and understory species composition, suggesting that the type of competition seen by the deciduous trees present in Harvard Forest may differ from that of the other three sites. The multi-dimensional scalar plots show the differences in combinations of species, meaning that the three overlapping convex hulls (representing the White Mountains, the Grant site, and St. Hippolyte in increasing latitudes) saw very similar combinations of species in both the overstory and the understory, whereas Harvard Forest saw very different combinations of species from the other three sites. The plots are dimensionless, and serve to represent the convex hull of unique combinations of species. For example, a larger polygon would indicate more unique combinations of species present (such as in the White Mountains site), whereas the Grant plot sees a smaller diversity of species combinations. The understory in general was composed of smaller diversity of species combinations as compared to the overstory.

The strength of competition index - latitude relationships differed by species, with no clear pattern across all six species. There was little to no correlation between increasing latitude (approaching northern edges) and the competitive index of the focal trees. *Acer pensylvanicum* (denoted by ACEPEN in the figures) showed a slight negative correlation ($R^2 = -0.011$) between the relative basal area of the focal individuals to the distance from the minimum latitude. This indicates a general trend of *Acer pensylvanicum* individuals growing less competitive as they grow more northward (closer to the species' northern range limit). *Betula papyrifera* (denoted by BETPAP in the figures), whose distribution lies predominantly north of Harvard Forest (with the center near the northern-most testing site), shows a slight negative correlation ($R^2 = -0.033$) between its relative basal area and its distance from its minimum latitude. Similarly, *Cornus alternifolia* (denoted by CORALT) showed the same

slight (though statistically insignificant) trend downwards ($R^2 = -0.068$), as did *Hamamelis virginiana* (HAMVIR, $R^2 = -2.049$). In other words, *A. pensylvanicum*, *B. papyrifera*, *C. alternifolia*, and *H. virginiana* ($R^2 = -2.049$) all show a decreasing level of competition (in the form of relative basal area to surrounding individuals larger than they are) as they go north in latitude.

In terms of species distributions playing a part in the competitiveness of a species at a location, this would indicate there is no significant correlation between a species' place within its range and its competitiveness. *B. papyrifera* approaches its range interior (its distribution lies mostly north of the Saint Hippolyte experimental site). Thus, the data suggests that *B. papyrifera* grows less competitive as it approaches its range interior. Similarly, *Fagus grandifolia* (FAGGRA) shows a trend towards increasing competitiveness as it approaches its range extreme. Both trends are inconsistent with my hypothesis that a species would decrease in competitiveness towards its range limit, and increase towards its range interior.

Like *B. papyrifera*, *Sorbus americana* (SORAME) also approaches its range interior across distance from the minimum latitude represented in the data. Yet unlike *B. papyrifera*, *Sorbus americana* shows a positive correlation between its position in its latitudinal range and its relative basal area - and thus its competition index ($R^2 = 0.250$). In other words, as *S. americana* approaches its range interior, the more competitive it becomes relative to trees around it. This is consistent with my hypothesis that a species would grow less competitive as it approached its range extreme due to constrained resources and niche limitations. This is mirrored as well in the results of the linear mixed effects model (Table 2), which shows a coefficient of fixed general effects to be -0.0317. Thus, the general trend is for species to decrease in competitiveness as they go north, taking into account the random effects.

In contrast to the biotic interactions that might be limiting a species at its range limit, measuring the euclidean distance from a species' climatic centroid in relation to its relative basal area shows how limiting the climatic factors are to a species. Since the main discussion

over what is limiting a species at its range limits often emphasizes climatic variables or biotic factors, it follows that a discussion of competitiveness should include its relationship with a plant's distance from its climatic centroid. Plotting this relationship reveals an inconsistent pattern across all six species (Fig. 3). *A. pensylvanicum* ($R^2 = -0.050$), *B. papyrifera* ($R^2 = -0.033$), and *F. grandifolia* ($R^2 = -0.049$) all show very little correlation between an individual's euclidean distance from its climatic centroid and its relative basal area to the surrounding individuals. *C. alternifolia*, on the other hand, shows a positive correlation between its distance from the climatic centroid and its relative basal area. *It is worth noting that CORALT is generally quite a small tree, and thus is usually the least competitive within a given plot* (include this?). This correlation indicates that as *C. alternifolia* individuals grow farther from their climatic centroid, the more competitive they become. This is contrary to my hypothesis that the farther an individual is from its climatic centroid (theorized as the place it grows best), the less basal area it occupies relative to the individuals in the surrounding area.

Unlike *C. alternifolia*, *H. virginiana* and *S. americana* show a negative correlation between the distance from their climatic centroid and the focal individuals' relative basal areas ($R^2 = 0.047$ and $R^2 = 0.187$ with a p value of 0.047 respectively). This indicates that for *H. virginiana* and *S. americana*, climate is correlated with limitations in its competitiveness. Although there could be other factors present in limiting the competitiveness of *S. americana*, this suggests that distance from climatic centroid could be a major factor in limiting the spread of the trees.

Figures




Figure 1: Map of all four sites mapped onto a raster image of their mean annual temperatures, compiled from the 30 arcsecond layer from <http://worldclim.org>.




Figure 2: Anthropogenic warming can drastically alter the environment to which a species has adapted. This shift in climate also means that a species' ideal "climate envelope" shifts. If a species is limited by its preferred environmental variables, it may have to shift its range in order to compensate for the change in climate. It must match its realized niche with the shift climate envelope or face extinction at its southern (warming) edge.

Range Maps




Figure 2: Representation of the range of *A. pensylvanicum*. Individuals were found at all sites, with functional trait data taken across all four. Used as one of four species approaching its northern range at the northernmost site at Saint Hippolyte. Image taken from <https://gec.cr.usgs.gov>.

Betula papyrifera




Figure 3: Representation of the range of *B. papyrifera*. Individuals were found at all sites, with functional trait data taken across all four. Used as one of two species entering its range interior away from Harvard Forest (taken as near its southern range extreme). Image taken from <https://gec.cr.usgs.gov>.




Figure 4: Representation of the range of *C. alternifolia*. Community composition and competition index data was collected at the White Mountains, Dartmouth Grant, and Saint Hippolyte sites. Functional trait data was taken at the Harvard Forest, Dartmouth Grant, and Saint Hippolyte sites (over the year between the trait data and community composition data collections, the individuals found at Harvard Forest became very unhealthy). Used as one of four species approaching its northern range at the northernmost site at Saint Hippolyte. Image taken from <https://gec.cr.usgs.gov>.

Fagus grandifolia




Figure 5: Representation of the range of *F. grandifolia*. Individuals were found at all four sites, with functional trait data taken across all four. Used as one of four species approaching its northern range at the northernmost site at Saint Hippolyte. Image taken from <https://gec.cr.usgs.gov>.




Figure 6: Representation of the range of *H. virginiana*. Individuals were found at only two sites, with functional trait data taken at both (though not enough to compute convex hull volumes). Used as one of four species approaching its northern range at the one of the middle sites in the White Mountains. Image taken from <https://gec.cr.usgs.gov>.




Figure 7: Representation of the range of *S. americana*. Individuals were found at all four sites, though not enough functional trait data was collected to determine convex hull volumes across all four sites (even though data was collected across all four). Used as one of two species that nears its southern range edge at the site near Harvard Forest. Image taken from <https://gec.cr.usgs.gov>.




Figure 8: A multidimensional scalar plot representing the species composition and variation of the overstory across all four sites. The shape of the polygon represents the different combinations of species that appear in the overstory in each plot within each site. The larger the polygon, the more unique combinations of species that appear together at that site. The overlap across the northern three sites (White Mountains, Dartmouth Grant, and Saint Hippolyte) signifies a great deal of similarity in combinations of species (species that appeared together in plots at those sites). There is a clear difference between Harvard Forest (LTER site), and the other three of the sites.




Figure 9: A multidimensional scalar plot representing the species composition and variation of the understory (under 1.3m and thus not measured for DBH) across all four sites. The shape of the polygon represents the different combinations of species that appear in the overstory in each plot within each site. The larger the polygon, the more unique combinations of species that appear together at that site. The overlap across the northern three sites (White Mountains, Dartmouth Grant, and Saint Hippolyte) signifies a great deal of similarity in combinations of species (species that appeared together in plots at those sites). There is a clear difference between Harvard Forest (LTER site), and the other three of the sites. Between the overstory and understory species composition, however, there are fewer unique combinations of species that occur in the understory. This is denoted by the slightly smaller polygons.





Figure 10: Competitiveness index across latitudes varies by species. Overall, there is a positive trend in competition index as individuals appear farther at sites located farther north. *B. papyrifera* and *S. americana* are separated from the other four species to emphasize that they are the two species that are moving away from their (southern) range extremes, rather than approaching their northern range extremes.






Figure 12: The suites of traits examined (SLA, %C:%N, DBH, Stem density) did not vary predictably with latitude. Across the four species that had enough trait data to calculate convex hull volume, two species (*B. papyrifera* and *A. pensylvanicum*) exhibited a decrease in their relative convex hull volume across latitudes. The other two increased across latitudes. All species appeared at all four sites, except for *C. alternifolia*, which appeared at three.




Figure 13: Relative convex hull volume decreased consistently across all four species examined as they moved away from their climatic centroids. Three of the species appeared at the four sites, and one appeared at three sites.




Figure 14: Across the four species with sufficient trait data, relative convex hull volume of species across all sites did not reliably predict the relative basal area. This suggests more complex factors at play in determining a species' suite of functional traits than an individual's competitiveness.




Figure 15: Convex hull volumes using a PCA across all species show marked variations in functional traits across sites.

Discussion

Ecologists have long understood that range limits are caused by complex dynamics between biotic and abiotic factors. These dynamics are especially important to understand in light of climate change. A warming world and shift in abiotic factors necessitates research into the limiting effect of both abiotic and biotic interactions at range limits. As a species' climatic envelope migrates north, looking at the limiting power of biotic and abiotic factors can provide insight into the future survival of that species.

My results suggest that the effect of biotic and abiotic factors on limiting trees at their range limits varies across tree species. They showed no clear pattern with regards to a species' competitiveness across latitudes or climatic envelopes, even within species positioned closer to their southern range limits.

There was a similar dearth of a clear pattern in examining the functional traits of four of the six focal species across the sites at which they were found. The functional trait space each species occupied relative to the average functional diversity of communities at each site reflected very little effect derived from latitudinal or climatic gradients.

Competition across latitudinal and climatic range limits

I found no strong support for my hypothesis that species would decrease in competitiveness as they approached their latitudinal and/or climatic range limits. The overall trend in competitiveness across latitudes was positive, indicating a slight increase in dominance as a species goes north. This lack of clear pattern probably indicates that latitudinal range limits are caused by a complex interplay of biotic and abiotic factors that is not fully accounted for in simply examining a species' competitiveness. This suggests that competition is not predicted by its position in its latitudinal range (**H1, A2**).

The overall lack of a significant relationship competitiveness and position in range, combined

with different relationships for certain species, indicate that there are other factors playing a part in determining a species' range limits. These other factors probably prevent a clear pattern from emerging in my data. It is possible that the species of trees, such as *H. virginiana*, *A. pensylvanicum*, and *S. americana*, that are mid-sized and never reach the same level of competitiveness in basal area as *B. papyrifera*, for example, are more susceptible to biotic factors as they near their latitudinal range limits. This could be due to biotic factors not examined, such as percentage cover in the canopy, which are widely recognized as constraining plant development over time due to possible loss of energy from sunlight (Biring & Dobbertin 1995, Contreras et. al 2011). To account for this, one could utilize a different competition index in a similar study on latitudinal and climatic ranges. More accurate competition indices are distance-dependent (whereas summing up the surrounding basal area of larger competitors than the focal individual is a distance-independent index). The best distance-dependent competition index is the sum of the horizontal angles multiplied by the ratios of the DBHs of the neighbor and the cored individuals (outlined in Rouvinen & Kuuluvainenas 1997 and as tested in Contreras et. al 2011). The downside of this method (and of distance-dependent competition indices in general) is that they require much more time to collect data. This is a major limiting factor in latitudinal studies, since so many resources are required to collect data in the field across such a large area. Additionally, studies have observed that the stronger limiting factors at a species' southern range limit are interspecific competition, while environmental factors limit species at their northern limits (Brown et. al 1996, Nakawatase & Peterson 2006). Due to the limited data I recorded of species with southern range limits close to the four sites, my study was lacking in this respect.

It is possible that community composition and local adaptation matter more. In this case, local community composition could be more predictive of a tree's success. This is a positive outlook in terms of the possibility of a tree species' potential to migrate north to account for population loss on its southern range edge, though this is not supported by current literature that suggests range shifts will probably lag the velocity of climate change (Loarie et. al 2009).

It is also possible that a species is more limited by competition at its southern edge, and thus my results would not reflect this as they are heavily reliant on species that lie closer to their northern range limits across our sites. Biotic interactions have been theorized as the stronger limiting factor at southern range edges. Additionally, ecological models predict that biotic interactions will create lags in climate tracking (Svenning et. al 2014). This has also been supported by experiments on the ground (Fisichelli et. al 2012).

I saw a similarly unclear pattern in competitiveness across species' climatic envelopes. The only species to exhibit a decrease in competitiveness was *B. papyrifera*, suggesting that environment may be more of a limiting factor in its establishment than the other species examined. Overall, there was a negative trend in a species' relative basal area in relation to the distance of the individual from its climatic "sweet spot." This confirms my hypothesis that a species will become less competitive as it approaches its climatic range limit (or moves away from the center of its climatic range, where environmental factors are most favorable for it). This suggests the species examined have climatic ranges that are not necessarily representative of their fundamental niche, and also that the latitudinal range limit as a proxy for harsh environmental conditions may not tell the whole story.

This could mean that climate is not a limiting factor for the species that do not show a negative correlation. Instead, it is possible that they do not need to track their habitat and thus will not see a range shift in the face of a warming climate. A positive correlation between habitat preference and performance in the preferred habitat promotes specialization to the preferred habitat (Holt 1987). Seeing habitat preference have little effect when examined in relation to performance suggests that these species have not specialized to their preferred habitats, and so may not see the population loss that some species forced to shift in their range due to climate changes have faced in the past (Davis et. al 2001).

There is evidence that plant distribution data does not match climatic tolerance estimates (Curtis and Bradley 2016). Additionally, climatic niche shifts are not reflected in species

distribution, and many species in naturalized distribution were found outside native climatic conditions (Early and Sax 2014). If a species is not at climatic equilibrium, its competitiveness to individuals around it could reflect the effect of climate from earlier in the individual's life. This could explain the absence of a strong effect of climate on a species' size (and competitiveness), and indicates climate could be much less of a limiting factor than initially thought. Climate has also been shown to be a strong limiting factor on tree species at their range extremes, depending on their growth stage (Ettinger and HillesRisLambers 2013).

Suites of functional traits across ranges

I hypothesized that trees at their range extremes would show altered suites of functional traits from trees in their range interior. To examine this, I examined how much of a community's functional richness was occupied by each focal species. Across sites, latitude appeared to have no discernible effect overall on the species' relative functional richness (Fig. 5). This could indicate that biotic factors or environmental factors not consistent across latitudes are more predictive in a species' suite of functional traits. Past studies have found that latitude does predict trends of functional traits (De Frenne et. al 2011), but the traits examined differ from those examined in this study.

Conversely, across a climatic range there is overall a decrease in species' relative functional richness. This suggests that environment, more than latitude, is a determining factor in the functional traits exhibited by a species. As a species moves farther away from its climatic "sweet spot," its functional traits become more similar and show less variability relative to the functional richness of its community. In a study conducted by Guittar et. al (2016), only certain functional traits showed a reliable pattern across climatic gradients (including SLA, as examined in this study), but fewer traits were consistently predictive of changes in species abundances in warmed turf used in the study.

This result confirms my hypothesis that trees moving away from their climatic range interior

exhibit altered suites of functional traits from trees in their range interior, and specifically *A1*. This suggests that such suites of traits are successful in that particular limiting environment (Burns and Straus 2012). However, functional trait similarity was not linked to a tree's relative size (i.e. competitiveness index) within a plot. This could suggest that abiotic factors play more of a part in the phenotypes observed in a species, though this is not supported by the literature (Burns and Straus 2012; Kunstler et. al 2015). In a study by Funk and Wolf (2016), certain suites of functional traits in grasses were a better predictor of competitive outcome than phylogenetic relatedness.

There was no discernible pattern across species in the changes of their suites of functional traits, based solely on the overlap of their convex hulls (Fig. AHL Add figure number here with convex hull pictures). Two species, *C. alternifolia* and *F. grandifolia*, exhibited a marked difference in the convex hulls of their traits between the Harvard Forest and the other three of the sites. This mirrors the difference in species composition between Harvard Forest and the other three sites observed in both the understory and overstory (Fig. 3). Such results suggest that the type of species that co-occur with the focal species could affect the suites of traits exhibited in an individual.

Areas of future research

In predicting the future distributions of species in light of climate change, much is still unclear about the effect of biotic and abiotic variables. Based on my results, the relative strength of the two factors in limiting ranges seems to vary across species. The species *B. papyrifera* tends to grow much larger than any of the other species examined (and was one of two trees selected for the focal species that neared its southern edge rather than its northern edge). It seems likely that a major factor in *B. papyrifera*'s consistently larger relative basal area is the size and age of its mature trees, which will likely not be indicative of a species at climatic equilibrium, and thus will not provide an accurate forecast of species distributions due to

climate change.

Yackulic et. al (2015) outlines this problem, and suggests that models and field work take into account colonization and extinction rates of species, thus looking at time-variant relationships between environmental conditions and changes in species distribution. With trees especially, it will be important to look at dispersal and seedling establishment within this, as plants in general are more likely to be limited by the extent of their dispersal when looking ahead at possible migrations due to climate change (Ackerly 2003). Future research into competitiveness across ranges should take this into account, as looking at current competitiveness cannot provide the full picture of the complex dynamics at a species' range edge.

Another factor that is crucial to an understanding of the relationship between a species, its competitors, and its environment is its growth stage. Indeed, growth stage has been shown to make a marked difference in how a species responds to a warming environment and competitive interactions (Zhu et. al 2012; Ettinger and HillesRisLambers 2013). In modeling future species distributions, taking into account growth stages for more long-lived species such as the deciduous trees examined in my study will provide a much clearer and more accurate idea of species' responses to climate change.

Additionally, my results emphasize the importance of examining species at their southern ranges in future work. Unfortunately, I was only able to examine two species that neared their southern range edges (*S. americana* and *B. papyrifera*). Studies have shown that interspecific competition sets lower limits (Brown et. al 1996), necessitating the inclusion of southern range limits as a way to understand the complex dynamics of biotic factors and future success of species.

Conclusion here — do I make a whole new section for this?

Note: these are the stipulations for the Discussion — The discussion section should first of all summarize and analyze the results that you have presented. Also recapitulate your overall argument(s) presented in the Introduction as well as strengths and weaknesses, and address

the theoretical issues that were used in approaching and analyzing the problem. You should also explain how you may have modified your view of the issues in the course of conducting the analysis. If your experiments did not produce significant or meaningful results, explain here your thoughts on why this might have occurred, and suggest how things could be done differently. Convince the reader that you have fully thought through and understand the implications of your work, no matter how individual experiments may have turned out.

Question: where should I integrate how things could be done differently?

Reference

- Ackerly, David D. "Community Assembly, Niche Conservatism, and Adaptive Evolution in Changing Environments." *International journal of plant sciences* 164 (2003): S165-84. Print.
- Bertrand, Romain, et al. "Changes in Plant Community Composition Lag Behind Climate Warming in Lowland Forests." *Nature* 479.7374 (2011): 517. Print.
- Biging, G. S., and M. Dobbertin. "Evaluation of Competition Indices in Individual Tree Growth Models." *Forest Science; Evaluation of competition indices in individual tree growth models.* 2 (1995): 360-77. Print.
- Brown, James H., George C. Stevens, and Dawn M. Kaufman. "The Geographic Range: Size, Shape, Boundaries, and Internal Structure." *Annual Review of Ecology and Systematics* 27 (1996): 597-623. Print.
- Burns, Jean H., and Sharon Y. Strauss. "Effects of Competition on Phylogenetic Signal and Phenotypic Plasticity in Plant Functional Traits." *Ecology* 93 (2012): S126-37. Print.
- Cavender-Bares, J. 2007. Chilling and freezing stress in live oaks (*Quercus* section *Virentes*): intra- and inter-specific variation in PSII sensitivity corresponds to latitude of origin. *Photosynthesis Research* 94. (2007): 437-53. Print.
- Chen, I. C., et al. "Rapid Range Shifts of Species Associated with High Levels of Climate Warming." *Science* 333.6045 (2011): 1024. Print.
- Contreras, Marco A., David Affleck, and Woodam Chung. "Evaluating Tree Competition Indices as Predictors of Basal Area Increment in Western Montana Forests." *Forest Ecology and Management* 262.11 (2011): 1939-49. Print.
- Cornelissen, J. H. C., et al. "A handbook of protocols for standardised and easy measurement of plant functional traits worldwide." *Australian Journal of Botany* 51.4 (2003): 335-380.
- Cornwell, William K., and David D. Ackerly. "Community Assembly and Shifts in Plant Trait Distributions Across an Environmental Gradient in Coastal California." *Ecological Monographs* 79.1 (2009): 109-26. Print.
- Darwin, Charles. *On the Origin of Species*. Ed. Jim Endersby. Cambridge ; New York:, 2009. Print.
- Davis, M. B., and R. G. Shaw. Range Shifts and Adaptive Responses to Quaternary Climate Change. 292 Vol, 2001. Print.
- De Frenne, Pieter, et al. "Latitudinal Gradients as Natural Laboratories to Infer Species' Responses to Temperature." *The Journal of Ecology* 101.3 (2013): 784. Print.
- Doak, Daniel F., and William F. Morris. "Demographic Compensation and Tipping Points

- in Climate- Induced Range Shifts.” *Nature* 467.7318 (2010): 959. Print.
- Funes, Guillermo, et al. “Seed Size and Shape are Good Predictors of Seed Persistence in Soil in Temperate Mountain Grasslands of Argentina.” *Seed Science Research; Seed Sci.Res.* 9.4 (1999): 341-5. Print.
- Grime, J. P. “Vegetation Classification by Reference to Strategies.” *Nature* 250.5461 (1974): 26. Print.
- Hargreaves, Anna L., Karen E. Samis, and Christopher G. Eckert. “Are Species’ Range Limits Simply Niche Limits Writ Large? A Review of Transplant Experiments Beyond the Range.” *The American Naturalist* 183.2 (2014): 157. Print.
- Holt, Robert. “Population Dynamics and Evolutionary Processes: The Manifold Roles of Habitat Selection.” *Evolutionary Ecology* 1.4 (1987): 331-47. Print.
- Jump, Alistair S., Csaba Mátyás, and Josep Peñuelas. “The Altitude-for- Latitude Disparity in the Range Retractions of Woody Species.” *Trends in Ecology & Evolution* 24.12 (2009): 694-701. Print.
- Jung, Vincent, et al. “Intraspecific Variability and Trait-Based Community Assembly.” *Journal of Ecology* 98.5 (2010): 1134-40. Print.
- Kraft, Nathan J. B., Renato Valencia, and David D. Ackerly. “Functional Traits and Niche-Based Tree Community Assembly in an Amazonian Forest.” *Science (New York, N.Y.)* 322.5901 (2008): 580. Print.
- Louhan, Am, Df Doak, and A. L. Angert. Where and when do Species Interactions Set Range Limits? 30 Vol., 2015. Print.
- MacArthur, Robert, and Richard Levins. “The Limiting Similarity, Convergence, and Divergence of Coexisting Species.” *The American Naturalist* 101.921 (1967): 377-85. Print.
- Matías, Luis, and Alistair S. Jump. “Asymmetric Changes of Growth and Reproductive Investment Herald Altitudinal and Latitudinal Range Shifts of Two Woody Species.” *Global Change Biology* 21.2 (2015): 882-96. Print.
- Mcgill, Brian J., et al. “Rebuilding Community Ecology from Functional Traits.” *Trends in Ecology & Evolution* 21.4 (2006): 178-85. Print.
- McKenney, Daniel W., et al. “Potential Impacts of Climate Change on the Distribution of North American Trees.” *Bioscience* 57.11 (2007): 939-48. Print
- Nakawatase, Jill, and David Peterson. “Spatial Variability in Forest Growth - Climate Relationships in the Olympic Mountains, Washington.” *Canadian Journal of Forest Research* 36.1 (2006): 77-91. Print.
- Nussey, D. H., et al. “Selection on Heritable Phenotypic Plasticity in a Wild Bird Population.” *Science Magazine* 310.5746 (2005): 304-6. Print.

Parmesan, Camille. "Ecological and Evolutionary Responses to Recent Climate Change." *Annual Review of Ecology, Evolution, and Systematics* 37 (2006): 637. Print.

Peterson, A. T., J. Soberón, and V. Sánchez-Cordero. "Conservatism of Ecological Niches in Evolutionary Time." *Science* 285.5431 (1999): 1265-7. Print.

Pigot, Alex L., and Joseph A. Tobias. "Species Interactions Constrain Geographic Range Expansion Over Evolutionary Time." *Ecology Letters* 16.3 (2013): 330-8. Print.

ADD REICH AND OLEKSYN 2008

Rouvinen, S., and T. Kuuluvainen. "Structure and Asymmetry of Tree Crowns in Relation to Local Competition in a Natural Mature Scots Pine Forest." *Canadian Journal of Forest Research* 27.6 (1997): 890-902. Print.

Sexton, Jason, et al. "Evolution and Ecology of Species Range Limits." *Annual Review of Ecology, Evolution, and Systematics* 40 (2009): 415. Print.

Skelly, David K., et al. "Evolutionary Responses to Climate Change." *Conservation Biology* 21.5 (2007): 1353-5. Print.

Stahl, Ulrike, Björn Reu, and Christian Wirth. "Predicting Species' Range Limits from Functional Traits for the Tree Flora of North America." *Proceedings of the National Academy of Sciences of the United States of America* 111.38 (2014): 13739. Print.

Suttle, K. B., Meredith A. Thomsen, and Mary E. Power. "Species Interactions Reverse Grassland Responses to Changing Climate." *Science* (New York, N.Y.) 315.5812 (2007): 640. Print.

Thomas, Chris D., Aldina M. A. Franco, and Jane K. Hill. "Range Retractions and Extinction in the Face of Climate Warming." *Trends in Ecology & Evolution* 21.8 (2006): 415-6. Print.

Thompson, K., S. R. Band, and J. G. Hodgson. "Seed Size and Shape Predict Persistence in Soil." *Functional Ecology* 7.2 (1993): 236-41. Print.

Urban, Mark C., Josh J. Tewksbury, and Kimberly S. Sheldon. "On a Collision Course: Competition and Dispersal Differences Create no-Analogue Communities and Cause Extinctions during Climate Change." *Proceedings of the Royal Society B* 279.1735 (2012): 2072-80. Print.

Weiher, Evan, and Paul A. Keddy. "Community Assembly Rules, Morphological Dispersion, and the Coexistence of Plant Species." *Oikos* 81.2 (1998): 309-22. Print.

Zhu, Kai, Christopher W. Woodall, and James S. Clark. "Failure to Migrate: Lack of Tree Range Expansion in Response to Climate Change." *Global Change Biology* 18.3 (2012): 1042-52. Print.