Draft version October 13, 2026
Typeset using LaTeX **twocolumn** style in AASTeX7

arXiv:submit/8220485 [astro-ph.HE] 13 Oct 2026

Making Reconnection Cooler: A relativistic resistive magnetohydrodynamics model for radiative collisionless reconnection

Alicia M. Savelli [1,2,3] Daniel Grošelj [4] Bart Ripperda [1,2,5,6] Amir Levinson [7] Alexander Philippov [8,9,10,11] and Norman Murray [1,2,5]

[1] *Canadian Institute for Theoretical Astrophysics, University of Toronto, 60 St. George Street, Toronto, ON M5S 3H8, Canada*
[2] *David A. Dunlap Department of Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada*
[3] *Dunlap Institute for Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada*
[4] *Centre for mathematical Plasma Astrophysics, Department of Mathematics, KU Leuven, Celestijnenlaan 200B, B-3001 Leuven, Belgium*
[5] *Department of Physics, University of Toronto, 60 St. George St, Toronto, ON M5S 1A7, Canada*
[6] *Perimeter Institute for Theoretical Physics, 31 Caroline St. North, Waterloo, ON N2L 2Y5, Canada*
[7] *School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel*
[8] *Department of Physics, University of Maryland, College Park, MD 20742, USA*
[9] *Joint Space-Science Institute, University of Maryland, College Park, MD 20742, USA*
[10] *Physics Department, Stanford University, 382 Via Pueblo Mall, Stanford, CA 94305, USA*
[11] *Kavli Institute for Particle Astrophysics and Cosmology, Department of Physics, Stanford University, Stanford, CA 94305, USA*

ABSTRACT

Relativistic magnetic reconnection around astrophysical compact objects is a likely site for particle energization and multiwavelength emission. Kinetic theory has substantially modelled such collisionless radiative relativistic reconnection using a radiative drag force countering the acceleration of the particles. Here, we systematically derive an energy-momentum sink term in relativistic resistive magnetohydrodynamics (RRMHD), informed by kinetic theory and coupled with a charge-starved resistivity prescription which reproduces the fast reconnection rate seen in kinetic calculations, to model radiative losses. We show that our radiative loss term accurately captures important macroscopic properties – including the temperature in the reconnection layer and the bulk motion energy distribution of the out-flowing plasma – for inverse Compton cooling applied to two-dimensional RRMHD simulations of a reconnection layer and benchmarked to kinetic simulations. Our results hold in various cooling regimes with a one-to-one mapping to kinetic theory. Our radiative model lays a foundation for future RRMHD simulations of plasma in the relativistic radiative regime, such as in reconnection sites in a local turbulent flow or globally in the magnetospheres of black holes and neutron stars.

Keywords: Plasma astrophysics (1261) — High energy astrophysics (739) — Magnetohydrodynamics (1964) — Magnetohydrodynamical simulations (1966) — Compact radiation sources (289) — Special relativity (1551)

1. INTRODUCTION

Magnetic reconnection is a canonical dissipation channel in astrophysical sources. One of the longest- and best-studied examples is the solar corona, where reconnection drives flares and contributes to the heating of the surrounding plasma (e.g., P. Antolin et al. 2021; X. Cheng et al. 2023; R. Patel et al. 2025). More recently, interest has expanded towards more distant and extreme sources, such as accreting black holes (e.g., K. Parfrey et al. 2019; L. Sironi & A. M. Beloborodov 2020; N. Sridhar et al. 2021; J. M. Mehlhaff et al. 2021; B. Ripperda et al. 2022; H. Hakobyan et al. 2023b; A. Galishnikova et al. 2023; S. Gupta et al. 2024; H. Hakobyan et al. 2025; N. Sridhar et al. 2025; J. M. Mehlhaff et al. 2026), neutron-star magnetospheres and their winds (e.g., B. Cerutti & A. A. Philippov 2017; A. Philippov et al. 2019; B. Cerutti et al. 2020; H. Hakobyan et al. 2023a; J. F. Mahlmann 2024), astrophysical jets (e.g., A. Levinson 2006; M. Petropoulou et al. 2016; L. Sironi et al. 2021), and gamma ray bursts (e.g., Y. Lyubarsky 2010; J. C. McKinney & D. A. Uzdensky 2012; J. Granot 2016). Reconnection in high-energy environments com-

Corresponding author: Alicia M. Savelli
Email: alicia.savelli@utoronto.ca

monly occurs in the relativistic regime, which differs from more conventional (e.g., solar) plasmas, such that the magnetic energy density upstream of the reconnection layer far exceeds the plasma rest mass energy density (i.e., a high upstream magnetization parameter $\sigma_0 \equiv B_0^2/4\pi\rho_0 c^2 \gg 1$, where B_0 is the strength of the upstream magnetic field, ρ_0 is the upstream mass density, and c is the speed of light), leading to efficient nonthermal particle acceleration (L. Sironi & A. Spitkovsky 2014; F. Guo et al. 2014; G. R. Werner et al. 2016).

Reconnection in high-energy sources is also frequently accompanied by rapid radiative cooling, which competes against the particle acceleration. This cooling introduces new effects that alter the physics of the reconnection layer and its emission signature. Such effects include cooling of the particles on timescales faster than the dynamic dissipation time, radiative drag on the bulk motions, and – in the most extreme cases – reprocessing of the plasma composition via electron-positron pair creation and annihilation (H. Hakobyan et al. 2019; K. M. Schoeffler et al. 2023; J. Mehlhaff et al. 2024). Therefore, in order to predict emission from high-energy sources powered by reconnection, there is a need to include, at some level, both radiative effects as well as relativistic kinetic physics.

One notable example of such a high-energy source undergoing rapid cooling is the magnetospheres of neutron stars, where reconnection can power multiwavelength emission (A. A. Philippov & A. Spitkovsky 2018; A. Philippov et al. 2019; Y. Lyubarsky 2020; J. F. Mahlmann et al. 2022; H. Hakobyan et al. 2023a; J. Parsons et al. 2026). Another prominent candidate for relativistic radiative reconnection is the black hole "corona" – hot plasma of moderate optical depth (F. Yuan & A. A. Zdziarski 2004; A. C. Fabian et al. 2015) near the event horizon of luminous accreting black holes. Here, radiative reconnection has been proposed as a mechanism for plasma energization (A. M. Beloborodov 2017; L. Sironi & A. M. Beloborodov 2020; N. Sridhar et al. 2021), powering hard X-ray emission (A. A. Zdziarski & M. Gierliński 2004) believed to be sourced from the Comptonization of soft seed photons by relativistically hot electrons of temperature $k_{\rm B}T_{\rm e} \sim 100$keV (D. M. Eardley et al. 1975; S. L. Shapiro et al. 1976), where $k_{\rm B}$ is the Boltzmann constant, as well as by the bulk plasma flow (A. M. Beloborodov 2017; N. Sridhar et al. 2025).

Reconnection in astrophysical plasmas also typically occurs in the collisionless regime (i.e., the mean free path of a particle is much larger than the system size). Kinetic particle-in-cell (PIC) simulations are required to accurately model the collisionless microsphysics of the plasma, but these simulations are computationally limited to a local reconnection layer and are able to capture neither their formation nor their interaction with the accretion flow of a global astrophysical system. This separation of scales is especially relevant in the case of black hole coronae, where there has been considerable effort towards understanding the plasma-radiation interaction from first principles with the corona structured as both a single large-scale reconnection layer (L. Sironi & A. M. Beloborodov 2020; N. Sridhar et al. 2021; J. M. Mehlhaff et al. 2021) and as a turbulent plasma driving multiple smaller current sheets (D. Grošelj et al. 2024; J. Nättilä 2024; D. Grošelj et al. 2026). Recent work has also shown large-scale reconnecting current sheets forming within the putative coronal region in global general relativistic magnetohydrodynamic (GRMHD) simulations of an accreting black hole (M. T. P. Liska et al. 2022; N. Sridhar et al. 2025); however, the GRMHD framework lacks essential microphysical components of collisionless reconnection, such as a fast reconnection rate or physically-informed models for coronal plasma-radiation interaction and cooling, which could significantly impact global accretion. There exist subgrid models for thermal IC cooling – neglecting bulk Comptonization – in ideal GRMHD simulations (B. E. Kinch et al. 2020), but these have never been applied directly to magnetic reconnection, utilized resistive magnetohydrodynamics, or been compared to kinetic results.

In this letter, we develop, from first principles, a new subgrid model for collisionless relativistic radiative reconnection in resistive relativistic magnetohydrodynamics (RRMHD). The model is based on a set of systematically derived sink terms in the RRMHD energy and momentum equations that capture losses from both IC scattering and bulk Comptonization. We also utilize a charge-starved effective resistivity prescription (A. Moran et al. 2025; B. Ripperda et al. 2026) to mimic the fast reconnection rate of collisionless plasmas. As a first step, we develop our model in the optically thin regime in an isotropic background radiation field. We validate our results against first-principles PIC simulations with the full microphysics of collisionless relativistic radiative reconnection, and find that our RRMHD subgrid model is able to reproduce macroscopic properties – including the fast reconnection rate, the temperature in the reconnection layer, and the bulk motions of the outflow – from PIC. This confirmation of kinetic theory sets the stage for physically-informed global simulations of radiation-dominated systems with emission powered through magnetic reconnection.

2. MODEL FOR RADIATIVE COOLING

2.1. *Radiative drag force on particles*

To model IC cooling in kinetic theory, particles are considered to be travelling through a photon bath described by an isotropic radiation energy density $u_{\rm rad}$, causing the particles to scatter off the background photons. Momentum losses due to IC scattering are calculated by subtracting a radiative drag force $\mathbf{F}_{\rm IC}$ from the Lorentz force experienced by the particles (L. D. Landau & E. M. Lifshitz 1971):

$$\mathbf{F}_{\rm IC} = -\frac{4}{3}\sigma_{\rm T} u_{\rm rad}\gamma^2\boldsymbol{\beta}, \tag{1}$$

where $\sigma_{\rm T}$ is the Thomson cross section, $\boldsymbol{\beta} = \mathbf{v}/c$ is the particle three-velocity $\mathbf{v}$ in units of c, and $\gamma = 1/\sqrt{1-\beta^2}$ is the particle Lorentz factor.

The magnitude of $\mathbf{F}_{\rm IC}$ can be expressed in terms of a characteristic Lorentz factor $\gamma_{\rm rad}$ at which the radiative drag is perfectly balanced against the acceleration supplied by the electric field $E_{\rm rec} = \beta_{\rm rec}B_0$ in the reconnection layer

$$e\beta_{\rm rec}B_0 = \frac{4}{3}\sigma_{\rm T} u_{\rm rad}\gamma_{\rm rad}^2, \tag{2}$$

where $\beta_{\rm rec}$ is the reconnection rate, e is the particle charge, and we have taken $\beta \to 1$ assuming that particles accelerate in the diffusion region (D. A. Uzdensky 2016; L. Sironi & A. M. Beloborodov 2020). Equation (2) gives a convenient parametrization for $u_{\rm rad}$ in terms of $\gamma_{\rm rad}$, so that the strength of the cooling can now be controlled by the characteristic Lorentz factor such that smaller $\gamma_{\rm rad}$ corresponds to larger momentum losses and thus stronger cooling, and vice versa.

This parametrization allows us to rewrite the drag force in terms of dimensionless parameters, the characteristic Lorentz factor, and a length scale, which together fully determine the physical regime of the radiative reconnection. The most meaningful length scale is the half-thickness of the reconnection layer, estimated to be the hot Larmor radius $\Delta \sim r_{\rm L} = \gamma m_{\rm e}c^2/eB_0 \sim B_0/4\pi n_{\rm e,0}e$, using that $\gamma \sim \sigma_0$ in the diffusion regions around the X-points (J. G. Kirk & O. Skjæraasen 2003; G. R. Werner et al. 2016; L. Comisso & L. Sironi 2019; A. Chernoglazov et al. 2023), and where $m_{\rm e}$ is the particle mass and $n_{\rm e,0}$ is the particle number density in the upstream (which we take to be the same for both species). Then, $\mathbf{F}_{\rm IC}$ can be written as

$$\mathbf{F}_{\rm IC} = -\frac{m_{\rm e}c^2\beta_{\rm rec}\sigma_0}{\Delta}\left(\frac{\gamma}{\gamma_{\rm rad}}\right)^2\boldsymbol{\beta}. \tag{3}$$

2.2. *Energy-momentum sink in fluid theory*

The radiative losses in a fluid description of the plasma can be modelled as energy and momentum losses in the form of a sink term applied to the RRMHD equations. Then, the evolution of the plasma is described by the continuity equation

$$\nabla_\nu\left(\rho u^\nu\right) = 0, \tag{4}$$

the equations for energy and momentum conservation,

$$\nabla_\nu T^{\mu\nu} = \frac{1}{c}S^\mu, \tag{5}$$

and Maxwell's equations

$$\nabla_\nu {}^*F^{\mu\nu} = 0, \tag{6}$$

$$\nabla_\nu F^{\mu\nu} = \frac{4\pi}{c}J^\mu. \tag{7}$$

where ρ is the fluid mass density; u^μ is the fluid four-velocity; T^μ_ν is the total stress-energy tensor, including both the fluid and electromagnetic parts; S^μ is the volumetric cooling rate sink term; $F^{\mu\nu}$ is the Faraday tensor and ${}^*F^{\mu\nu}$ is its Hodge dual; and

$$J^\mu = \left\{\varrho c, \varrho\mathbf{v} + \frac{\Gamma}{\eta}\left[\mathbf{E} + \frac{\mathbf{v}\times\mathbf{B}}{c} - \frac{(\mathbf{E}\cdot\mathbf{v})\,\mathbf{v}}{c^2}\right]\right\} \tag{8}$$

is the resistive four-current, with charge density ϱ, Lorentz factor Γ, resistivity η, and velocity, electric, and magnetic field vectors $\mathbf{v}$, $\mathbf{E}$, and $\mathbf{B}$, respectively. This system of equations is closed by the ideal gas equation of state $p = (\hat{\gamma}-1)\rho\epsilon$, relating the pressure p and mass density ρ via the specific internal energy density ϵ and the adiabatic index $\hat{\gamma}$, which we set to 4/3.

The IC losses have both a component due to individual relativistic electrons upscattering the background photons, as well as a bulk Comptonization component due to photons getting upscattered instead by bulk motions of the plasma (i.e., the outflow in a reconnection layer) (A. M. Beloborodov 2017), both of which require a counterpart in a fluid theory. In adapting a fluid representation of IC cooling from kinetic theory, it is necessary to average over the particle distribution function so that individual particle quantities may be written in their analogous fluid form. In particular, averaging over the individual particle motions translates to a temperature and a bulk velocity. To model IC losses, the energy-momentum sink term S^μ should thus have a dependence on both temperature (capturing losses due to electron IC scattering) and bulk velocity (capturing losses due to bulk Comptonization).

Following O. Golan & A. Levinson (2015), the IC radiative sink term S^μ as a volumetric cooling rate for an isotropic radiation field is given by

$$S^0 = -\frac{4}{3}\frac{\sigma_{\rm T}c}{m_{\rm e}}u_{\rm rad}\rho_{\rm e}\Gamma^2\left[\beta_\Gamma^2 + 12\Theta_{\rm e}^2\left(1+\beta_\Gamma^2\right)\right]; \tag{9}$$

$$S^i = -\frac{4}{3}\frac{\sigma_{\rm T}c}{m_{\rm e}}u_{\rm rad}\rho_{\rm e}\Gamma^2\beta_\Gamma^i\left[1 + 4\Theta_{\rm e}^2\left(5+\beta_\Gamma^2\right)\right], \tag{10}$$

where ρ_e is the electron mass density, $\Gamma = 1/\sqrt{1-\beta_\Gamma^2}$ is the Lorentz factor of the bulk motions, β_Γ^i is the bulk three-velocity v_Γ^i in units of c, and $\Theta_e = k_b T_e/m_e c^2$ is the dimensionless electron temperature. A full derivation of Equations (9)-(10) can be found in Appendix A[1].

In order to quantitatively compare the RRMHD radiative losses S^μ to the kinetic radiative drag $\mathbf{F}_{IC}$ (Equation (3)), we use the same force-balance parametrization from Equation (2). We therefore require that the electric field in the reconnection layer E_{rec} matches that in kinetic theory, and thus we introduce an effective resistivity to model the subgrid kinetic length scales. A. Moran et al. (2025) (hereafter M25) proposes such a charge-starved resistivity: if the diffusion regions surrounding the X-points are sufficiently charge-starved, then the corresponding current density there is given by $|\mathbf{J}| \sim 2 n_e e c$, where n_e is the particle number density and the 2 arises from counter-streaming species.

These charge-starved regions can be represented in RRMHD by a localized, effective resistivity $\eta_{eff} = |\mathbf{E}^*|/|\mathbf{J}| \sim \beta_{rec} B_0/2 n_e e c$, where $\mathbf{E}^* = \Gamma\left[\mathbf{E} + (\mathbf{v}\times\mathbf{B})/c - (\mathbf{E}\cdot\mathbf{v})\mathbf{v}/c^2\right]$ (E. G. Blackman & G. B. Field 1993; S. S. Komissarov 2007). Such a prescription has been shown to reproduce the structure and magnitude of $\mathbf{E}^*$, and hence the reconnection rate, from PIC simulations in the non-radiative regime (B. Ripperda et al. 2026, hereafter (R26)). The charge-starved resistivity introduces an effective cold upstream skin depth $\lambda_e = \sqrt{m_e^2 c^2/4\pi\rho_0 e^2}$, parameterized in the numerical prescription by a mass-to-charge ratio m_e/e and upstream mass density ρ_0, such that the typical sheet half-thickness can be written concisely in RRMHD as a product of fundamental plasma parameters $\Delta \sim \sqrt{\sigma_0}\lambda_e$.

To write the force-balance parametrization (Equation (2)) fully in fluid form, we average over the thermal electron distribution function, such that $\langle\gamma^2\rangle = 12\Theta_e^2$ in the relativistic limit (G. B. Rybicki & A. P. Lightman 1979). In the case of a pair plasma[2], we can take $\rho_e = \rho/2$ and $\Theta_e = \Theta$, where ρ and Θ are the RRMHD mass density and dimensionless temperature, respectively. Then, u_{rad} can be written in terms of a characteristic Θ_{rad} as the force-balance in Equation (2) becomes $e\beta_{rec}B_0 = 16\sigma_T u_{rad}\Theta_{rad}^2$. Reparametrizing Equations (9) and (10) in terms of Δ and Θ_{rad}, S^μ becomes

$$S^0 = -\frac{c^3\beta_{rec}\sigma_0}{24\Delta}\rho\Gamma^2\left(\frac{\Theta}{\Theta_{rad}}\right)^2\left[\left(\frac{\beta_\Gamma}{\Theta}\right)^2 + 12\left(1+\beta_\Gamma^2\right)\right]; \tag{11}$$

$$S^i = -\frac{c^3\beta_{rec}\sigma_0}{24\Delta}\rho\Gamma^2\beta_\Gamma^i\left(\frac{\Theta}{\Theta_{rad}}\right)^2\left[\frac{1}{\Theta^2} + 4\left(5+\beta_\Gamma^2\right)\right]. \tag{12}$$

Thus, in the relativistic regime, where the current density in diffusion regions approaches charge starvation (M25; R26), the radiative loss term for RRMHD recovers the parametrization of the radiative drag from kinetic theory, allowing a one-to-one mapping of cooling strengths from PIC → RRMHD via $\gamma_{rad}^2 \to 12\Theta_{rad}^2$.

2.3. *Radiative cooling regimes*

When defining the strength of the cooling in models of radiative reconnection, there are two characteristic length scales of interest: (1) the typical size of a diffusion region, relevant to the distance over which individual particles are accelerated and cooled; and (2) the typical length of the reconnection layer, relevant to bulk Comptonization. To identify the cooling regimes associated with each of these length scales, one must compare the respective crossing time with the typical cooling time (L. Sironi & A. M. Beloborodov 2020):

$$t_{IC} = \frac{m_e c\gamma\boldsymbol{\beta}}{\mathbf{F}_{IC}} = \frac{\Delta\gamma_{rad}^2}{\beta_{rec}\sigma_0 c\gamma}, \tag{13}$$

using Equation (3).

First considering cooling on the scale of a diffusion region, the associated light-crossing time is given by $t_D = \ell_D/c$, where $\ell_D \sim \Delta/\beta_{rec}$ is the length of the diffusion region measured parallel to the reconnection layer, and we estimate the aspect ratio of the diffusion region as $\Delta/\ell_D \sim \beta_{rec}$ (R26). Then the condition for strong cooling on this scale is given by $t_{IC}/t_D \lesssim 1$, or $\gamma_{rad}^2 \lesssim \sigma_0^2$, using that electrons are typically accelerated up to a Lorentz factor $\gamma \sim \sigma_0$ in the diffusion regions in the absence of cooling. Conversely, weak cooling on this scale occurs when $t_{IC}/t_D \gg 1$, or $\gamma_{rad}^2 \gg \sigma_0^2$.

In the case of cooling along the length of the reconnection layer, strong cooling occurs when the cooling time scale is much shorter than the timescale for plasma advection $t_{IC}/t_{adv} \ll 1$, where $t_{adv} \sim L/v_A \sim L/c$, L is the half-length of the sheet, and v_A is the Alfvén speed. In terms of the characteristic Lorentz factor, the condition for strong cooling is then $\gamma_{rad}^2 \ll \beta_{rec} L\sigma_0/\Delta$, where we have assumed $\gamma \sim 1$. Likewise, the cooling on this scale is weak when $t_{IC}/t_{adv} \gg 1$, or $\gamma_{rad}^2 \gg \beta_{rec} L\sigma_0/\Delta$.

We therefore study four separate cooling regimes to cover our entire parameter space: (1) **strong cooling**

[1] Note that in the limit of no bulk motions (i.e., $\Gamma^2 \to 1$, $\beta_\Gamma^2 \to 0$), S^0 recovers the volumetric cooling rate derived by B. E. Kinch et al. (2020) (their Equation (8)) in the relativistic limit, $S^0 = (4/3)\sigma_T n_e c u_{rad}(12\Theta_e^2)$.

[2] If the plasma also consists of ions, an additional electron temperature equation may be included (S. M. Ressler et al. 2015).

Table 1. The characteristic Lorentz factors $\gamma_{\rm rad}$ and dimensionless temperatures $\Theta_{\rm rad}$ corresponding to each of the four cooling regimes defined in Section 2.3 chosen for the simulation runs with magnetization $\sigma_0 = 15$ and half-sheet length $L \approx 3000\lambda_{\rm e}$. The analogous cooling strengths between PIC and RRMHD are calculated using the mapping $\gamma_{\rm rad}^2 \to 12\Theta_{\rm rad}^2$.

Regime	$\gamma_{\rm rad}$ (PIC)	$\Theta_{\rm rad}$ (RRMHD)
Strong cooling $t_{\rm IC} \lesssim t_{\rm D}$	10	2.9
	12	3.5
	14	4.0
	15	4.3
Moderate cooling $t_{\rm D} \ll t_{\rm IC} \ll t_{\rm adv}$	17	4.9
	20	5.8
	32	9.2
Weak cooling $t_{\rm adv} \gg t_{\rm IC} \ll t_{\rm adv}$	60	17.3
No cooling	∞	∞

($t_{\rm IC} \lesssim t_{\rm D}$), (2) **moderate cooling** ($t_{\rm D} \ll t_{\rm IC} \ll t_{\rm adv}$), (3) **weak cooling** ($t_{\rm IC} \gg t_{\rm adv}$), and (4) **no cooling** ($t_{\rm IC} \to \infty$). We calculate values for $\gamma_{\rm rad}$ for each of the four cooling regimes using simulation parameter values $\sigma_0 = 15$, $L \approx 3000\lambda_{\rm e}$ with roughly 5 cells per $\lambda_{\rm e}$, and fiducial $\beta_{\rm rec} = 0.1$. Our corresponding choices for $\gamma_{\rm rad}$ and its mapping to $\Theta_{\rm rad}$ in each regime are summarized in Table 1. For the remainder of this paper, we shall list cooling strenghts using the PIC $\gamma_{\rm rad}$ only, for brevity, and readers may refer to Table 1 for the corresponding RRMHD $\Theta_{\rm rad}$.

These regimes map onto distinct astrophysical sources depending on the role IC scattering plays in each. In black hole coronae, where IC dominates the radiative losses (J. Nättilä 2024; D. Grošelj et al. 2024), particles leave the diffusion regions before radiating (A. M. Beloborodov 2017; L. Sironi & A. M. Beloborodov 2020), yet the IC cooling time is far shorter than the advection time along the layer, placing coronal current sheets in the moderate regime; in this regime, Compton drag on the plasmoids converts most of the dissipated energy into radiation through bulk Comptonization (L. Sironi & A. M. Beloborodov 2020; N. Sridhar et al. 2021). Gamma-ray burst jets, where the radiation energy density exceeds the magnetic energy density, are expected to be qualitatively similar, while the IC-dominated jets of flat-spectrum radio quasars (G. Ghisellini & F. Tavecchio 2009) lie near the boundary between moderate and weak cooling. In contrast, strong cooling at the diffusion regions occurs in the magnetospheric current sheets of young pulsars and M87* through synchrotron losses (A. Chernoglazov et al. 2023; H. Hakobyan et al. 2023b), where IC scattering is dynamically negligible yet can produce the observed TeV emission (H. Hakobyan et al. 2023b).

3. SIMULATION SETUP

We apply our radiative model to RRMHD simulations of a single reconnection layer with a charge-starved resistivity prescription (M25; R26) in Minkowski spacetime using the `Black Hole Accretion Code` (`BHAC`, O. Porth et al. 2017; H. Olivares et al. 2019; B. Ripperda et al. 2019), and benchmark the results against first-principles radiative PIC simulations of pair plasma run with `TRISTAN-MP v2` (A. Spitkovsky 2005; H. Hakobyan et al. 2023).

Both simulation setups feature a 2D Harris current sheet (E. G. Harris 1962) in Cartesian (x, y) coordinates with the sheet parallel to the x-axis. We initialize the PIC and RRMHD simulations such that the initial conditions are as similar as possible. The magnetic field is initialized by the reconnecting magnetic field $B_x = B_0 \tanh(y/d)$ – reversing sign at $y = 0$ over a thickness d of order a few tens of skin depths – and by the uniform guide field aligned with the electric current $B_z = B_g = 0.5B_0$. We include a moderate guide field to represent reconnection in turbulent systems, where idealized anti-parallel configurations (i.e., $B_g \approx 0$) are rare (e.g., V. Zhdankin et al. 2013; A. Chernoglazov et al. 2021; M. Nurisso et al. 2023; D. Grošelj et al. 2024; T. Ha et al. 2025; D. Grošelj et al. 2026; Z. Davis et al. 2026), and to provide magnetic pressure to support plasmoids against collapsing due to radiative losses. The electric and velocity fields are initialized to 0. The initial mass density profile is given by $\rho = \rho_0 + \rho_{\rm s}/\cosh^2(y/d)$ with an over-density in the sheet $\rho_{\rm s} = 3\rho_0$ relative to the upstream density set by the upstream magnetization $\sigma_0 = 15$, chosen to be in the relativistic regime. In RRMHD, the pressure profile is then $p = p_0 + B_0^2/(2\cosh^2(y/d))$ such that the plasma pressure inside the sheet balances the magnetic pressure, and where the upstream pressure is $p_0 = 10^{-2}\rho_0 c^2$ such that the upstream dimensionless temperature in both PIC and RRMHD is set to $\Theta_0 = 10^{-2}$. To trigger reconnection, the pressure balance is disturbed at the centre of the current sheet by a small perturbation, generating an X-point.

Our numerical domain has a length of $2L$ and a comparable effective resolution of $N_x \times N_y = 32,768^2$ in RRMHD (using a base resolution of $N_x \times N_y = 256^2$ with eight levels of adaptive mesh refinement applied to the current sheet region of the domain) and $N_x \times N_y =$

$30,720 \times 30,000$ in PIC. In both RRMHD and PIC, we set a half-box size of $L \approx 3000\lambda_e$, such that we resolve roughly 5 cells per skin depth. We confirm in Appendix B that the effective skin depth scale is well-resolved and hence so is the diffusion region. The boundaries in the outflow direction are initially periodic and switch to absorbing after one light-crossing time $t_c = L/c$ (W. Daughton et al. 2006; L. Sironi et al. 2016; R26), while the boundaries parallel to the sheet employ "plasma injectors" in PIC and absorbing boundaries (effectively replenishing the initial magnetic field and density) in RRMHD. The radiation prescription also turns on after one t_c, where we start solving the radiative source term S^μ (Equations (9)-(10)) explicitly after the resistive source term (B. Ripperda et al. 2019) in RRMHD, and $\mathbf{F}_{\rm IC}$ (Equation (3)) during the particle push in PIC (M. Tamburini et al. 2010; M. Vranic et al. 2016; M. Zou et al. 2025). We set a threshold of $\gamma > \gamma_{\rm thresh} = 1.1$ and $\Gamma > \Gamma_{\rm thresh} = 1.02$ for which to apply the radiation prescription in PIC and RRMHD, respectively[3], in order to avoid cooling the upstream plasma.

4. RESULTS

Here, we demonstrate that our RRMHD cooling model can faithfully reproduce – both qualitatively and quantitatively – the essential macroscopic results of radiative PIC simulations, performed in matching conditions for a given relative cooling strength. For our analysis, we select one representative cooling strength per regime for which to compare: $\gamma_{\rm rad} \in \{\infty, 60, 32, 15\}$ for no, weak, moderate, and strong cooling, respectively. We focus on macroscopic features important for future large-scale GRMHD simulations, rather than microscopic physics only probeable through kinetic theory.

We first examine the structure of the reconnection layer across varying cooling strengths in both the RRMHD and PIC models in Figure 1. For all cooling regimes, RRMHD accurately reproduces the overall morphology of the reconnection layer as well as the typical magnitudes of the bulk speed, plasma temperature, and density in the sheet with respect to the PIC result. Here, $T = m_e p/k_B\rho$ is the plasma temperature, and "bulk speeds" (i.e., the square of the bulk four-velocity, $\Gamma^2\beta_\Gamma^2 = \Gamma^2 - 1$) are found from the bulk Lorentz factor via the bulk three-velocities $\Gamma \equiv \Gamma(v_\Gamma)$; in PIC v_Γ is calculated from the magnitude of the current density $|\mathbf{J}| = \varrho|\mathbf{v}_\Gamma|$, where ϱ is the charge density. The effect of stronger cooling is clearest in temperature – the temperature of the sheet drastically drops with decreasing $\gamma_{\rm rad}$ – with a secondary effect decreasing the bulk motion speeds for decreasing $\gamma_{\rm rad}$ – most noticeable in the plasmoids and at the far edge of the outflow. There is also a decrease in density on the plasmoid faces opposing the direction of the bulk flow with stronger cooling (L. Sironi & A. M. Beloborodov 2020).

We next consider characteristics of the inflow and outflow speeds with respect to the reconnection layer. The inflow speed is found from averaging over slices in the velocity perpendicular to the current sheet, $\beta_{\rm in} = \langle\beta_y\rangle$; while the outflow speed is taken to be the maximum speed parallel to the current sheet, $\beta_{\rm out} = \max(|\beta_x|)$. With the charge-starved resistivity model, RRMHD is able to recover the kinetic reconnection rate $\langle\beta_{\rm rec}\rangle = \langle\beta_{\rm in}/\beta_{\rm out}\rangle \sim 0.08$ when time-averaged over the steady state past the initial transient ($t > 3.5t_c$) and over all four cooling regimes, as shown in the top panel of Figure 2. Note that the reconnection rate is mediated by the guide field and thus is slightly lower than the typically quoted ~ 0.1 rate for zero-guide reconnection (S. Gupta et al. 2024). The steady-state bulk outflow speed is shown in the bottom panel of Figure 2, and is consistent between PIC and RRMHD for all cooling strengths, well-matching the analytic Alfvénic outfow $\Gamma_A\beta_A = \sqrt{B_0^2\sigma_0/(B_0^2 + B_g^2\sigma_0)} \sim \sqrt{\sigma_{\rm eff}}$ (Y. E. Lyubarsky 2005; M. Takamoto & A. Lazarian 2016; L. Sironi et al. 2025), where β_A is the Alfvén speed in units of c and Γ_A is the corresponding Lorentz factor. The strength of the cooling has negligible effect on the reconnection rate or outflow speed, in agreement with previous studies (L. Sironi & A. M. Beloborodov 2020).

The most quantitative comparison between the two models is in the energy spectra. We compare both the kinetic energy $E_{\rm kin}(\gamma)$ spectra and bulk motion energy $E_{\rm bulk}(\Gamma)$ spectra from PIC and RRMHD for each cooling regime in Figure 3. The kinetic energy spectra are calculated from the particle energies in PIC ($E_{\rm kin} = (\gamma - 1)mc^2$). In RRMHD, the kinetic energy spectra are calculated assuming a drifting Maxwellian in each cell, weighted by that cell's density relative to the upstream and cell size relative to the base grid resolution (to account for adaptive mesh refinement), and integrated over the cells in the relevant simulation domain (described below) at each output time step. Additional details regarding the computation of the kinetic energy spectra in RRMHD are provided in Appendix C. The bulk motion energy spectra ($E_{\rm bulk} = (\Gamma - 1)mc^2$) are found from the bulk Lorentz factor in each cell.

[3] Note that the thresholds differ as $\gamma_{\rm thresh}$ in PIC refers to the particle Lorentz factor – which includes the higher-energy particles in the Maxwell-Jüttner distribution, while $\Gamma_{\rm thresh}$ in RRMHD is the bulk Lorentz factor – which is generally lower than the maximum particle Lorentz factor γ.




Figure 1. Comparison of the reconnection layer structure in PIC (**left**) and MHD (**right**) for a range in cooling strengths. Three blocks each display a different quantity: bulk speed $\Gamma^2 - 1$ (**top**), temperature normalized by the upstream T/T_0 **(middle)**, and density normalized by the upstream ρ/ρ_0 overlaid with white contours of the out-of-plane vector potential pointing in the direction of the in-plane magnetic field (**bottom**). A scale-bar indicating the relative size of 30Δ is provided in the top-right panel, indicating the order of the size of the sheet thicknesses. Each block shows the given quantity in three cooling regimes: no cooling ($\gamma_{\rm rad} = \infty$; **top**), weak cooling ($\gamma_{\rm rad} = 60$; **middle**), and strong cooling ($\gamma_{\rm rad} = 15$; **bottom**). With stronger cooling, the temperature in the sheet is noticeably reduced, and the sides of plasmoids opposing the out-flowing motion become less dense. The effect of cooling on the bulk motions is secondary to temperature, but the sheet and plasmoids both show lower Γ with stronger cooling.

Each spectrum is calculated in cells located in the reconnection layer; for the weak and no cooling regimes (where we expect $\Theta \sim \sigma_0$ in the layer) this is identified by cells with a temperature above a threshold $\Theta > \Theta_{\rm tresh} = 3 \times 10^{-2}$, while for the moderate and strong cases the sheet becomes too cold to be identified by temperate alone, so an additional bulk four-velocity threshold is applied $u_x^2 > u_{x,\rm thresh}^2 = 0.1$. The spectra are time-averaged over the steady state of the simulation run from $3.5 \leq t/t_c \leq 10$. Finally, each spectrum is normalized by its own $N_{\rm tot}$ such that it integrates to 1.

The kinetic energy spectra from RRMHD and PIC exhibit similar peaks in the thermal components, indicating good agreement in temperature in all four cooling regimes between the two models. The disparity in the tails between PIC and RRMHD – especially evident




Figure 2. Top: The time-evolution of the reconnection rate $\beta_{\rm rec}$ in each of the four cooling regimes (colours), showing good agreement between PIC (dashed lines) and MHD (solid lines). When time-averaged over the steady state ($t \geq 3.5t_c$; indicated by grey dotted line), we find a rate of $\beta_{\rm rec} = 0.08$ across all simulations shown. **Bottom:** The time-evolution of the four-velocity in the outflow direction $\Gamma_{\rm out}\beta_{\rm out}$. Both MHD and PIC converge to the analytic Alfvénic outflow $\Gamma_{\rm A}\beta_{\rm A} \sim \sqrt{\sigma_{\rm eff}}$ (dotted navy line). The strength of the cooling has negligible effect on $\beta_{\rm rec}$ or $\Gamma_{\rm out}\beta_{\rm out}$.

without cooling in the top panel – is explained by the single-fluid collisional nature of RRMHD and its inability to self-consistently capture particle acceleration as in PIC. The shape of the PIC tail is described by particle acceleration in the diffusion region up to $\gamma \sim \sigma_0$ (dotted vertical blue line in Figure 3), causing a break in the tail, followed by secondary acceleration processes up to $\gamma \sim \gamma_{\rm rad}$ (solid vertical blue line in Figure 3), where the tail is cut off (L. Sironi & A. Spitkovsky 2014). The slope after the break steepens with stronger cooling which has a more significant effect on the nonthermal particles in PIC, driving the distribution closer to that in RRMHD. Thus, stronger cooling shrinks the gap between the spectra from the two models, with both tails fixing to energies described by $\gamma \sim \gamma_{\rm rad}$ and specifically $\gamma \sim \sigma_0$ in the strongest cooling case.

The bulk motion energy spectra in PIC and RRMHD show excellent agreement in both the peak and slope for all cooling strengths, and are consistent with previous studies of guide field-mediated reconnection (S. Gupta et al. 2024). The peak occurs just below the analytic Alfvén Lorentz factor estimate $\Gamma_{\rm A} = 1/\sqrt{1-\beta_{\rm A}^2}$, with $\beta_{\rm A}^2 = \sigma_0/\left[1+\sigma_0\left(1+B_g^2/B_0^2\right)\right]$ (L. Sironi et al. 2025) (dotted vertical blue line in Figure 3), indicating the bulk outflow.




Figure 3. The kinetic energy ($E_{\rm kin}(\gamma)$; solid lines) and bulk motion energy ($E_{\rm bulk}(\gamma)$; dash-dotted lines) spectra for each cooling regime (top four panels) in PIC (pink) and RRMHD (light pink). Vertical lines show $\Gamma = \Gamma_{\rm A}$ (dotted), $\gamma = \sigma_0$ (dashed) and $\gamma = \gamma_{\rm rad}$ (solid). The spectra peak at the Alfvénic outflow $\Gamma_{\rm A}$, and the tails of the energy spectra cool to $\gamma \sim \gamma_{\rm rad}$. All spectra are normalized by $E_{\rm tot}$, illustrating the fraction of total energy supplied by bulk motions. The bulk motion energy spectra match well between RRMHD and PIC for all four cooling regimes; the kinetic energy spectra agree best with smaller $\gamma_{\rm rad}$ when the high energy tail in PIC is cooled to match the thermal RRMHD distribution. The bottom panel shows the kinetic energy and bulk motion energy spectra for all cooling strengths outlined in Table 1 for RRMHD only, highlighting the work of cooling to bring the distributions to lower energies.

RRMHD only recovers the bulk motions energy spectrum with the fast reconnection rate $\beta_{\rm rec} \sim 0.1$, and therefore the charge-starved resistivity model is essential to accurately model the outflow. The tension of the magnetic field lines that determines the outflow and resulting bulk motions is regulated by the aspect ratio of the diffusion regions, which itself is controlled by the reconnection rate (R26). Matching the kinetic rate is therefore not only essential to capture the bulk motion energy spectra accurately, but also directly affects plasmoid dynamics that can result in observable radiative signatures (A. Philippov et al. 2019; J. F. Mahlmann et al. 2022). A comparison to the bulk motion energy spectra produced with the ~ 0.01 reconnection rate with uniform resistivity is provided in Appendix D.

5. CONCLUSIONS

This work marks a first step in modelling kinetic aspects of radiative reconnection in RRMHD with applications to extreme radiative astrophysical environments. We have developed a subgrid model for IC cooling in RRMHD that reproduces the kinetic physics of radiative reconnection – the reconnection rate, plasma temperature, and Compton drag on bulk motions – in agreement with PIC simulations of identical setup across all cooling regimes. The model couples a systematically derived radiative energy and momentum sink, valid in the optically thin regime, with a localized charge-starved resistivity. Our main findings are:

1. **The charge-starved resistivity is essential to reproduce kinetic properties.** The charge-starved resistivity introduces the effective skin depth which links fluid and kinetic cooling strengths, giving the exact mapping $\gamma_{\rm rad}^2 \to 12\Theta_{\rm rad}^2$ for a thermal distribution. It is also required to reproduce the fast reconnection rate (R26) and the bulk-motion statistics (Appendix D).

2. **Our RRMHD radiative losses combined with the charge-starved resistivity successfully recover the plasma temperature, reconnection rate, and bulk motions distribution from kinetic theory.** Our cooling model is also able to match the peak in the kinetic energy spectrum, indicating agreement in plasma temperature in the reconnection layer; and although the non-thermal tail present in kinetic theory is not achievable through RRMHD, we find better agreement between the PIC and RRMHD kinetic energy spectra with stronger cooling, as the cooling acts to reduce the energies to a more thermal-like distribution.

Our framework is built to be extended with additional physical effects:

- **Electron-ion plasmas.** Our simulations invoked a pair ($e^{\pm}$) plasma, and thus our model is missing effects related to multiple species with a mass difference, such as the division of heating from reconnection between ions and electrons. Recent work has focused on two-temperature MHD (S. M. Ressler et al. 2015; M. T. P. Liska et al. 2022) or electron-ion kinetic simulations (D. Grošelj et al. 2026; N. Mancini et al. 2026), and the inclusion of such a model with an adapted two-temperature version of our radiative cooling could make for more realistic radiative reconnection physics in global simulations of compact objects.

- **Additional emission processes and gemoetries.** Our cooling sink can be extended to include other types of emission, such as losses from synchrotron or synchrotron self Compton cooling , or applied to more complex systems, such as a turbulent flow , where reconnection layers form dynamically.

- **Non-thermal particles.** Since RRMHD with charge-starved resistivity captures the magnitude and shape of the non-ideal electric field in the diffusion region (M25; R26), we expect that test particles in RRMHD can faithfully reproduce the non-thermal tail of the PIC spectra even in the non-radiative case, which we will explore in future work.

- **Optically thick systems.** Our cooling model only applies in the optically thin regime; to study optically thicker systems – which may occur in black hole accretion disk coronae, gamma-ray bursts, or magnetar magnetospheres – the plasma-radiation interaction (i.e., the feedback of a dynamically changing photon field on the plasma) should be taken into account (J. C. McKinney et al. 2014; M. T. P. Liska et al. 2022; L. Zhang et al. 2026; C. Nagele et al. 2026). Physical models for pair production and annihilation (e.g., as a source in the continuity equation) or synchrotron self Compton radiation may become important in such applications as well (A. M. Beloborodov 2017; J. Nättilä 2024; H. Hakobyan et al. 2025).

- **Global systems.** Particularly in global simulations, a functional form for the energy density $u_{\rm rad}$

evolved in time and space and sourced by e.g., an accretion disk is required, rather than a constant, isotropic photon field as we have assumed here for simplicity.

Our model is designed for application to global GRMHD simulations of luminous systems – such as in the coronae of accreting black holes, astrophysical jets, or the magnetospheres and winds of compact objects – where the strength of the cooling can be tuned to a specific environment or regime via the free parameter $\gamma_{\rm rad}$. The radiative losses capture IC cooling in diffusion regions as well as Compton drag limiting bulk motions, combined with a charge-starved reconnection rate that sets realistic flaring timescales, altogether supplying physically-motivated cooling to reconnection sites in global simulations.

ACKNOWLEDGMENTS

The authors thank Braden Seefeldt-Gail, Gibwa Musoke, Maëlle Magnan, Charlotte Savelli, Lorenzo Sironi, Andrei Beloborodov, Chris Thompson, Michael Grehan, Sasha Chernoglazov, and Joonas Nättilä for insightful discussions.

AMS, BR and NM are supported in part by the Natural Sciences & Engineering Research Council of Canada (NSERC) [funding reference numbers 568580, RGPIN-2023-04844, and RGPIN-2023-04901], and the Canadian Space Agency (23JWGO2A01). AMS acknowledges the support of NSERC [funding reference numbers 585825-2023, 589240-2024]. Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG), [numéros de référence 568580, RGPIN-2023-04901, 585825-2023, 589240-2024]. AMS was also supported by the Walter C. Sumner Memorial Fellowship and the Mary and Ron Martin Graduate Fellowship in Astrophysics. AMS, BR, AP, and AL acknowledges support by a grant from the Simons Foundation (MP-SCMPS-00001470). DG was supported by the Research Foundation–Flanders (FWO) Senior Postdoctoral Fellowship 12B1424N. AL and AP acknowledge support from a BSF grant (2022314)

The computational resources and services used in this work were partially provided by facilities supported by the VSC (Flemish Supercomputer Center), funded by the Research Foundation Flanders (FWO) and the Flemish Government -– department EWI, and by Compute Ontario and the Digital Research Alliance of Canada (alliancecan.ca) compute allocation rrg-ripperda.

APPENDIX

A. DERIVATION OF THE ENERGY-MOMENTUM SINK TERM

Here we derive for completeness the energy-momentum sinks in Equations (9)–(10) following O. Golan & A. Levinson (2015, hereafter GL15). The sink term for Compton drag acting on a "blob" with lab-frame bulk velocity β^i_Γ is found by averaging over the particle and photon energy distribution functions, and is given by Equation (34) in GL15

$$S^\mu = -\sigma_{\rm T} c^2 \int \frac{{\rm d}^3 k}{k^0} \int \frac{{\rm d}^3 p}{p^0} f_{\rm r} f_{\rm e} p_\nu k^\nu \left[k^\mu + \frac{(p_\nu k^\nu)\, p^\mu}{m_{\rm e}^2 c^2} \right], \tag{A1}$$

where $p^\mu = (\gamma m_{\rm e} c, \gamma m_{\rm e} c \beta^i)$ is the particle four-momentum, $k^\mu = (k, k^i)$ is the photon four-momentum, $f_{\rm e} \equiv f_{\rm e}(p^i)$ is the electron distribution function, and $f_{\rm r}$ is the photon distribution function. We calculate the source term in the rest (primed) frame of the blob where $f_{\rm e}$ is isotropic, and then transform back to the lab (unprimed) frame at the end.

We begin with the energy component of the sink term S'^0. Using $p'_\nu k'^\nu = -p'^0 k'^0 + p'_i k'^i = -p'^0 k'^0 + p' k' \mu' = -p'^0 k'^0 (1 - \beta' \mu')$, since $k'^0 = k'$, and where μ' is the cosine of the scattering angle, S'^0 becomes

$$S'^0 = -\sigma_{\rm T} c^2 \int {\rm d}^3 k' f'_{\rm r} \int {\rm d}^3 p' f'_{\rm e} \frac{p'_\nu k'^\nu}{p'^0 k'^0} \left[k'^0 + \frac{(p'_\nu k'^\nu)\, p'^0}{m_{\rm e}^2 c^2} \right] \tag{A2}$$

$$= -\sigma_{\rm T} c^2 \int {\rm d}^3 k' f'_{\rm r} \int {\rm d}^3 p' f'_{\rm e} \left[-k'^0 (1 - \beta' \mu') + \frac{\left(p'^0\right)^2 k'^0 \left(1 - \beta' \mu'\right)^2}{m_{\rm e}^2 c^2} \right] \tag{A3}$$

$$= \sigma_{\rm T} c^2 \int {\rm d}^3 k'\, k'^0 f'_{\rm r} \int {\rm d}^3 p' f'_{\rm e} \left[(1 - \beta' \mu') - \gamma'^2 \left(1 - \beta' \mu'\right)^2 \right]. \tag{A4}$$

Rewriting $f'_{\rm e}(p'^i)$ as $\tilde{f}'_{\rm e}(p',\mu')$, the momentum integrals become $\iiint {\rm d}^3p' f'_{\rm e} \to \iint {\rm d}\mu'\,{\rm d}p'\,p'^2\tilde{f}'_{\rm e}$, where the factor of 2π from averaging over ϕ' has been absorbed into $\tilde{f}'_{\rm e}$, and the energy component of the sink term further simplifies to

$$S'^0 = \sigma_{\rm T}c^2 \int {\rm d}^3k'\,k'^0 f'_{\rm r} \int {\rm d}p'\,p'^2\tilde{f}'_{\rm e}\left[2 - 2\gamma'^2\left(1+\frac{\beta'^2}{3}\right)\right] \tag{A5}$$

$$= -2\sigma_{\rm T}c^2 \int {\rm d}^3k'\,k'^0 f'_{\rm r} \int {\rm d}p'\,p'^2\tilde{f}'_{\rm e}\left[\frac{4}{3}\gamma'^2\beta'^2\right], \tag{A6}$$

where we used $\gamma'^2 - 1 = \gamma'^2\beta'^2$.

Now, with $n'_{\rm e} = \int {\rm d}^3p'\,f'_{\rm e} = 2\int {\rm d}p'\,p'^2\tilde{f}'_{\rm e}$ and the radiation energy-momentum tensor $T'^{\mu\nu}_{\rm r} = c\int {\rm d}^3k'(k'^\mu k'^\nu/k'^0)f'_{\rm r}$, the energy component of the sink term becomes

$$S'^0 = -\frac{4}{3}\sigma_{\rm T}cT'^{00}_{\rm r}n'_{\rm e}\langle\gamma'^2\beta'^2\rangle. \tag{A7}$$

For an isotropic radiation field in the lab frame, the radiation energy-momentum tensor is given by

$$T^{00}_{\rm r} = u_{\rm rad}; \qquad T^{0i}_{\rm r} = 0; \qquad T^{ij}_{\rm r} = \frac{u_{\rm rad}}{3}\delta_{ij}; \tag{A8}$$

where δ_{ij} is the Kronecker delta function. Then, to find $T'^{00}_{\rm r}$ in the rest frame, we apply the Lorentz transformation such that $T'^{00}_{\rm r} = \Lambda^0_\mu\Lambda^0_\nu T^{\mu\nu}_{\rm r} = \Gamma^2 u_{\rm rad}(1+\beta^2_\Gamma/3)$, where Γ is the bulk Lorentz factor and β^i_Γ is the bulk three-velocity in units of c. This gives the final form of the energy component in the rest frame

$$S'^0 = -\frac{4}{3}\sigma_{\rm T}cn'_{\rm e}u_{\rm rad}\left\langle\gamma'^2\beta'^2\right\rangle\Gamma^2\left[1+\frac{1}{3}\beta^2_\Gamma\right], \tag{A9}$$

as in Equation (45) from GL15.

Turning now to the momentum component of the sink term, we follow the same steps:

$$S'^i = -\sigma_{\rm T}c^2 \int {\rm d}^3k'\,f'_{\rm r}\int {\rm d}^3p'\,f'_{\rm e}\frac{p'_\nu k'^\nu}{p'^0k'^0}\left[k'^i + \frac{(p'_\nu k'^\nu)\,p'^i}{m^2_{\rm e}c^2}\right] \tag{A10}$$

Transforming again $f'_{\rm e}(p') \to \tilde{f}'_{\rm e}(p',\mu')$, we average both k'^i and β'^i over ϕ'. Choosing the local z'-axis to be in the $\hat{k}'$ direction, $\left\langle k'^i\right\rangle_\phi = \int {\rm d}\phi' k'^i = k'\mu'\hat{k}' = k'^i$ and $\left\langle\beta'^i\right\rangle_\phi = \int {\rm d}\phi'\beta'^i = \beta'\mu'\hat{k}'$, with the 2π factor again absorbed into $\tilde{f}'_{\rm e}$, and so

$$S'^i = \sigma_{\rm T}c^2 \int {\rm d}^3k'\,f'_{\rm r}\int {\rm d}p'p'^2\tilde{f}'_{\rm e}\int {\rm d}\mu'\,(1-\beta'\mu')\left[\left\langle k'^i\right\rangle_\phi - \gamma'^2k'\left\langle\beta'^i\right\rangle_\phi(1-\beta'\mu')\right] \tag{A11}$$

$$= \sigma_{\rm T}c^2 \int {\rm d}^3k'\,f'_{\rm r}\int {\rm d}p'p'^2\tilde{f}'_{\rm e}\int {\rm d}\mu'\,(1-\beta'\mu')\left[k'^i - \gamma'^2\beta'\mu'k'^i(1-\beta'\mu')\right] \tag{A12}$$

$$= \sigma_{\rm T}c^2 \int {\rm d}^3k'k'^i f'_{\rm r}\int {\rm d}p'p'^2\tilde{f}'_{\rm e}\left(2+\frac{4}{3}\gamma'^2\beta'^2\right) \tag{A13}$$

$$= \sigma_{\rm T}cn'_{\rm e}T'^{0i}_{\rm r}\left\langle 1+\frac{2}{3}\gamma'^2\beta'^2\right\rangle. \tag{A14}$$

After boosting $T'^{0i}_{\rm r}$ in the same way, we arrive at the final expression for the momentum component of the sink term in the blob's rest frame

$$S'^i = -\frac{4}{3}\sigma_{\rm T}cn'_{\rm e}u_{\rm rad}\Gamma^2\beta^i_\Gamma\left\langle 1+\frac{2}{3}\gamma'^2\beta'^2\right\rangle \tag{A15}$$

as in Equation (46) from GL15.

We perform one final Lorentz transformation to S'^μ to obtain the sink term in the lab frame. Starting with S'^0, this is simply

$$S^0 = \Lambda^0_0 S'^0 + \Lambda^0_i S'^i = \Gamma S'^0 + \Gamma\beta^i_\Gamma S'^i, \tag{A16}$$




Figure 4. Right: Zoom-in of a diffusion region around an X-point in the RRMHD simulation with $\gamma_{\mathrm{rad}} = 15$, showing contours of the out-of-plane vector potential pointing in the direction of the in-plane magnetic field (white lines). **Left:** Slice through the diffusion region (indicated by the pink dashed line in the right panel), showing $\langle\beta_{\mathrm{rec}}\rangle \sim c\eta|J_z|/v_{\mathrm{A}}\left(B_0 + B_g^2/B_0\right)$. The separation between two matching horizontal lines illustrates critical length scales – the cell size $\mathrm{d}x$ (dotted blue), the effective skin depth $\lambda_{\mathrm{e},0}$ (dash-dotted violet), and the sheet half-thickness $\Delta = \lambda_{\mathrm{e},0}\sqrt{\sigma_{\mathrm{eff}}}$ (solid green) – indicating the resistive scale is well-resolved in our RRMHD simulations.

since we are now boosting by $-\beta_\Gamma^i$. For S'^i, we first apply the boost:

$$S^i = \Lambda_0^i S'^0 + \Lambda_j^i S'^j \tag{A17}$$

$$= \Gamma\beta_\Gamma^i S'^0 + S'^j\left(\delta_{ij} + (\Gamma-1)\frac{\beta_\Gamma^i\beta_\Gamma^j}{(\beta_\Gamma)^2}\right) \tag{A18}$$

$$= \Gamma\beta_\Gamma^i S'^0 + S'^i + (\Gamma-1)\,S'^j\frac{\beta_\Gamma^i\beta_\Gamma^j}{(\beta_\Gamma)^2}. \tag{A19}$$

Focusing now on the last two terms, we notice $S'^j\beta_\Gamma^j\beta_\Gamma^i/(\beta_\Gamma)^2 = S'^j\hat{\beta}_\Gamma\hat{\beta}_\Gamma = S'^j_\parallel$. Then the above becomes

$$S^i = \Gamma\beta_\Gamma^i S'^0 + S'^i + \Gamma S'^i_\parallel - S'^i_\parallel \tag{A20}$$

$$= \Gamma\beta_\Gamma^i S'^0 + \Gamma S'^i_\parallel + S'^i_\perp. \tag{A21}$$

Since the bulk velocity is entirely along $S'^i_\parallel$, then $S'^i_\perp = 0$ and $S'^i_\parallel = S'^i$, and so

$$S^i = \Gamma\beta_\Gamma^i S'^0 + \Gamma S'^i. \tag{A22}$$

Applying the Lorentz transformations from Equations (A16) and (A22) to S'^0 and S'^i (Equations (A9) and (A15), respectively) and rewriting the number density n_{e} as a mass density ρ_{e}, we finally arrive at the sink terms in the lab frame shown in Equations (9) and (10) of the main text:

$$S^0 = -\frac{4}{3}\frac{\sigma_{\mathrm{T}}c}{m_{\mathrm{e}}}u_{\mathrm{rad}}\rho_{\mathrm{e}}\Gamma^2\left(\beta_\Gamma^2 + 12\Theta_{\mathrm{e}}^2\left(1+\beta_\Gamma^2\right)\right); \tag{A23}$$

$$S^i = -\frac{4}{3}\frac{\sigma_{\mathrm{T}}c}{m_{\mathrm{e}}}u_{\mathrm{rad}}\rho_{\mathrm{e}}\Gamma^2\beta_\Gamma^i\left(1 + 4\Theta_{\mathrm{e}}^2\left(5+\beta_\Gamma^2\right)\right), \tag{A24}$$

where we have used $\left\langle\gamma'^2\beta'^2\right\rangle = 12\Theta_{\mathrm{e}}^2$ (dropping the prime as temperature is defined in the rest frame).

B. RESOLVING THE DIFFUSION REGION IN RRMHD

To confirm the resistive scale is resolved even with strong cooling in RRMHD, we must ensure we are resolving the effective skin depth λ_{e} with a sufficient number of numerical cells. Based on requirements from PIC simulations (L.

Sironi et al. 2025), as well as RRMHD without cooling (R26), we employ 5 cells/λ_e. Our base resolution in RRMHD is 256^2 with eight levels of adaptive mesh refinement, resulting in an effective resolution of $32,768^2$. The effective skin depth is $\lambda_e \sim m/e/\sqrt{\rho_0}$, with the initial upstream density $\rho_0 = B_0^2/\{\sigma_0[1+\hat{\gamma}\,\Theta_0/(\hat{\gamma}+1)]\}$, and where m/e is a free parameter, which should be set such that $\lambda_e \ll L$ (R26), similar as in PIC simulations. With our simulation parameters for initial upstream magnetic field $B_0 = 1$, magnetization $\sigma_0 = 15$, dimensionless temperature $\Theta_0 = 0.01$, adiabatic index $\hat{\gamma} = 4/3$, and mass-to-charge ratio $m/e = 8\times10^{-5}$, we find $\lambda_e \approx 5\mathrm{d}x \approx 0.0003L$, or equivalently $\Delta_0 = \lambda_{e,0}\sqrt{\sigma_{\rm eff}} \approx 0.001L$, where L is the half-length of the simulation domain, Δ_0 is the initial half-thickness of the current sheet, $\lambda_{e,0}$ is the initial upstream cold effective skin depth, and $\sigma_{\rm eff} \sim \Gamma_A\beta_A$ is the effective magnetization with a guide field (L. Sironi et al. 2025).

We further illustrate the resolution of our RRMHD simulations in Figure 4 through a zoom-in of a diffusion region surrounding an X-point. The right panel shows the effective reconnection electric field $c\eta|J_z|/v_A\left(B_0 + B_g^2/B_0\right) \sim \beta_{\rm rec}$ in the diffusion region. Here, $|J_z|$ is the magnitude out-of-plane resistive current density:

$$J_z = \varrho v_z + \frac{\Gamma}{\eta}\left[E_z + \frac{(\mathbf{v}\times\mathbf{B})_z}{c} - \frac{(\mathbf{E}\cdot\mathbf{v})\,v_z}{c^2}\right], \tag{B25}$$

where ϱ is the charge density (E. G. Blackman & G. B. Field 1993; S. S. Komissarov 2007). In the left panel, we show a vertical slice through the diffusion region and compare it to the time-averaged $\langle\beta_{\rm rec}\rangle \sim 0.08$. Overtop of the slice of the diffusion region, we show three length scales of interest to the resolution through the distance between pairs of lines: the cell size $\mathrm{d}x$, the initial effective skin depth $\lambda_{e,0}$, and the initial current sheet half-thickness $\Delta_0 = \lambda_{e,0}\sqrt{\sigma_{\rm eff}}$. The thickness of the diffusion region is larger than $\lambda_{e,0}$ and is shown to approach the expected Δ_0 (R26). The size of $\mathrm{d}x$ compared to both $\lambda_{e,0}$ and Δ_0 confirms the diffusion region is well-resolved for our effective resolution in RRMHD.

C. DERIVATION OF THE DRIFTING MAXWELLIAN

Here we derive the drifting Maxwellian calculated in each RRMHD cell to produce the kinetic energy spectrum shown in Figure 3. We first derive the expression analytically by boosting the local MJ distribution to the lab frame and integrating over the angle between the particle's motion and the bulk motion. We then show the simplified expression used to calculate the spectrum numerically.

C.1. *Analytical expression*

S. Zenitani (2015) has shown that the MJ distribution function in terms of four-velocity $\mathbf{u}$ transforms as

$$f'(\mathbf{u}') = f(\mathbf{u}) = \frac{1}{4\pi\Theta K_2\left(\frac{1}{\Theta}\right)}\exp\left(-\frac{\gamma'}{\Theta}\right) \tag{C26}$$

$$= \frac{1}{4\pi\Theta K_2\left(\frac{1}{\Theta}\right)}\exp\left(-\frac{\Gamma\gamma\,(1-\beta_\Gamma\beta\mu)}{\Theta}\right), \tag{C27}$$

where $\Theta = k_B T/m_e c^2$ is the dimensionless electron temperature, K_2 is the modified Bessel function of the second kind, Γ is the bulk lorentz factor, β_Γ is the bulk three-velocity in units of c, γ is the particle Lorentz factor, β is the particle three-velocity in units of c, μ is the cosine of the angle between β_Γ and β, and we again indicate the rest frame as primed and the lab frame as unprimed. Averaging over μ yields

$$f(\mathbf{u}) = \frac{1}{4\pi\Theta K_2\left(\frac{1}{\Theta}\right)}\frac{1}{2}\int_{-1}^{1}\exp\left(-\frac{\Gamma\gamma\,(1-\beta_\Gamma\beta\mu)}{\Theta}\right)\mathrm{d}\mu \tag{C28}$$

$$= \frac{1}{4\pi K_2\left(\frac{1}{\Theta}\right)}\frac{\sinh\left(\sqrt{\Gamma^2-1}\sqrt{\gamma^2-1}/\Theta\right)}{\sqrt{\Gamma^2-1}\sqrt{\gamma^2-1}}\exp\left(-\frac{\Gamma\gamma}{\Theta}\right), \tag{C29}$$

$$\tag{C30}$$

using $\gamma\beta = \sqrt{\gamma^2-1}$.

Since $f(\mathbf{u})\mathrm{d}^3u = f(\gamma)\mathrm{d}\gamma\,\mathrm{d}^3u/\mathrm{d}\gamma = 4\pi\gamma\sqrt{\gamma^2-1}f(\gamma)\mathrm{d}\gamma$, the final expression for the drifting Maxwellian in terms of γ in the lab frame is given by

$$f(\gamma)\mathrm{d}\gamma = \frac{\gamma\sinh\left(\sqrt{\Gamma^2-1}\sqrt{\gamma^2-1}/\Theta\right)}{K_2\left(\frac{1}{\Theta}\right)\sqrt{\Gamma^2-1}}\exp\left(-\frac{\Gamma\gamma}{\Theta}\right)\mathrm{d}\gamma. \tag{C31}$$

C.2. *Numerical calculation*

Equivalently, the boosted MJ distribution can be derived as follows. We begin with the normalized energy distribution in the fluid rest frame, where the fluid is isotropic:

$$\mathrm{d}N = n' f'(\gamma')\,\mathrm{d}\gamma' = \frac{n'}{\Theta K_2\left(\frac{1}{\Theta}\right)}\gamma'^2\beta'(\gamma')\exp\left(-\frac{\gamma'}{\Theta}\right)\mathrm{d}\gamma'. \tag{C32}$$

Since particle number is invariant,

$$\mathrm{d}N = \left(\frac{\mathrm{d}N}{\mathrm{d}\gamma'}\right)_{\mathrm{rest}}\mathrm{d}\gamma' = \left(\frac{\mathrm{d}N}{\mathrm{d}\gamma}\right)_{\mathrm{lab}}\mathrm{d}\gamma, \tag{C33}$$

so,

$$f(\gamma) = \frac{n'}{n}f'(\gamma')\frac{\mathrm{d}\gamma'}{\mathrm{d}\gamma} = \frac{f'(\gamma')}{\Gamma}\frac{\mathrm{d}\gamma'}{\mathrm{d}\gamma}. \tag{C34}$$

In the relativisitic limit where $\beta'(\gamma') \to 1$, we have $\mathrm{d}\gamma'/\mathrm{d}\gamma \to \gamma'/\gamma$, so that the boosted MJ in the lab frame is given by

$$f(\gamma)\mathrm{d}\gamma = \frac{\gamma'}{\Gamma\gamma}f'(\gamma')\mathrm{d}\gamma' \tag{C35}$$

$$= \frac{1}{\Gamma^2(1-\beta_\Gamma\beta'\mu')}f'(\gamma')\mathrm{d}\gamma' \tag{C36}$$

$$= \frac{\gamma'^2\beta'}{\Gamma^2\left(1-\beta_\Gamma\beta'\mu'\right)\Theta K_2\left(\frac{1}{\Theta}\right)}\exp\left(-\frac{\gamma}{\Theta}\right)\mathrm{d}\gamma', \tag{C37}$$

which can be numerically binned over a range of rest-frame γ' and μ' to obtain the energy spectrum for a given cell Θ and lab-frame γ.

D. UNIFORM RESISTIVITY

The main result of this paper is the reproduction of the PIC macroscopic quantities – i.e., the reconnection rate, temperature in the reconnection layer, and bulk motions energy spectra – in RRMHD with IC cooling. The charge-starved resistivity model is essential in replicating PIC results particularly in the reconnection rate and bulk motion energy distribution. In Figure 5 we compare the reconnection rate (left panel) and bulk motions spectrum (right panel) for an RRMHD simulation with the uniform resistivity prescription to an RRMHD simulation using the charge-starved resistivity and a PIC simulation. All simulations have cooling turned off to isolate the effect of the η prescription and a comparable resolution of $N_x \times N_y = 32,768^2$ for both RRMHD simulations and $N_x \times N_y = 30,720 \times 30,000$ in PIC. With the uniform resistivity model, the reconnection rate is ~ 4 times slower, and while the maximum value for the bulk spectra remains the same (as it is governed by the Alfvénic outflow) the peak is less defined and orders of magnitude less energetic. The reconnection rate governs the aspect ratio of the diffusion region and thus the field line tension which drives the outflows, and so the slower rate from the uniform resistivity prescription produces a slower average bulk outflow. Therefore, uniform resistivity prescriptions or ideal MHD with a uniform numerical resistivity (M. P. Grehan et al. 2025) are unable to capture both the kinetic β_{rec} and thus the correct bulk motions, and hence ideal GRMHD (e.g., N. Sridhar et al. 2025) is not sufficient to model bulk motions due to reconnection in global accretion simulations.

REFERENCES

Antolin, P., Pagano, P., Testa, P., Petralia, A., & Reale, F. 2021, Nature Astronomy, 5, 54, doi: 10.1038/s41550-020-1199-8

Beloborodov, A. M. 2017, ApJ, 850, 141, doi: 10.3847/1538-4357/aa8f4f

Blackman, E. G., & Field, G. B. 1993, PhRvL, 71, 3481, doi: 10.1103/PhysRevLett.71.3481

Cerutti, B., & Philippov, A. A. 2017, A&A, 607, A134, doi: 10.1051/0004-6361/201731680

Cerutti, B., Philippov, A. A., & Dubus, G. 2020, A&A, 642, A204, doi: 10.1051/0004-6361/202038618

Cheng, X., Priest, E. R., Li, H. T., et al. 2023, Nature Communications, 14, 2107, doi: 10.1038/s41467-023-37888-w




Figure 5. Left: The effect of the charge-starved resistivity on the reconnection rate $\beta_{\rm rec}$. Here we plot $\beta_{\rm rec}$ for the uniform resistivity model with no cooling (green) over the top panel of Figure 2, comparing it to the fast rate from kinetic simulations or MHD with charge-starved resistivity. For the kinetic or charge-starved RRMHD, the steady-state time-averaged reconnection rate is $\langle\beta_{\rm rec}\rangle = 0.08$ (dotted navy horizontal line); for the uniform resistivity model, $\langle\beta_{\rm rec}\rangle = 0.02$ (horizontal green line), four times slower. **Right:** The bulk motions spectrum $E_{\rm bulk}(\Gamma)$ for the no cooling regime in PIC (pink dotted line), charge-starved resistive RRMHD (light pink solid line), and uniform resistivity RRMHD (dash-dotted green line). The uniform resistivity model cannot reproduce the same peak in the bulk motions spectrum as the kinetic simulation ($\sim \Gamma_{\rm A}$), but the charge-starved resistivity model can. Note the lower energy peak in the charge-starved RRMHD spectrum corresponds to the Lorentz factor of the inflow $\Gamma \sim \Gamma_{\rm in} = 1/\sqrt{1-\beta_{\rm in}^2} \approx 2 \times 10^{-3}$.

Chernoglazov, A., Hakobyan, H., & Philippov, A. 2023, ApJ, 959, 122, doi: 10.3847/1538-4357/acffc6

Chernoglazov, A., Ripperda, B., & Philippov, A. 2021, ApJL, 923, L13, doi: 10.3847/2041-8213/ac3afa

Comisso, L., & Sironi, L. 2019, ApJ, 886, 122, doi: 10.3847/1538-4357/ab4c33

Daughton, W., Scudder, J., & Karimabadi, H. 2006, Physics of Plasmas, 13, 072101, doi: 10.1063/1.2218817

Davis, Z., Comisso, L., Haggerty, C., & Nättilä, J. 2026, ApJ, 999, 261, doi: 10.3847/1538-4357/ae3d94

Eardley, D. M., Lightman, A. P., & Shapiro, S. L. 1975, ApJL, 199, L153, doi: 10.1086/181871

Fabian, A. C., Lohfink, A., Kara, E., et al. 2015, MNRAS, 451, 4375, doi: 10.1093/mnras/stv1218

Galishnikova, A., Philippov, A., Quataert, E., et al. 2023, PhRvL, 130, 115201, doi: 10.1103/PhysRevLett.130.115201

Ghisellini, G., & Tavecchio, F. 2009, MNRAS, 397, 985, doi: 10.1111/j.1365-2966.2009.15007.x

Golan, O., & Levinson, A. 2015, ApJ, 809, 23, doi: 10.1088/0004-637X/809/1/23

Granot, J. 2016, ApJL, 816, L20, doi: 10.3847/2041-8205/816/2/L20

Grehan, M. P., Ghosal, T., Beattie, J. R., et al. 2025, PhRvD, 112, 063046, doi: 10.1103/8xf2-x2nq

Grošelj, D., Hakobyan, H., Beloborodov, A. M., Sironi, L., & Philippov, A. 2024, PhRvL, 132, 085202, doi: 10.1103/PhysRevLett.132.085202

Grošelj, D., Philippov, A., Beloborodov, A. M., & Mushotzky, R. 2026, ApJ, 1001, 64, doi: 10.3847/1538-4357/ae50fc

Guo, F., Li, H., Daughton, W., & Liu, Y.-H. 2014, PhRvL, 113, 155005, doi: 10.1103/PhysRevLett.113.155005

Gupta, S., Sridhar, N., & Sironi, L. 2024, MNRAS, 527, 6065, doi: 10.1093/mnras/stad3573

Ha, T., Nättilä, J., Davelaar, J., & Sironi, L. 2025, ApJL, 985, L31, doi: 10.3847/2041-8213/add47b

Hakobyan, H., Levinson, A., Sironi, L., Philippov, A., & Ripperda, B. 2025, ApJL, 995, L73, doi: 10.3847/2041-8213/ae286a

Hakobyan, H., Philippov, A., & Spitkovsky, A. 2019, ApJ, 877, 53, doi: 10.3847/1538-4357/ab191b

Hakobyan, H., Philippov, A., & Spitkovsky, A. 2023a, ApJ, 943, 105, doi: 10.3847/1538-4357/acab05
Hakobyan, H., Ripperda, B., & Philippov, A. A. 2023b, ApJL, 943, L29, doi: 10.3847/2041-8213/acb264
Hakobyan, H., Spitkovsky, A., Chernoglazov, A., et al. 2023,, v2.6 Zenodo, doi: 10.5281/zenodo.7566725
Harris, E. G. 1962, Il Nuovo Cimento, 23, 115, doi: 10.1007/BF02733547
Kinch, B. E., Noble, S. C., Schnittman, J. D., & Krolik, J. H. 2020, ApJ, 904, 117, doi: 10.3847/1538-4357/abc176
Kirk, J. G., & Skjæraasen, O. 2003, ApJ, 591, 366, doi: 10.1086/375215
Komissarov, S. S. 2007, MNRAS, 382, 995, doi: 10.1111/j.1365-2966.2007.12448.x
Landau, L. D., & Lifshitz, E. M. 1971, The classical theory of fields
Levinson, A. 2006, International Journal of Modern Physics A, 21, 6015, doi: 10.1142/S0217751X06035063
Liska, M. T. P., Musoke, G., Tchekhovskoy, A., Porth, O., & Beloborodov, A. M. 2022, ApJL, 935, L1, doi: 10.3847/2041-8213/ac84db
Lyubarsky, Y. 2010, ApJL, 725, L234, doi: 10.1088/2041-8205/725/2/L234
Lyubarsky, Y. 2020, ApJ, 897, 1, doi: 10.3847/1538-4357/ab97b5
Lyubarsky, Y. E. 2005, MNRAS, 358, 113, doi: 10.1111/j.1365-2966.2005.08767.x
Mahlmann, J. F. 2024, arXiv e-prints, arXiv:2405.02773, doi: 10.48550/arXiv.2405.02773
Mahlmann, J. F., Philippov, A. A., Levinson, A., Spitkovsky, A., & Hakobyan, H. 2022, ApJL, 932, L20, doi: 10.3847/2041-8213/ac7156
Mancini, N., Comisso, L., & Sironi, L. 2026, arXiv e-prints, arXiv:2609.21118, doi: 10.48550/arXiv.2609.21118
McKinney, J. C., Tchekhovskoy, A., Sadowski, A., & Narayan, R. 2014, MNRAS, 441, 3177, doi: 10.1093/mnras/stu762
McKinney, J. C., & Uzdensky, D. A. 2012, MNRAS, 419, 573, doi: 10.1111/j.1365-2966.2011.19721.x
Mehlhaff, J., Werner, G., Cerutti, B., Uzdensky, D., & Begelman, M. 2024, MNRAS, 527, 11587, doi: 10.1093/mnras/stad3863
Mehlhaff, J. M., Chen, A. Y., Luepker, M., & Yuan, Y. 2026, arXiv e-prints, arXiv:2602.22168, doi: 10.48550/arXiv.2602.22168
Mehlhaff, J. M., Werner, G. R., Uzdensky, D. A., & Begelman, M. C. 2021, MNRAS, 508, 4532, doi: 10.1093/mnras/stab2745
Moran, A., Sironi, L., Levis, A., et al. 2025, ApJL, 978, L45, doi: 10.3847/2041-8213/ada158
Nagele, C., Krolik, J. H., Liu, R., Kinch, B. E., & Schnittman, J. D. 2026, ApJ, 1001, 190, doi: 10.3847/1538-4357/ae4ec7
Nättilä, J. 2024, Nature Communications, 15, 7026, doi: 10.1038/s41467-024-51257-1
Nurisso, M., Celotti, A., Mignone, A., & Bodo, G. 2023, MNRAS, 522, 5517, doi: 10.1093/mnras/stad1348
Olivares, H., Porth, O., Davelaar, J., et al. 2019, A&A, 629, A61, doi: 10.1051/0004-6361/201935559
Parfrey, K., Philippov, A., & Cerutti, B. 2019, PhRvL, 122, 035101, doi: 10.1103/PhysRevLett.122.035101
Parsons, J., Spitkovsky, A., Philippov, A., & Hakobyan, H. 2026, arXiv e-prints, arXiv:2604.22059, doi: 10.48550/arXiv.2604.22059
Patel, R., Niembro, T., Xie, X., et al. 2025, Nature Astronomy, 9, 1444, doi: 10.1038/s41550-025-02623-6
Petropoulou, M., Giannios, D., & Sironi, L. 2016, MNRAS, 462, 3325, doi: 10.1093/mnras/stw1832
Philippov, A., Uzdensky, D. A., Spitkovsky, A., & Cerutti, B. 2019, ApJL, 876, L6, doi: 10.3847/2041-8213/ab1590
Philippov, A. A., & Spitkovsky, A. 2018, ApJ, 855, 94, doi: 10.3847/1538-4357/aaabbc
Porth, O., Olivares, H., Mizuno, Y., et al. 2017, Computational Astrophysics and Cosmology, 4, 1, doi: 10.1186/s40668-017-0020-2
Ressler, S. M., Tchekhovskoy, A., Quataert, E., Chandra, M., & Gammie, C. F. 2015, MNRAS, 454, 1848, doi: 10.1093/mnras/stv2084
Ripperda, B., Grehan, M. P., Moran, A., et al. 2026, arXiv e-prints, arXiv:2601.02460, doi: 10.48550/arXiv.2601.02460
Ripperda, B., Liska, M., Chatterjee, K., et al. 2022, ApJL, 924, L32, doi: 10.3847/2041-8213/ac46a1
Ripperda, B., Bacchini, F., Porth, O., et al. 2019, ApJS, 244, 10, doi: 10.3847/1538-4365/ab3922
Rybicki, G. B., & Lightman, A. P. 1979, Radiative processes in astrophysics
Schoeffler, K. M., Grismayer, T., Uzdensky, D., & Silva, L. O. 2023, MNRAS, 523, 3812, doi: 10.1093/mnras/stad1588
Shapiro, S. L., Lightman, A. P., & Eardley, D. M. 1976, ApJ, 204, 187, doi: 10.1086/154162
Sironi, L., & Beloborodov, A. M. 2020, ApJ, 899, 52, doi: 10.3847/1538-4357/aba622
Sironi, L., Giannios, D., & Petropoulou, M. 2016, MNRAS, 462, 48, doi: 10.1093/mnras/stw1620
Sironi, L., Rowan, M. E., & Narayan, R. 2021, ApJL, 907, L44, doi: 10.3847/2041-8213/abd9bc
Sironi, L., & Spitkovsky, A. 2014, ApJL, 783, L21, doi: 10.1088/2041-8205/783/1/L21

Sironi, L., Uzdensky, D. A., & Giannios, D. 2025, ARA&A, 63, 127, doi: 10.1146/annurev-astro-020325-115713

Spitkovsky, A. 2005, in American Institute of Physics Conference Series, Vol. 801, Astrophysical Sources of High Energy Particles and Radiation, ed. T. Bulik, B. Rudak, & G. Madejski (AIP), 345–350, doi: 10.1063/1.2141897

Sridhar, N., Ripperda, B., Sironi, L., Davelaar, J., & Beloborodov, A. M. 2025, ApJ, 979, 199, doi: 10.3847/1538-4357/ada385

Sridhar, N., Sironi, L., & Beloborodov, A. M. 2021, MNRAS, 507, 5625, doi: 10.1093/mnras/stab2534

Takamoto, M., & Lazarian, A. 2016, ApJL, 831, L11, doi: 10.3847/2041-8205/831/2/L11

Tamburini, M., Pegoraro, F., Di Piazza, A., Keitel, C. H., & Macchi, A. 2010, New Journal of Physics, 12, 123005, doi: 10.1088/1367-2630/12/12/123005

Uzdensky, D. A. 2016, in Astrophysics and Space Science Library, Vol. 427, Magnetic Reconnection: Concepts and Applications, ed. W. Gonzalez & E. Parker, 473, doi: 10.1007/978-3-319-26432-5_12

Vranic, M., Martins, J. L., Fonseca, R. A., & Silva, L. O. 2016, Computer Physics Communications, 204, 141, doi: 10.1016/j.cpc.2016.04.002

Werner, G. R., Uzdensky, D. A., Cerutti, B., Nalewajko, K., & Begelman, M. C. 2016, ApJL, 816, L8, doi: 10.3847/2041-8205/816/1/L8

Yuan, F., & Zdziarski, A. A. 2004, MNRAS, 354, 953, doi: 10.1111/j.1365-2966.2004.08262.x

Zdziarski, A. A., & Gierliński, M. 2004, Progress of Theoretical Physics Supplement, 155, 99, doi: 10.1143/PTPS.155.99

Zenitani, S. 2015, Physics of Plasmas, 22, 042116, doi: 10.1063/1.4919383

Zhang, L., Stone, J. M., Davis, S. W., et al. 2026, arXiv e-prints, arXiv:2603.05588, doi: 10.48550/arXiv.2603.05588

Zhdankin, V., Uzdensky, D. A., Perez, J. C., & Boldyrev, S. 2013, ApJ, 771, 124, doi: 10.1088/0004-637X/771/2/124

Zou, M., Hakobyan, H., Mbarek, R., et al. 2025, PhRvD, 111, 083032, doi: 10.1103/PhysRevD.111.083032