



Editorial

Gender differences in athlete ECGs



In the first of this month's featured articles Bessem, de Bruijn and Nieuwland discuss the gender differences that occur in young athlete's ECG's and examine possible implications of these differences. Mucha and colleagues report a systematic review of the literature in the area of hip adductor strength and injury in distance runners and Adams and Peiffer describe a randomised trial that shows now effect of nasal dilation on a cycling 20 km time trial.

The second article in the sports injury section, by Fowler's group describes the effect of west to east long haul flights on players participating in the 2014 FIFA World Cup in Brazil confirming potential significant impact of "jet lag" on player wellness scores. Philipou and co-workers examine the relationship between nutrition knowledge and adherence to the Mediterranean Diet in young competitive swimmers. The final paper in the section, by Hindieh and colleagues, reviews current evidence and national guidelines for exercise in hypertrophic cardiomyopathy patients.

The first paper in the sports injury section, from Mosler and colleagues, adds to the literature of normative values in football for hip strength and range of motion with data from a professional football league. Howard and co-workers add support to the use of humeral elevation as a component of dynamic shoulder rehabilitation. Wikstrom and McKeon provide a report that suggests increased focus on ankle sensory rehabilitation in individuals with chronic ankle instability increases postural control following rehabilitation. In the final article, Pazzinatto's group report on a study examining the relationship between running volume and pressure pain thresholds.

In the first article in this month's physical activity section Rowlands and co-workers describe an algorithm that enables effective accelerometer wear site discrimination between the hip and wrist.

Cadenas-Sanchez reports a component of the European HELENA study examining the relationship between fitness and fatness and attention capacity in adolescents. Cairney and colleagues report a longitudinal study of a cohort of 2278 Canadian children examining the relationship between cardiorespiratory fitness and activity in children both with and without motor coordination problems. Reed's group report a negative effect of wearing compression garments in individuals with cardiovascular disease risk factors.

In the sport and exercise science section, Beckman, Connick and Tweedy review the literature surrounding strength assessment approaches for classification in Paralympic sport. Claudino and co-workers provide a meta-analysis of the literature examining the countermovement jump as a measure of neuromuscular status, suggesting average CMJ height is the best index of performance. De Andrade Kratz and colleagues report on a randomised controlled trial of beta-alanine supplementation and performance measures in judo athletes and in the final article in the section Fontana's group suggest a mix of variables that may be of use in predicting future performance in young rugby players.

I commend to you this issue which continues to reflect the Journal's readership players.

I commend to you this issue which continues to reflect the Journal's readership with an excellent mix of articles across the spectrum of science and medicine in sport.

Gordon S. Waddington, PhD
Chief in Editor