

# Discrete Structures



## Lecture 6: More on predicate calculus

Susan L. Epstein



1

## Last time

★ Predicate calculus extends logical representation

★ Validity  $\neq$  truth

- Invalid argument forms
- How to express facts in predicate calculus
- How to translate both ways between English and predicate calculus
- How to negate quantified statements
- How to demonstrate uniqueness

Fall 2023

CSCI 150

2/22

2

## Today's outline

- Multiple quantifiers in predicate calculus
- Argumentation in predicate calculus
- Formal verbal proofs



Predicate calculus extends logical representation



Fall 2023

CSCI 150

3/22

3

## Multiple quantifiers

For  $x$ ,  $D = \{x | x \text{ is a person}\}$ , for  $y$ ,  $E = \{y | y \text{ is a dog}\}$   
 $P(x, y)$  means "x loves y"

- $\forall x \in D, \forall y \in E P(x, y) \equiv \forall x, y P(x, y)$  Everyone loves all dogs  
 To prove this, test every possible pair of person and dog
- $\exists x \in D, \exists y \in E P(x, y) \equiv \exists x \in D, y \in E P(x, y)$  Someone loves some dog  
 To prove this, test every possible pair of person and dog
- $\forall x \in D, \exists y \in E P(x, y)$  Everyone loves some dog  
 To prove this, pick an arbitrary  $x \in D$  and find its  $y$
- $\exists x \in D \exists \forall y \in E P(x, y)$  Someone loves all dogs  
 To prove this, pick an arbitrary  $x \in D$  and show it works for every  $y$
- $\forall x \in D, \exists! y \in E P(x, y)$  Everyone loves exactly one dog
- $\exists! x \in D \exists \forall y \in E P(x, y)$  There is only one person who loves all dogs

Fall 2023

CSCI 150

4/22

4

## Negation of multiple quantifiers

- Recall that  $\sim[\forall x \in D, P(x)] \equiv \exists x \in D \ni \sim P(x)$   
 $\sim[\exists x \in D \ni P(x)] \equiv \forall x \in D \ni \sim P(x)$
- $\sim(\forall x \in D, \exists y \in E \ni P(x, y))$  For  $x$ ,  $D = \{x | x \text{ is a person}\}$ ,  
 $\equiv \exists x \in D \ni \sim(\exists y \in E \ni P(x, y))$  why? for  $y$ ,  $E = \{y | y \text{ is a dog}\}$   
 $\equiv \exists x \in D \ni \forall y \in E \ni \sim P(x, y)$  why?  $P(x, y)$  means "x loves y"  
 It is false that everyone loves some dog  $\equiv$  Someone loves no dogs
- $\sim(\exists x \in D, \forall y \in E \ni P(x, y))$   
 $\equiv \forall x \in D \ni \sim(\forall y \in E \ni P(x, y))$  why?  
 $\equiv \forall x \in D \ni \exists y \in E \ni \sim P(x, y)$  why?  
 It is false that someone loves all dogs  $\equiv$   
 Everyone has some dog they do not love

Fall 2023

CSCI 150

5/22

5

## What's the negation of ...?

$\forall x \in D \exists y \in E$  such that  $x + y = 1$   
 $\sim(\forall x \in D \exists y \in E \ni x + y = 1) \equiv$   
 $\exists x \in D \ni \sim \exists y \in E \ni x + y = 1 \equiv$   
 $\exists x \in D \ni \forall y \in E \ni x + y \neq 1$

$\exists x \in D$  such that  $\forall y \in E \ni x + y = -y$   
 $\sim(\exists x \in D \ni \forall y \in E \ni x + y = -y) \equiv$   
 $\forall x \in D \ni \sim(\forall y \in E \ni x + y = -y) \equiv$   
 $\forall x \in D \ni \exists y \in E \ni x + y \neq -y$



Any questions?

Fall 2023

CSCI 150

6/22

6

## Today's outline

- ✓ Multiple quantifiers in predicate calculus
- Argumentation in predicate calculus
- Formal verbal proofs

Fall 2023

CSCI 150

7/22

7

## Valid argument forms in predicate calculus

- Valid argument form in predicate calculus = for any predicate symbols used in the premises, when the resultant premises are all true the conclusion is true

 $A = \{x | x \text{ is alive}\}$ 
 $Lola \in A$ 
 $P(x) = x \text{ is a dog}$ 
 $Q(x) = x \text{ is a mammal}$ 

- Universal modus ponens

 $\forall x P(x) \rightarrow Q(x)$ 

All dogs are mammals

 $P(a)$  for a particular  $a \in A$ 

Lola is a dog

 $\therefore Q(a)$ 
 $\therefore$  Lola is a mammal


- Universal modus tollens

 $\forall x P(x) \rightarrow Q(x)$ 
Booby  $\in A$ 

All dogs are mammals

 $\sim Q(a)$  for a particular  $a \in A$ 

Booby is not a mammal

 $\therefore \sim P(a)$ 
 $\therefore$  Booby is not a dog


Fall 2023

CSCI 150

8/22

8

## Fallacies in predicate calculus

### Converse error

$\forall x P(x) \rightarrow Q(x)$   
 $Q(d)$  for a particular  $d \in D$   
 $\therefore P(d)$

### Inverse error

$\forall x P(x) \rightarrow Q(x)$   
 $\sim P(d)$  for a particular  $d \in D$   
 $\therefore \sim Q(d)$

$D = \{x \mid x \text{ is alive}\}$   
 $P(x) = \text{is a dog}$   
 $Q(x) = \text{is a mammal}$   
 $Sally \in D$



All dogs are mammals  
 Sally is a mammal  
 $\therefore$  Sally is a dog

All dogs are mammals  
 Sally is not a dog  
 $\therefore$  Sally is not a mammal

Fall 2023

CSCI 150

9/22

9

## Predicate calculus and mathematical statements

Not proving these, just getting ready to...

- The sum of 2 positive integers is always positive  
 $\forall x, y (x > 0 \wedge y > 0) \rightarrow (x + y) > 0$   
 $\forall x, y \in \mathbf{Z}^+ (x + y) > 0$
- Every real number except 0 has a multiplicative inverse  
 $\forall x \in \mathbf{R} x \neq 0 \exists y \in \mathbf{R} \exists xy = 1$
- There is no smallest positive real number  
 $\forall x \in \mathbf{R}^+ \exists y \in \mathbf{R}^+ \exists y < x$

Fall 2023

CSCI 150

10/22

10

## There's so much more

- Predicate calculus can be extended
  - Add operators for numbers  $=, \neq$
  - Add operators for sets  $\cup, \cap$
  - Quantify not just over variables but also over predicates
- Proofs can introduce variables dependent on others for their existence
- **Prolog** is a programming language that, given the premises of a true theorem in a limited version of predicate calculus, **can prove the theorem**
- $\exists$  many other valid argument forms and many other ways to construct a proof ... we will use them on additional foundational mathematics



Any questions?

Fall 2023

CSCI 150

11/22

11

## Today's outline

- ✓ Multiple quantifiers in predicate calculus
- ✓ Argumentation in predicate calculus
- Formal verbal proofs

Fall 2023

CSCI 150

12/22

12

## Motivation

- A proof is an argument that a statement is true
- Thus far, proofs have been by
  - Truth table
  - Substitution of logical equivalents
  - Substitution into valid argument forms
- Now we look at formal arguments directed at a variety of targets
- **Proof formats** are intended to help the reader follow the argument

Fall 2023

CSCI 150

13/22

13

## What a good (verbal) proof looks like

**Theorem:** (copy the statement here)

**Proof:**

**Let/Assume/Suppose:** Name variables and state what they stand for  
                                   be general: any                   state any assumptions

**We must show that...**  
                                   multiple grammatically correct sentences

Give a reason for every assertion

**Clarify your logic:** connect statements with reasons   Thus   Then  
                                   Therefore   So   Hence   Consequently   It follows that

Display equations and inequalities centered and on separate lines

By definition of           By substitution           Because           Since

**QED** = *quod erat demonstrandum* = that which we intended to show  
 = □ = ■ (aka tombstone)

Fall 2023

CSCI 150

14/22

14

## Proof frame

**Theorem:** (copy the statement here)

**Proof:**

**Let/Assume/Suppose:** Name variables and state what they stand for  
 be general                      state any assumptions

**We must show that...**  
 multiple grammatically correct sentences

Clarify your logic with a reason for every assertion

Display equations and inequalities clearly

**QED**

Fall 2023

CSCI 150

15/22

15

## Definitions as raw material

- Often a proof relies on formal definitions

We will use these as an example:

$n \in \mathbb{Z}$  is **even** iff  $\exists k \in \mathbb{Z}$  such that  $n = 2k$       30      516

$n \in \mathbb{Z}$  is **odd** iff  $\exists k \in \mathbb{Z}$  such that  $n = 2k + 1$       899      -5

- Proofs often rely on properties of sets as well

We will use these as an example:

**Closure:** Set  $S$  is **closed** under operation  $\diamond$  iff  $\forall x, y \in S, x \diamond y \in S$   
 $\mathbb{Z}$  is closed under addition

The square of any negative number is positive

Fall 2023

CSCI 150

16/22

16



## A correct, formal proof

**Theorem:** The sum of any 2 odd integers is even.

**Proof:**

Let  $x$  and  $y$  be any odd integers.

We must show that  $x + y$  is even.

By definition of odd number,  $\exists a, b \in \mathbb{Z}, x = 2a + 1$  and  $y = 2b + 1$ .

By substitution,  $x + y = 2a + 1 + 2b + 1 = 2a + 2b + 2$   
which factors into  
 $2(a + b + 1)$ .

Because  $\mathbb{Z}$  is closed under addition,  $(a + b + 1) \in \mathbb{Z}$ .

By definition of even number,  $2(a + b + 1)$  is even.

And since  $x + y = 2(a + b + 1)$ ,  $x + y$  is even.

**QED**

Fall 2023

CSCI 150

17/22

17

## These examples are incorrect!

- Do not argue for a theorem about an infinite set with examples.  
 $13$  is odd and  $19$  is odd and their sum  $32$  is even. So...
- Do not reuse the same variable name for 2 different values.  
 $\exists k \in \mathbb{Z}, x = 2k + 1$  and  $y = 2k + 1$ .
- Do not skip steps.  
By substitution  $x + y = 2(a + b + 1)$
- Do not use circular reasoning.  
Suppose  $x$  and  $y$  are any odd integers. When any odd integers are added, their sum is even. Hence  $x + y$  is even.
- Do not incorporate the conclusion with a variable.  
To show  $x + y$  is even we must show  $\exists k \in \mathbb{Z}, x + y = 2k$
- "any"  $\neq$  "some" Any is like  $\forall$ ; suggests a particular element  
By definition of odd,  $m = 2a + 1$  for any integer  $a$ .
- Do not use "if" when you mean "because."  
Suppose  $p$  is a prime number. If  $p$  is prime, then  $p$  cannot be written as a product of two smaller positive integers.

Fall 2023

CSCI 150

18/22

18

## Proof by cases skeleton

**Theorem:** If  $A_1$  or  $A_2$  or ... or  $A_n$  then  $C$ .

**Proof:**

**Let/Assume/Suppose:** Name variables and state what they stand for  
be general: **any**                      **state any assumptions**

**We must show that  $C$  is true in each of the following cases:**

**Case 1:** If  $A_1$  then  $C$ .

**Case 2:** If  $A_2$  then  $C$ .

...

**Case  $n$ :** If  $A_n$  then  $C$

Clarify your logic with a reason for every assertion  
Display equations and inequalities clearly

Thus  $C$  is true regardless of which is the case.

**QED**

Fall 2023

CSCI 150

19/22

19

## A simple proof by cases

**Theorem:** If  $n \in \mathbb{Z}$  then  $n^2 \geq n$ .

**Proof:**

**Let**  $n \in \mathbb{Z}$ .

**We must show that  $n^2 \geq n$  is true in each of the following cases:**

$n = 0$ ,  $n \geq 1$ , or  $n \leq -1$ .

**Case 1:**  $n = 0$

$0^2 = 0 \geq 0$  so true for  $n = 0$

**Case 2:**  $n \geq 1$

Multiplying both sides of  $n \geq 1$  by  $n$ :  $n^2 \geq n$  so true for  $n \geq 1$

**Case 3:**  $n \leq -1$

Since the square of any negative number is positive,  $n^2 > 0$  and

$0 > n$ , so  $n^2 \geq n$  and true for  $n \leq -1$

Thus  $n^2 \geq n$  is true regardless of which is the case.

**QED**

Fall 2023

CSCI 150

20/22

20

## Proof methods (so far)

Truth table

Sequence of statements with reasons

Valid argument forms (modus ponens, modus tollens,...)

Method of exhaustion

Predicate logic (quantification, existence, uniqueness)

Proof by cases

Fall 2023

CSCI 150

21/22

21

## What you should know

### ★ Predicate calculus extends logical representation

- How to translate multiply quantified statements from logic to English and from English to logic
- How to construct and negate multiply quantified predicate statements
- Valid arguments in FOPC
- Basic formal proof structures
- Proof by cases



Any questions?

**Next up: Introduction to number theory**

**Time to finish up that Opening sheet!**

**Problem set 5,6 is due on Thursday, September 21 at 11PM**

Fall 2023

CSCI 150

22/22

22