

Testing and Isolation Efficacy; Insights from a Simple Epidemic Model

January 27, 2021

1 Abstract

2 Introduction

The observed dynamics of the COVID-19 epidemic are driven by both epidemiological processes (infection and recovery) and testing processes (testing and test reporting). In addition to shaping epidemic observations (via case reports), testing processes can also affect epidemiological dynamics. In particular, individuals who are awaiting the results of a test may partially or fully self-isolate, and individuals who have tested positive are highly likely to self-isolate. We developed a model that incorporates mechanistic epidemic processes and testing in order to explore the effects of testing and isolation on epidemic dynamics.

Because individuals who have been tested or received a positive test are likely to self-isolate, thus reducing or eliminating their potentially infectious contacts, the dynamics of an epidemic will depend on who gets tested and why. For a given testing intensity (tests performed per day), testing will have a stronger effect on infection rates the more strongly it is focused on people who are infectious for SARS-CoV-2. This level of focus depends in turn on the purpose and design of testing programs. (1) When testing is done for the purposes of disease surveillance (Foddai et al., 2020) we might expect tests to be assigned randomly across the population (in an ideal world: some bias in selection for testing is probably inevitable), although a stratified sampling design would be preferred for statistical efficiency (Cleevely et al., 2020). *[JD: This is a bit confusing; cleevely is specifically about control, and not just surveillance.]* Over the course of the COVID19 pandemic, however, the vast majority of testing has been done with other goals – primarily diagnosis (determining

COVID19 infection status for clinical purposes), or control (determining COVID19 status in order to quickly isolate cases that have been found by contact tracing), which we characterize as *targeted* testing strategies. In these cases, testing probabilities vary widely across epidemiological compartments; in our dynamical model, we will characterize these probabilities by assigning a *per capita* testing weight to each compartment that determines the *relative* probability that an individual in that compartment will be selected for testing (see Methods).

[JD: *note that we have not yet introduced the term “testing weights”.*] When testing is used primarily for diagnosis it will focus on people with COVID19-like symptoms; thus the testing weights for people with vs. without SARS-CoV-2 infections will depend on the relative magnitudes of the probability of having COVID19-like symptoms when infected with SARS-CoV-2 (i.e. the relative proportions of presymptomatic or asymptomatic vs. symptomatic infections) and the probability of having COVID19-like symptoms when *not* infected (e.g. due to other respiratory illnesses). Testing for epidemic control will focus on people who are known to have been in contact with known COVID19 cases; in this case the testing weights for infected vs. uninfected people will depend on the probability of infection given contact, as well as the thoroughness and effectiveness of the system for identifying suspicious contacts.

[BMB: *save stuff about flights, surgery, LTC for Discussion?*]

In addition to the overall testing intensity, the testing weights for different compartments, and the degree of self-isolation of test-pending and test-confirmed people, our models will also need to include test specificity and sensitivity, as well as the distribution of times from testing to reporting. [JD: *This seems weak. Why are we studying \mathcal{R}_0 ? What does it cover and what not?*] Here, we focus on the the effect of testing intensity, different levels of testing “focus” (from random to highly targeted), and rate of test processing on a single index of epidemiological dynamics, the basic reproduction number \mathcal{R}_0 .

The early dynamics of an epidemic are determined by \mathcal{R}_0 . This index is defined as the expected number of secondary infections arising from a typical infective individual in a completely susceptible population (Dietz, 1993). In the early stages of an epidemic the number of infected individuals is expected grow exponentially over time when $\mathcal{R}_0 > 1$. To control the spread of disease in the population, \mathcal{R}_0 must be < 1 . Although the value of \mathcal{R}_0 cannot completely characterize the dynamics of even the simplest epidemic model (Shaw and Kennedy, 2021), it does give a simple and widely accepted index for the difficulty of control, as well as some indication of the likely final size of an epidemic (Ma and Earn, 2006).

[Fady: *Not sure if interesting:*

one point we can make when discussing the relative weight of testing in different compartments is to discuss pre-testing screening tools, such as surveys or questionnaires. If we have a quantitative description of how the testing intensities affect the dynamics, we can make statements like “our results suggest that employing a pre-testing screening tool can help target infected individuals more effectively. In particular, doubling the sensitivity of the pre-screening tool would *do something* to \mathcal{R}_0 .] [BMB: possibly interesting, but **belongs in the Discussion** if anywhere.]

3 Methods

We developed a deterministic model, Eqs. (S1), which groups individuals based on disease status and testing status. Disease states include Susceptible, Infectious and Recovered (thus this is an SIR-type model), and testing status categorizes people as *untested*, waiting-for-*positive*, waiting-for-*negative*, or *confirmed positive* (Figure 1). Two ‘accumulator’ compartments, N and P , were also incorporated in the model in order to collect cumulative reported negative or positive tests. The model and details of calculation of \mathcal{R}_0 are presented in the Supplementary Materials.




Figure 1: Flowchart of the SIR (Susceptible-Infectious-Recovered) model, S1. For the description of the parameters see Table 1.

Table 1 defines the model parameters, which are generally straightforward *per capita* flows between compartments, or modifiers to these flow rates. The novel component of the model comes in through the compartment-specific relative testing weights W_S , W_I and W_R ; these give the relative rates at which people in the S , I ,

and R compartments are tested, respectively. For example, $W_I/W_S = 2$ means that infected individuals are tested at twice the *per capita* rate of susceptible individuals.

In order to link to more applied models, we constructed this model so we could specify the total *per capita* testing rate. We do this by defining the weighted size of the testing pool $W = W_S S_u + W_I I_u + W_R R_u$, and calculating the scaling parameter for testing as:

$$\sigma = \frac{\rho N_0}{W}. \quad (1)$$

Thus, *per capita* testing rate for compartment Z is $F_Z = \sigma W_Z$, where $Z \in \{S, I, R\}$. For a high-sensitivity test, infected people typically flow through to the “confirmed positive” (I_c/R_c) compartments and are thus unavailable for further testing. Over the course of the epidemic, a fixed testing rate as specified in (1) can (if large enough) exhaust the pool of people available for testing, leading to a singularity when no one is left untested. Although this phenomenon does not affect our analysis of \mathcal{R}_0 , it does sometimes occur in numerical solutions of the time dynamics: we discuss an adjustment to the model that solves this problem in the Supplementary material.

The classical SIR model is based on the following implicit assumptions; well-mixed population, homogeneity of the population, exponentially distributed duration of infection and large population size (see, e.g., Keeling and Rohani (2011)). In addition to these standard assumptions, our model, S1, assumes: (i) there is a single force of infection, Λ , defined as follows

$$\Lambda = \beta \frac{(I_u + \eta_w I_n + \eta_w I_p + \eta_c I_c)}{N_0}, \quad (2)$$

across all susceptible pools (see Table 1 for detail), (ii) $\eta_c \leq \eta_w$, i.e., the individuals awaiting test results have a higher transmission probability than the reported individuals. For this analysis, we also assume a perfectly specific test, ($p_S = 0$). This last assumption combined with the assumption that no individual is in waiting-for-positive or confirmed positive compartments, i.e., $S_p(0) = S_c(0) = 0$, reduces the model to 10 equations (equations c and d in (S1) are eliminated).

The Disease-Free Equilibrium (DFE) for the SIR model, Eqs. (S1), is given by solving the coupled system including $S_u + S_n = N_0$ and $F_S S_u - \omega S_n = 0$. The DFE is

$$S_n^* = \frac{\rho}{\omega} N_0, \quad S_u^* = N_0 - S_n^*, \text{ and } I_j = R_j = 0 \text{ for all } j. \quad (3)$$

The basic reproduction number, \mathcal{R}_0 , was calculated by using the next generation

Symbol	Description	Unit	Value
N_0	Total population size	people	10^6
ω	Rate of onward flow from “waiting” to “confirmed” or “untested” compartments	1/day	-
γ	Recovery rate	1/day	1/3
ρ	Per capita testing intensity	1/day	0.01
η_w	Relative probability of transmission for “waiting” individuals	-	-
η_c	Relative probability of transmission for “confirmed positive” individuals	-	-
β	Transmission rate	1/day	0.39
Λ	Force of infection	1/day	-
p_S	Probability of positive tests for S ($= 1 - \text{specificity}$)	-	0
p_I	Probability of positive tests for I ($= \text{sensitivity}$)	-	1
p_R	Probability of positive tests for R ($= 1 - \text{specificity}$)	-	0.5
W_S, W_I, W_R	Relative testing weights	-	Random: $\{1, 1, 1\}$ Targeted: $\{0.3, 1, 1\}$

Table 1: Parameters of the model (S1).

matrix method developed by Van den Driessche and Watmough (2002). \mathcal{R}_0 is

$$\mathcal{R}_0 = (A \times S_u^* + B \times S_n^*) \times C, \quad (4)$$

where

$$\begin{aligned} A &= \gamma(\omega + \gamma) + (\gamma\eta_w + \omega\eta_c p_I)F_I, \\ B &= (\omega + (F_I + \gamma)\eta_w)\gamma + \frac{(\eta_w\gamma + \eta_c\omega)\omega p_I F_I}{\omega + \gamma}, \\ C &= \frac{\beta/\gamma}{N_0(\gamma(\omega + \gamma) + F_I(\gamma + \omega p_I))}. \end{aligned}$$

Further details are provided in the Supplementary material section.

The analytical calculation of the next generation matrix, G , and the derivation of \mathcal{R}_0 was carried out in Maple (ref?) by using simple linear Algebra package. We used R for the simulation part which included using the explicit expression of \mathcal{R}_0 4 as a function of the underlying parameters, specifying the parameters a realistic range, calculating the corresponding \mathcal{R}_0 and plotting the contours of \mathcal{R}_0 by using ggplot package in R. The related result is presented in Figure 2, where panel (2a) represents the random testing, and panel (2b) is representing non-random testing. The parameter values or ranges are as follows (I like using a table here, with the param, their units and values). It is notable that in the simulation part, the explicit expression of \mathcal{R}_0 was used and not the approximation (S6). This simulation reflects the behavior of \mathcal{R}_0 with respect to the selected parameters for two different testing strategies: (i) random testing, represented by all testing weights to be the same, $W_S = W_I = W_R$, and (ii) non-random testing, when testing weight are not equal. For simulation purposes we chose $W_S = W_I = W_R = 1$ for random testing, and $W_S = 0.3$ and $W_I = W_R = 1$ for non-random testing strategy. Note that the critical contour of $\mathcal{R}_0 = 1$ is plotted in solid line in Figure 2.

4 Results

The explicit formula for the basic reproduction number, \mathcal{R}_0 (4), provides an opportunity to study the influence of changes in the underlying parameters on the critical index of epidemic dynamics. We are interested in understanding the effect of parameters that can be realistically controlled by changing in isolation, testing intensity and test resulting, i.e., η_c and η_w , ρ and ω , respectively. The following Proposition is the direct result of taking partial derivative of \mathcal{R}_0 (4) with respect to selected parameters.

Proposition 1. Using the expression of \mathcal{R}_0 (4),

1. $\partial R_0/\partial \eta_c \geq 0$ and $\partial R_0/\partial \eta_w \geq 0$.
2. $\partial \mathcal{R}_0/\partial \rho \leq 0$ when $\rho \approx 0$.
3. $\partial R_0/\partial \omega$ can be positive or negative when $\rho \approx 0$.

Given that the perfect isolation occurs at $\eta_c = \eta_w = 0$, Prop. 1 means that lifting the isolation from awaiting group results in an increased \mathcal{R}_0 and consequently a greater number of infected individuals. We used Taylor approximation of \mathcal{R}_0 at $\rho = 0$ due to the complexity of the expression and thus inconclusive. See the Appendix for details. It is straightforward to derive 2 and 3 from the linear approximation of \mathcal{R}_0 at $\rho = 0$.

Prop. 2 indicates that increasing testing intensity reduces \mathcal{R}_0 . This is sensible since as people are moved to test compartments (I_p, I_n **Fady: the susceptibles also reduce contact and transmission**) the higher probability of being subject to isolation and the lower the transmission (2) will become.

[BMB: move to discussion?? this applies to this ¶, also most of what's below here. To a first approximation, any text that discusses the context and meaning of the results, rather than the results themselves, can go in the discussion.]

Of these results, prop. 3 is the most surprising. It states that returning test results more rapidly (i.e., increasing ω) does not necessarily lower \mathcal{R}_0 ; whether increasing ω lowers \mathcal{R}_0 depends on the precise combination of model parameters (test reporting rate, testing processes/strategies (W_*), testing efficacy (P_I) and isolation (η_*). Specifically, the $\partial R_0/\partial \omega > 0$ suggests that reproduction number may increase as the test reporting process become faster. This can be seen from expression (S7) in the case of perfect isolation, when $\eta_w = 0$ and $\eta_c = 0$. The underlying mechanism for this result is that individuals may take more precautionary behaviors (e.g., physical distancing) when they are awaiting test results compared to when they are untested or their test has returned negative. An example of a situation that would favour slow return of tests is if the test being employed produces many false negatives, because many infected individuals will believe they are negative and thus may unknowingly spread the virus to many others. For an example of the behaviour of \mathcal{R}_0 see Figure 2 at varying values of ω . This property also provides a tool to quantify the amount of delay required in the test reporting process as a strategy to reduce \mathcal{R}_0 and consequently control the epidemic.

Our numerical solutions in support of the analytical results in Proposition 1 are presented in Figure 2. It can be inferred that when random testing strategy is applied, Figure 2a, comparing to the non-random testing strategy, Figure 2a,




Figure 2: Behaviour of the basic reproduction number, \mathcal{R}_0 , with respect to changes in the underlying parameters; **[BMB: remind the reader about the parameters and their values]**. The solid line in each panel represents the critical contour of $\mathcal{R}_0 = 1$. **[BMB: we still have some collisions in the y-axis tick labels]**

the critical testing intensity of the corresponding panels are lower in non-random testing. Also, it appears that speeding the test reporting, i.e., increasing ω , does not significantly lower \mathcal{R}_0 when awaiting people follow the isolation perfectly, i.e., η_w is closer to 0. However, speeding the test reporting reduces the epidemic more when the awaiting people less follow the isolation, i.e., η_w is closer to 1.

Furthermore, we derived an inequality that quantifies the exact relationships between model parameters that result in returning tests more rapidly being favorable (S13). To give a biological interpretation, we describe the qualitative trends predicted by the inequality. Returning test results more rapidly tends to be more favorable when

- individuals who have tested positive lower their contact much more than individuals who are waiting for their test results (i.e., $\eta_c \ll \eta_w$).
- the test is sensitive.
- testing of infected individuals is more intense than testing of susceptible individuals.

An in-depth discussion of these results is presented in the discussion section.

The use of tests cheaper than RT-PCR has been proposed as a potential strategy for containing the COVID-19 pandemic. While cheaper tests may be less sensitive and reliable than RT-PCR, they allow for broader and more intense testing. Using our Taylor approximation of \mathcal{R}_0 near $\rho = 0$, we examined what circumstances (i.e., model parameters) make the use of one test more favourable than another, and give a complete description through inequality S14. In general, we found that the expensive test tends to more effectively lower \mathcal{R}_0 when

- individuals who test positive self-isolate much more than individuals who are waiting for their test result.
- the time it takes to return tests is much shorter than the mean infectious period.
- the testing intensity is much greater for infected individuals than susceptible individuals.

Our model also predicts an asymmetry between cheap and expensive tests: if a cheap test can identify on average more infected individuals as an expensive test, then our model predicts that the cheap test will lower \mathcal{R}_0 more. In contrast, if an expensive test can identify on average more infected individuals, it will not necessarily lower \mathcal{R}_0 more than the cheap test. *[Fady: we can state this as a formal proposition in equations rather than in words, if that would be better.]*

5 Discussion

From JD’s email:

- We’re basically looking at the potential advantage of slow tests: people waiting for test results may be more careful than those who have received negatives. These advantages are real, and neglected. We also compare to an individual-level advantage of fast tests: people who test positive may be even more careful. But we are missing out on community-level advantages of fast testing: better assessment of the situation, identification of hot spots, contact-tracing, etc.
- Maybe something about the testing rate/intensity, now the per capita testing intensity is very low ($\rho \approx 0$). In near future new test kits may be widely accessible, our model provide insights in this case (what are those?)
- “delay negative results” result

5.1 Rate of returning tests

Our model predicts that when the difference in self-isolation between individuals waiting for their tests and individuals who are confirmed positive is small, returning tests rapidly is not beneficial. The underlying mechanism for this prediction is that the benefit in returning tests rapidly so (i.e., the extra self-isolation an individual does when they test positive) is minimal and outweighed by the fact that those who test negative may increase their contacts. Thus, in such a case, returning tests more rapidly is expected to increase \mathcal{R}_0 . Similarly, if the test is not sensitive, many infected individuals will test negative, and thus may increase their contacts under the false belief that they are not infectious, so returning tests rapidly in this case may also contribute to increased transmission.

When testing of infected individuals is more intense than susceptible individuals, returning test results rapidly is beneficial because those infected individuals will be informed and self-isolate. Conversely, if testing of susceptible individuals is much greater, very few tests will be positive, and the benefit of informing those few positive cases will be outweighed by the potential increase in contacts of the individuals testing negative.

5.2 Expensive vs. cheap tests

[Fady: *I think we should discuss this before I write it. I'm a little hesitant to really expand on these results because of the points JD brought up over email.*]

References

- Arino, J. and Portet, S. (2020). A simple model for COVID-19. *Infectious Disease Modelling*.
- Bergstrom, T., Bergstrom, C. T., and Li, H. (2020). Frequency and accuracy of proactive testing for COVID-19. *medRxiv*.
- Cleevely, M., Susskind, D., Vines, D., Vines, L., and Wills, S. (2020). A workable strategy for COVID-19 testing: stratified periodic testing rather than universal random testing. *Oxford Review of Economic Policy*, 36(Supplement_1):S14–S37.
- de Celles, M. D., Casalegno, J.-S., Lina, B., and Opatowski, L. (2020). Influenza may facilitate the spread of SARS-CoV-2. *medRxiv*.

- Dietz, K. (1993). The estimation of the basic reproduction number for infectious diseases. *Statistical methods in medical research*, 2(1):23–41.
- Elbanna, A., Wong, G. N., Weiner, Z. J., Wang, T., Zhang, H., Liu, Z., Tkachenko, A. V., Maslov, S., and Goldenfeld, N. (2020). Entry screening and multi-layer mitigation of COVID-19 cases for a safe university reopening. *medRxiv*.
- Endo, A., Leclerc, Q. J., Knight, G. M., Medley, G. F., Atkins, K. E., Funk, S., Kucharski, A. J., et al. (2020). Implication of backward contact tracing in the presence of overdispersed transmission in COVID-19 outbreaks. *Wellcome Open Research*, 5(239):239.
- Foddai, A., Lubroth, J., and Ellis-Iversen, J. (2020). Base protocol for real time active random surveillance of coronavirus disease (COVID-19)—adapting veterinary methodology to public health. *One Health*, page 100129.
- Jenness, S. M., Willebrand, K. S., Malik, A. A., Lopman, B. A., and Omer, S. B. (2020). Modeling dynamic network strategies for SARS-CoV-2 control on a cruise ship. *medRxiv*.
- Keeling, M. J. and Rohani, P. (2011). *Modeling infectious diseases in humans and animals*. Princeton University Press.
- Larremore, D. B., Wilder, B., Lester, E., Shehata, S., Burke, J. M., Hay, J. A., Tambe, M., Mina, M. J., and Parker, R. (2020). Test sensitivity is secondary to frequency and turnaround time for COVID-19 screening. *Science Advances*, page eabd5393.
- Lopman, B., Liu, C., Le Guillou, A., Handel, A., Lash, T. L., Isakov, A., and Jenness, S. (2020). A model of COVID-19 transmission and control on university campuses. *medRxiv*.
- Ma, J. and Earn, D. J. D. (2006). Generality of the final size formula for an epidemic of a newly invading infectious disease. *Bulletin of Mathematical Biology*, 68(3):679–702.
- Rice, K., Wynne, B., Martin, V., and Ackland, G. J. (2020). Effect of school closures on mortality from coronavirus disease 2019: old and new predictions. *BMJ*, 371.
- Ruszkiewicz, D. M., Sanders, D., O’Brien, R., Hempel, F., Reed, M. J., Riepe, A. C., Bailie, K., Brodrick, E., Darnley, K., Ellerkmann, R., et al. (2020). Diagnosis of

- COVID-19 by analysis of breath with gas chromatography-ion mobility spectrometry: a feasibility study. *EClinicalMedicine*, page 100609.
- Shaw, C. L. and Kennedy, D. A. (2021). What the reproductive number R_0 can and cannot tell us about COVID-19 dynamics. *Theoretical Population Biology*.
- Tuite, A. R., Fisman, D. N., and Greer, A. L. (2020). Mathematical modelling of COVID-19 transmission and mitigation strategies in the population of Ontario, Canada. *CMAJ*, 192(19):E497–E505.
- Van den Driessche, P. and Watmough, J. (2002). Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical biosciences*, 180(1-2):29–48.

Supplemental Materials: Title for main text

5.3 Model and calculation of \mathcal{R}_0

The model is

$$dS_u/dt = -\Lambda S_u - F_S S_u + \omega S_n, \quad (\text{S1a})$$

$$dS_n/dt = -\Lambda S_n + (1 - p_S) F_S S_u - \omega S_n, \quad (\text{S1b})$$

$$dS_p/dt = -\Lambda S_p + p_S F_S S_u - \omega S_p, \quad (\text{S1c})$$

$$dS_c/dt = -\Lambda S_c + \omega S_p, \quad (\text{S1d})$$

$$dI_u/dt = \Lambda S_u - F_I I_u + \omega I_n - \gamma I_u, \quad (\text{S1e})$$

$$dI_n/dt = \Lambda S_n + (1 - p_I) F_I I_u - \omega I_n - \gamma I_n, \quad (\text{S1f})$$

$$dI_p/dt = \Lambda S_p + p_I F_I I_u - \omega I_p - \gamma I_p, \quad (\text{S1g})$$

$$dI_c/dt = \Lambda S_c + \omega I_p - \gamma I_c, \quad (\text{S1h})$$

$$dR_u/dt = \gamma I_u - F_R R_u + \omega R_n, \quad (\text{S1i})$$

$$dR_n/dt = \gamma I_n + (1 - p_R) F_R R_u - \omega R_n, \quad (\text{S1j})$$

$$dR_p/dt = \gamma I_p + p_R F_R R_u - \omega R_p, \quad (\text{S1k})$$

$$dR_c/dt = \gamma I_c + \omega R_p, \quad (\text{S1l})$$

$$dN/dt = \omega(S_n + I_n + R_n), \quad (\text{S1m})$$

$$dP/dt = \omega(I_p + R_p), \quad (\text{S1n})$$

where β , (in units of 1/day), is the disease transmission rate, η_w and η_c are the isolation parameters for awaiting and reported individuals, respectively. Λ is the force of infection defined in Eq. (2).

The next generation matrix is $G = FV^{-1}$, where matrix F represents the inflow of new infection to the infected compartments and matrix V represents the flow in the infected compartments when the population is totally susceptible. For model, ??-??, F and V are

$$F = \beta/N_0 \begin{bmatrix} S_u & \eta_w S_u & \eta_w S_u & \eta_c S_u \\ S_n & \eta_w S_n & \eta_w S_n & \eta_c S_n \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (\text{S2})$$

$$V = \begin{bmatrix} F_I + \gamma & -\omega & 0 & 0 \\ -(1-p_I)F_I & \omega + \gamma & 0 & 0 \\ -p_I F_I & 0 & \omega + \gamma & 0 \\ 0 & 0 & -\omega & \gamma \end{bmatrix}, \text{ thus} \quad (\text{S3})$$

$$V^{-1} = \begin{bmatrix} \frac{\omega + \gamma}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & \frac{\omega}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & 0 & 0 \\ \frac{(1-p_I)F_I}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & \frac{F_I + \gamma}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & 0 & 0 \\ \frac{p_I F_I}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & \frac{\omega p_I F_I}{(\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I)(\omega + \gamma)} & \frac{1}{\omega + \gamma} & 0 \\ \frac{\omega F_I p_I}{(\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I)\gamma} & \frac{\omega^2 F_I p_I}{(\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I)(\omega + \gamma)\gamma} & \frac{\omega}{(\omega + \gamma)\gamma} & \frac{1}{\gamma} \end{bmatrix}. \quad (\text{S4})$$

The particular form of F with two rows of zeros at the bottom, simplifies G as

$$G = \begin{bmatrix} G_{11} & G_{12} \\ 0 & 0 \end{bmatrix}, \text{ where } G_{11} = C \begin{bmatrix} A S_u & B S_u \\ A S_n & B S_n \end{bmatrix}. \quad (\text{S5})$$

The block matrix G_{12} does not influence \mathcal{R}_0 (defined as the spectral radius of G). All that matters here are the eigenvalues of G_{11} , which are 0 and \mathcal{R}_0 (4).

5.4 On Testing Rate and Numerical Singularity

[BMB: need a little bit of context here, saying that we didn't do any numerical solutions for the trajectories in our analysis, but if you do try them you run into this problem ...]

The testing rate, σ , should be formulated such that people from the untested compartments will not be tested if they are not there. The numerical singularity issue with the chosen σ (1) is that the population in S compartments appeared to blow up when the DFE is achieved. This is once the only untested people are susceptibles, the FOI will become $\Lambda = 0$, testing rate $F_s = \rho N_0 / S_u$. Thus, eq(1) of the model will be $dS_u/dt = -\rho N_0 + \omega S_n$ which is no longer dependent on S_u and a linear rate of leaving the S_u compartment.

One way to fix this issue, is to consider a maximum testing rate, τ (1/day). In general, we want to test at a rate of ρ across the whole population. This won't

always be possible, so we impose a maximum rate of τ per testable person and redefine $\sigma = \frac{\tau \rho N_0}{\tau W + \rho N_0}$, with the assumption that $\tau \gg \rho$. This alteration in σ , does not change any results related to \mathcal{R}_0 , thus we only impose it in the simulation of the epidemic dynamic.

5.5 Taylor Approximation of \mathcal{R}_0 at $\rho = 0$

The basic reproduction number, \mathcal{R}_0 , close to $\rho = 0$ can be approximated linearly in ρ by using Taylor approximation. it follows

$$R_0 \approx \beta/\gamma + \frac{\beta\rho}{\omega(\omega + \gamma)\gamma^2 W_S} \left(\gamma(\eta_w - 1)(\gamma W_S + \omega W_I) + (\eta_c - 1)P_i W_i \omega^2 \right) + \mathcal{O}(\rho^2). \quad (\text{S6})$$

Also, to analyze the influence of ω on \mathcal{R}_0 , the approximation (S6) was used. Then it is straight forward to have

$$\partial R_0 / \partial \omega = \frac{-\beta\rho}{\gamma W_S \omega^2 (\gamma + \omega)^2} (a\omega^2 + b\omega + c), \quad (\text{S7})$$

where $a = (\eta_w - 1)W_I - (\eta_c - 1)P_i W_I = ((s - p_I)\eta_c + (p_I - 1))W_I$, $b = 2(\eta_w - 1)\gamma W_S$ and $c = (\eta_w - 1)\gamma^2 W_S$. Given that $0 \leq \eta_c \leq \eta_w \leq 1$, one can easily derive $b \leq 0$ and $c \leq 0$.

Note that in general, the necessary and sufficient condition for $a \geq 0$ is $(s - p_I)\eta_c \geq (1 - p_I)$, where $s = \frac{\eta_w}{\eta_c} \geq 1$.

In the case of “perfect isolation”, i.e., when $\eta_w = 0$ and consequently $\eta_c = 0$, it is straightforward to see that $a \leq 0$, $b < 0$ and $c < 0$. Thus, $\partial \mathcal{R}_0 / \partial \omega \geq 0$. As an example, in the case of a very accurate testing regime, i.e., $P_I = 1$, $a \geq 0$ is achieved. If $a \geq 0$, the quadratic expression in (S7), has Real roots. Assuming that $\omega_1 < 0$ and $\omega_2 > 0$ be the roots of the quadratic expression in $\partial R_0 / \partial \omega$. Thus, $\partial R_0 / \partial \omega > 0$ for $0 < \omega < \omega_2$ and $\partial R_0 / \partial \omega < 0$ for $\omega > \omega_2$.

[Fady: this subsection and most of the next belong to the Appendix since they are very detailed. Would you clean up the writing, there are some repeated expressions that can be referenced?]

5.6 rate of returning tests

The linearization of \mathcal{R}_0 around $\rho = 0$ is

$$\mathcal{R}_0 \approx \beta/\gamma + \frac{\beta\rho}{\omega(\omega + \gamma)\gamma^2 W_S} \left(\gamma(\eta_w - 1)(\gamma W_S + \omega W_i) + (\eta_c - 1)P_i W_i \omega^2 \right). \quad (\text{S8})$$

So when $\rho \approx 0$ we have

$$\partial \mathcal{R}_0 / \partial \omega \approx \frac{-\beta \rho}{\gamma W_s \omega^2 (\gamma + \omega)^2} (a \omega^2 + b \omega + c).$$

where $a = (\eta_w - 1)W_i - (\eta_c - 1)P_i W_i$, $b = 2(\eta_w - 1)\gamma W_s$ and $c = (\eta_w - 1)\gamma^2 W_s$.

Perhaps counter-intuitively, the equation above does not predict that \mathcal{R}_0 is monotone decreasing with respect to ω . In other words; our model does not predict that returning test results more rapidly *always* lower \mathcal{R}_0 . In order to gain insight into this intriguing behavior, we examine the zeroes of $\frac{\partial \mathcal{R}_0}{\partial \omega}(\omega)$.

Defining the following quantity, Q , will help us write the roots of $\partial \mathcal{R}_0 / \partial \omega$ neatly.

$$Q = \frac{W_i}{W_s} \left(1 - \frac{n_t - 1}{n_w - 1} P_i \right) \quad (\text{S9})$$

$$(\text{S10})$$

With that in mind, we can write the roots of $\partial \mathcal{R}_0 / \partial \omega$ as

$$\omega_1 = \frac{\gamma}{-\sqrt{1-Q}-1} \quad (\text{S11})$$

$$\omega_2 = \frac{\gamma}{\sqrt{1-Q}-1} \quad (\text{S12})$$

Note that the zeroes are real if and only if $Q < 1$. Note that have $\eta_c < \eta_w$, so if $P_i \approx 1$, we will have $Q < 0 < 1$. Thus, if we assume near-perfect test sensitivity, ω_1 and ω_2 will be real.

Assuming ω_1, ω_2 are real, it is easy to confirm that $\omega_1 < 0$ by looking at the denominator. To see that $\omega_2 > 0$, recall that $Q < 0$, so $\sqrt{1-Q} > 1$ and so $\sqrt{1-Q} - 1 > 0$. Knowing that $\omega_1 < 0$, the only root of interest (i.e., biologically relevant quantity) is ω_2 .

We can prove that $\partial \mathcal{R}_0 / \partial \omega > 0$ when $\omega \in (0, \omega_2)$ and $\partial \mathcal{R}_0 / \partial \omega < 0$ when $\omega \in (\omega_2, \infty)$ by computing the limits of $\partial \mathcal{R}_0 / \partial \omega$ at 0 and ∞ respectively. So it follows that \mathcal{R}_0 has a global maximum with respect to ω at $\omega = \omega_2$.

Now we want to characterize the parameter regions on which $\partial \mathcal{R}_0 / \partial \omega < 0$ (i.e., the conditions under which returning test results more rapidly is favorable). By the previous analysis, this is equivalent to solving for $\omega > \omega_2$. So

$$\begin{aligned}
& \omega > \omega_2 \\
& \omega > \frac{\gamma}{\sqrt{1-Q}-1} \\
& \vdots \\
& \frac{1-n_t}{1-n_w} P_i > \frac{W_s}{W_i} \left(\frac{\gamma}{\omega} + 1 \right)^2
\end{aligned} \tag{S13}$$

5.7 Expensive vs. cheap tests

The use of tests cheaper than RT-PCR has been proposed as a potential strategy for containing the COVID-19 pandemic. While cheaper tests may be less sensitive and reliable than RT-PCR, they allow for broader and more intense testing. In the analysis below, we compare the \mathcal{R}_0 predicted by our model depending on the testing strategy.

Consider a test that allows us to test at rate ρ_1 and has sensitivity $P_{i,1}$, and another test that allows us to test at ρ_2 and has sensitivity $P_{i,2}$. Suppose that $\rho_1 > \rho_2$. Recall that the linearization of \mathcal{R}_0 around $\rho \approx 0$ is given by

$$\mathcal{R}_0 \approx \beta/\gamma + \frac{\beta\rho}{\omega(\omega+\gamma)\gamma^2 W_s} \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_i W_i \omega^2 \right).$$

Treating \mathcal{R}_0 as a function of ρ and P_i , we can reduce the inequality

$$\mathcal{R}_0(\rho_2, P_{i,2}) < \mathcal{R}_0(\rho_1, P_{i,1})$$

into

$$\rho_1 \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_{i,1} W_i \omega^2 \right) - \rho_2 \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_{i,2} W_i \omega^2 \right) > 0$$

\vdots

$$\frac{\rho_2 P_{i,2} - \rho_1 P_{i,1}}{\rho_1 - \rho_2} > \frac{1 - \eta_w}{1 - \eta_c} \cdot \frac{\gamma(\gamma W_s + \omega W_i)}{\omega^2 W_i} \tag{S14}$$

Note that the RHS is positive, thus a necessary condition for the inequality above to hold is that $\rho_2 P_{i,2} > \rho_1 P_{i,1}$, equivalently

$$\frac{P_{i,2}}{P_{i,1}} > \frac{\rho_1}{\rho_2}. \tag{S15}$$

To state an example of this, if test A is three times as expensive as test B (and hence one can test three times as many people with test B), using test A rather than B will be favorable only if test A is at least 3 times more sensitive than test B . Note that this is a necessary but not sufficient condition, so even if test A is three times more sensitive, it is still possible for test B to be more effective.

Eq. (S14) tells us precisely when a test corresponding to $\rho_2, P_{i,2}$ will yield a lower \mathcal{R}_0 than a test corresponding to $\rho_1, P_{i,1}$, where $\rho_1 > \rho_2$. Some of the qualitative *trends* that favor test 2 (the higher-sensitivity test) include

- individuals who test positive self-isolate much more than individuals who are waiting for their test result.
- the time it takes to return tests is much shorter than the mean infectious period.
- the testing intensity is much greater for infected individuals than susceptible individuals.

5.8 Literature Review

5.8.1 Explicit models of TTI (trace/test/isolate) based on network or agent-based models

(Endo et al., 2020) [**Ali:** *It seems to me that this is just a statistical model to estimate the parent-offspring of an infected index, not sure if it fits into agent-based group!*] Used simulation on a branching process model to assess the forward and backward contact tracing efficiency. Assuming a negative-binomial branching process with a mean R , reproduction number, and overdispersion parameter k , the mean total number of generation G_3 and averted G_3 are estimated. The effectiveness of TTI is defined as the ratio of averted to the mean.

(Jenness et al., 2020) developed a network-based transmission model for SARS-CoV-2 on the Diamond Princess outbreak to characterize transmission dynamics and to estimate the epidemiological impact of outbreak control and prevention measures.

(Elbanna et al., 2020) [seems similar to MacPan model!]

(de Celles et al., 2020) Was discussed in the Math 747 SEIR Asymptomatic and symptomatic I_1, I_2 . Used linear chain trick Stringency index as a control force lowering β .

(Rice et al., 2020) Effect of school closures on mortality. Reproduce Report 9 results by spatial agent based CovidSim.

5.8.2 Models of repeated random testing of isolated populations

Bergstrom et al. (2020) (1) Model, assumptions: They developed a function, namely expected exposure $E(C, \tau)$, to approximate trade-offs between the frequency of testing, n , the sensitivity of testing, q , and the delay between testing and results, d . This function is explicitly derived and was connected the effective reproduction number $R = R_0 S$, where S is the proportion of population susceptible. assumption that transmission rates are a step function: individuals who have COVID go from non-infectious to fully infectious instantaneously, and remain fully infectious until they are no longer able to transmit disease. Test sensitivity takes the same form over the course of infection. More sophisticated models could allow varying infectiousness and varying sensitivity over time, as in (Larremore et al., 2020).

(Lopman et al., 2020) Used a Deterministic SEIR model, incorporated TTI, applicable to a university setting. They assumed a fairly high reproductive number that is not reduced through social distancing measures. They found that community-introduction of SARS-CoV-2 infection onto campus can be relatively controlled with effective testing, isolation, contract tracing and quarantine.

(Tuite et al., 2020) used an age-structured compartmental model of COVID-19 transmission in the population of Ontario, Canada. We compared a base case with limited testing, isolation and quarantine to different scenarios.

5.8.3 Other maybe-related works

(Arino and Portet, 2020) developed a SLIAR compartmental model to study the spread of an epidemic, specifically COVID-19, in a population. The model incorporates an Erlang distribution of times of sojourn in incubating, symptomatically and asymptotically infectious compartments. Basic reproduction number is derived. Also, sensitivity analysis with respect to the underlying parameters for the following two outputs was carried out; (i) the number of observable cases during the course of the epidemic and at the peak, and (ii) the timing of the peak of the outbreak. Sensitivity analysis is performed using the R package multisensi.

(Ruszkiewicz et al., 2020) novel with-in-a-minute breath testing with 80% accuracy.