

Testing and Isolation Efficacy: Insights from a Simple Epidemic Model

February 8, 2021

1 Abstract

The potential advantage of a slow rate of test return on the dynamics of an epidemic is real, often neglected, and needs to be quantified. At the individual level, this advantage can manifest in the following sense: individuals awaiting their test results or who have tested positive may partially or fully self-isolate, thus reducing or eliminating their potential in the transmission process.

At the community-level (e.g., nursing homes, LTC facilities, etc.), fast test reporting may be more beneficial. It gives a rapid assessment of the situation, identifies hot spots, and may enable rapid contact-tracing. In this paper, we investigated this individual-level effect of testing processes on the epidemic dynamics by developing a SIR-type model.

Although the model development was motivated by the COVID-19 epidemic, the model has general epidemiological and testing structures. The realistic components of the model include testing intensity, test sensitivity and specificity, rate of test return, and isolation. The novel component is the compartment-specific relative testing weights, which reflect the testing strategies—surveillance, diagnosis, or control. Here, we showed that the slow test reporting can be beneficial on the dynamics of an epidemic. In particular, it is possible for the basic reproduction number, \mathcal{R}_0 , to be increasing with respect to the rate of test return.

Further, we compare two testing strategies, random vs. targeted, in this framework. We conclude that the targeted testing strategy is more effective than the random one in the sense that a lower range of testing intensity is required to keep $\mathcal{R}_0 < 1$. Also, targeted testing reduces the individual-level of advantage of a slow rate of test return.

[Fady: General thoughts on the abstract (take with a grain of salt): I think the abstract is eye-catching and interesting. I think leading in by highlighting potential advantages to a slow test return may be “provocative” (which may not necessarily be bad, I’m not sure). I think the second paragraph in some sense undermines or clashes with the first paragraph, especially since we don’t actually address these potential advantages in this model. Maybe one way to spin this around would be to lead in with a discussion of the advantages of a FAST test return (i.e., the second paragraph), and then say “but situations where returning tests more rapidly is BAD have not been studied”. I think that could have better harmony or coordination.]

2 Introduction

The observed dynamics of the COVID-19 epidemic are driven by both epidemiological processes (infection and recovery) and testing processes (testing and test reporting). In addition to shaping epidemic observations (via case reports), testing processes can also affect epidemiological dynamics. In particular, individuals who are awaiting the results of a test may partially or fully self-isolate, and individuals who have tested positive are more likely to self-isolate. We developed a model that incorporates mechanistic epidemic processes and testing in order to explore the effects of testing and isolation on epidemic dynamics.

Because individuals who have been tested or received a positive test are likely to self-isolate—reducing or eliminating their potentially infectious contacts—the dynamics of an epidemic will depend on who gets tested and why. For a given testing intensity (tests performed per day), testing will have a stronger effect on infection rates the more strongly it is focused on people who are infectious. [DJDE: *deleted “for SARS-CoV-2”. Since this is an idealized model intended for insights, it is probably better to use more generic language most of the time and not say covid unless necessary in a specific context.*] This level of focus depends in turn on the purpose and design of testing programs. (1) When testing is done for the purposes of disease surveillance (?) we might expect tests to be assigned randomly across the population (in an ideal world: some bias in selection for testing is probably inevitable), although a stratified sampling design would be preferred for statistical efficiency (?) [Ali: *better ref?*].

Over the course of the COVID-19 pandemic, however, the vast majority of testing has been done with other goals – primarily diagnosis (determining COVID19 infection status for clinical purposes), or control (determining COVID19 status in order to quickly isolate cases that have been found by contact tracing), which we characterize as *targeted* testing strategies. In these cases, testing probabilities vary widely across epidemiological compartments; in our dynamical model, we will characterize these probabilities by assigning a *per capita* testing weight to each compartment that determines the *relative* probability that an individual in that compartment will be selected for testing (see Methods).

When testing is used primarily for diagnosis it will focus on people with COVID19-like symptoms; thus the testing weights for people with [DJDE: *escape spaces after dots unless it is a full stop*] vs. without SARS-CoV-2 infections will depend on the relative magnitudes of the probability of having COVID19-like symptoms when infected with SARS-CoV-2 (i.e., the relative proportions of presymptomatic or asymptomatic vs. symptomatic infections) and the probability of having COVID19-like symptoms when *not* infected (e.g., due to other respiratory illnesses). Testing for epidemic control will focus on people who are known to have been in contact with known COVID19 cases; in this case the testing weights for infected vs. uninfected people will depend on the probability of infection given contact, as well as the thoroughness and effectiveness of the system for identifying suspicious contacts.

In addition to the overall testing intensity, the testing weights for different compartments, and the degree of self-isolation of test-pending and test-confirmed people, our models will also need to include test specificity and sensitivity [DJDE: *I presume you mean that models would ideally take account of these things, which should not be mentioned unless they are defined since the terms are often misunderstood*], as well as the distribution of times from testing to reporting. Here, we focus on the the effect of testing intensity, different levels of

testing “focus” (from random to highly targeted), and rate of test processing on a single index of epidemiological dynamics, the basic reproduction number \mathcal{R}_0 .

The early dynamics of an epidemic are determined by \mathcal{R}_0 . This index is defined as the expected number of secondary infections arising from a typical infective individual in a completely susceptible population (?). In the early stages of an epidemic the number of infected individuals is expected grow exponentially over time when $\mathcal{R}_0 > 1$. To control the spread of disease in the population, [DJDE: *the effective*] \mathcal{R}_0 must be < 1 . Although the value of \mathcal{R}_0 cannot completely characterize the dynamics of even the simplest epidemic model (?), it does give a simple and widely accepted index for the difficulty of control, as well as some indication of the likely final size of an epidemic (?). [DJDE: *This paragraph does not flow from the rest of the intro, and is a weird way to end when the point of the paper is about testing.*]

[DJDE: *There is some back and forth between generic discussion and COVID-specific comments. I think COVID should be used as motivation, but that the whole paper should be framed generically.*]

3 Methods

We developed a deterministic model, Eqs. (S1)[DJDE: *relegating the equations to supp is not OK. You don't want to publish a paper in which the model is specified exactly only in online supplementary material. I don't think there is any need for supp material. You can have appendices, but I would keep everything as part of the paper.*], which groups individuals based on disease status and testing status. Disease states include Susceptible, Infectious and Recovered (thus this is an SIR-type model), and testing status categorizes people as *untested*, *waiting-for-positive*, *waiting-for-negative*, or *confirmed positive* (Figure 1). Two ‘accumulator’ compartments, N and P , were also incorporated in the model in order to collect cumulative reported negative or positive tests. The model and details of calculation of the basic reproduction number \mathcal{R}_0 are presented in the Supplementary Materials.




Figure 1: Flowchart of the SIR (Susceptible-Infectious-Recovered) model, S1. For the description of the parameters see Table 1.

Table 1 defines the model parameters, which are generally straightforward *per capita* flows between compartments, or modifiers to these flow rates. [DJDE: *you need to define what you mean by “testing intensity” ρ*] The novel component of the model comes in through the compartment-specific relative testing weights W_S , W_I and W_R ; these give the relative rates at which people in the S , I , and R compartments are tested, respectively. For example, $W_I/W_S = 2$ means that infected individuals are tested at twice the *per capita* rate of susceptible individuals.

In order to link to more applied models [DJDE: *what does that mean? what is an “applied model”?*], we constructed this model so we could specify the total *per capita* testing rate. We do this by defining the weighted size of the testing pool $W = W_S S_u + W_I I_u + W_R R_u$, and calculating a scaling parameter for testing as:

$$\sigma = \frac{\rho N_0}{W}. \quad (1)$$

Thus, the *per capita* testing rate for compartment Z is $F_Z = \sigma W_Z$, where $Z \in \{S, I, R\}$. [DJDE: *For a generic compartment, I would suggest X rather than Z , since Z is often the final size.*] For a high-sensitivity test, infected people typically flow through to the “confirmed positive” (I_c , R_c) compartments and are thus unavailable for further testing. Over the course of the epidemic, a fixed testing rate as specified in (1) can (if large enough) exhaust the pool of people available for testing, leading to a singularity when no one is left untested. Although this phenomenon does not affect our analysis of \mathcal{R}_0 , it can affect the temporal dynamics (we discuss an adjustment to the model that solves this problem in the Supplementary material).

[DJDE: *A comment on notation: You haven’t actually defined S_u , I_u etc. In any case, I urge you to use macros and use Roman font for subscripts that are textual as opposed to variables. So S_u and I_p etc. Then it is clear the subscript isn’t a variable with a value.*]

The classical SIR model is based on the following implicit assumptions; well-mixed population, homogeneity of the population [DJDE: *What’s the difference btw “well-mixed” and “homogeneous”?*], exponentially distributed duration of infection and large population size (see, e.g., ?). In addition to these standard assumptions, our model, S1, assumes: (i) there is a single force of infection, Λ , defined as follows

$$\Lambda = \beta \frac{(I_u + \eta_w I_n + \eta_w I_p + \eta_c I_c)}{N_0}, \quad (2)$$

across all susceptible pools (see Table 1 for detail [DJDE: *there are no details in the table; you need to do a better job of explaining what’s going on*]), (ii) $\eta_c \leq \eta_w$, i.e., the individuals awaiting test results have a higher transmission probability than the reported individuals. For this analysis, we also assume a perfectly specific test ($p_S = 0$). This last assumption combined with the assumption that no individual is in waiting-for-positive or confirmed positive compartments, i.e., $S_p(0) = S_c(0) = 0$, reduces the model to 10 equations (equations c and d in (S1) are eliminated).

The Disease-Free Equilibrium (DFE) for the SIR model, Eqs. (S1), is given by solving the coupled system including $S_u + S_n = N_0$ and $F_S S_u - \omega S_n = 0$. [DJDE: *you mean by solving the algebraic equations obtain by setting the DEs to 0. not clear why you bother*

Symbol	Description	Unit	Value
N_0	Total population size	people	10^6
ω	Rate of onward flow from “waiting” to “confirmed” or “untested” compartments	1/day	-
γ	Recovery rate	1/day	1/3
ρ	Per capita testing intensity	1/day	0.01
η_w	Relative probability of transmission for “waiting” individuals	-	-
η_c	Relative probability of transmission for “confirmed positive” individuals	-	-
β	Transmission rate	1/day	0.39
Λ	Force of infection	1/day	-
p_S	Probability of positive tests for S ($= 1 - \text{specificity}$)	-	0
p_I	Probability of positive tests for I ($= \text{sensitivity}$)	-	1
p_R	Probability of positive tests for R ($= 1 - \text{specificity}$)	-	0.5
W_S, W_I, W_R	Relative testing weights	-	Random: $\{1, 1, 1\}$ Targeted: $\{0.3, 1, 1\}$

Table 1: Parameters of the model (S1).

to mention two of these algebraic eqns.] The DFE is [DJDE: don't put blank lines around equations unless you are started a new paragraph. comment out blank lines if want the .tex file to be more readable]

$$S_n^* = \frac{\rho}{\omega} N_0, \quad S_u^* = N_0 - S_n^*, \quad \text{and} \quad I_j = R_j = 0 \quad \text{for all } j. \quad (3)$$

[DJDE: above you had j in Roman font even though it *is* a variable, so I changed to j]

The basic reproduction number, \mathcal{R}_0 , was calculated by using the next generation matrix method developed by ?. \mathcal{R}_0 is

$$\mathcal{R}_0 = (A \times S_u^* + B \times S_n^*) \times C, \quad (4)$$

where

$$\begin{aligned} A &= \gamma(\omega + \gamma) + (\gamma\eta_w + \omega\eta_c p_I) F_I, \\ B &= (\omega + (F_I + \gamma)\eta_w) \gamma + \frac{(\eta_w \gamma + \eta_c \omega) \omega p_I F_I}{\omega + \gamma}, \\ C &= \frac{\beta/\gamma}{N_0(\gamma(\omega + \gamma) + F_I(\gamma + \omega p_I))}. \end{aligned} \quad (5)$$

[DJDE: put β/γ in front of the expression in eq (4), so it is clear that the rest is a modification of the simplest expression. It would be worth spending some time trying to write the expression in a way that helps the reader understand where the different factors come from.] Further details of derivation of \mathcal{R}_0 are provided in the Supplementary material section.

The analytical calculation of the next generation matrix, G [DJDE: don't waste the symbol G on a quantity we don't really need; G is often used for generation interval], and the derivation of \mathcal{R}_0 was carried out in Maple (?) by using simple linear Algebra package. We used R (?) for the simulation part which included using the explicit expression of \mathcal{R}_0 as a function of the underlying parameters, specifying the parameters a realistic range, calculating the corresponding \mathcal{R}_0 and plotting the contours of \mathcal{R}_0 by using ggplot package (ref?) in R. The related result is presented in Figure 2, where panel (2a) represents the random testing, and panel (2b) is representing non-random testing. The parameter values or ranges are as follows (I like using a table here, with the param, their units and values). [DJDE: use comment macro for comments] It is notable that in the simulation part, the explicit expression of \mathcal{R}_0 was used and not the approximation (S6). [DJDE: that's the first mention of "the approximation"] This simulation reflects the behavior of \mathcal{R}_0 with respect to the selected parameters for two different testing strategies: (i) random testing, represented by all testing weights to be the same, $W_S = W_I = W_R$, and (ii) non-random testing, when testing weight are not equal. For simulation purposes we chose $W_S = W_I = W_R = 1$ for random testing, and $W_S = 0.3$ and $W_I = W_R = 1$ for non-random testing strategy. Note that the critical contour of $\mathcal{R}_0 = 1$ is plotted in solid line in Figure 2.

[DJDE: Details such as the fact that you used ggplot are not important to state in the main text. On the other hand, there isn't a clear narrative here. The reader should feel that a story is being told.]

4 Results

The explicit formula for the basic reproduction number, \mathcal{R}_0 (4), provides an opportunity to study the influence of changes in the underlying parameters on the critical index of epidemic dynamics. We are interested in understanding the effect of parameters that can be realistically controlled by changing in isolation, *per capita* testing intensity and test resulting, i.e., η_c and η_w , ρ and ω , respectively. The following Proposition is the direct result of taking partial derivative of \mathcal{R}_0 (4) with respect to selected parameters. Note that we used Taylor approximation of \mathcal{R}_0 at $\rho = 0$ due to the complexity of the expression and thus inconclusive. **[DJDE: in that case it would be good to establish numerically that the relationships given in the proposition are valid more generally, at least for reasonable parameter ranges]** See the Supplementary material for details.

[DJDE: make sure to use \mathcal{R}_0 not R_0 below (and often in the supp material section). More importantly, you really need to state in words what this proposition is saying. The reader has to check the param table and think it through themselves.]

Proposition 1. Using the expression of \mathcal{R}_0 (4),

1. $\partial \mathcal{R}_0 / \partial \eta_c \geq 0$ and $\partial \mathcal{R}_0 / \partial \eta_w \geq 0$.
2. $\partial \mathcal{R}_0 / \partial \rho \leq 0$ when $\rho \approx 0$.
3. $\partial \mathcal{R}_0 / \partial \omega$ can be positive or negative when $\rho \approx 0$.

Given that the perfect isolation occurs at $\eta_c = \eta_w = 0$, **[DJDE: That's a perfect example of the kind of phrasing that really must be changed. You need to say something like "occurs when no one transmits while waiting for test results (i.e., $\eta_c = \eta_w = 0$)". It makes the text so much easier to process and doesn't require the reader to keep looking at the table to make sense of what you're saying.]** Prop. 1 means that lifting the isolation from awaiting group results in an increased \mathcal{R}_0 and consequently a greater number of infected individuals. Prop. 2 indicates that increasing the *per capita* testing intensity, ρ , reduces \mathcal{R}_0 . Lastly, Prop. 3 means that speeding up the test reporting may or may not reduce \mathcal{R}_0 , it will depend on other parameters. It is straightforward to derive 2 and 3 from the Taylor approximation of \mathcal{R}_0 at $\rho = 0$. **[DJDE: can we gain some insight from the expressions for the partial derivatives in the proposition, as opposed to just their signs?]**

Our numerical solutions in support of the analytical results in Proposition 1 are presented in Figure 2. It can be inferred that when random testing strategy is applied, Figure 2a, comparing to the targeted testing strategy, Figure 2b, the critical *per capita* testing intensity of the corresponding panels are lower in non-random testing. Also, it appears that speeding the test reporting, i.e., increasing ω , does not significantly lower \mathcal{R}_0 when awaiting people (including people in I_p and I_n compartments) follow the isolation perfectly, i.e., η_w is closer to 0. However, speeding the test reporting reduces the epidemic more when the awaiting people less follow the isolation, i.e., η_w is closer to 1.

Furthermore, we derived an inequality that quantifies the exact relationships between model parameters that result in returning tests more rapidly being favorable (S18). **[DJDE: What do we learn from this inequality? Why is it relegated to an appendix?]**




Figure 2: **A comparison of the behaviour of the basic reproduction number, \mathcal{R}_0 , between random versus targeted testing strategies at different levels of testing and isolation.** We numerically evaluate \mathcal{R}_0 (4) using the following parameters (listed in Table 1): $N_0 = 1 \times 10^6$, $\omega \in [0.1, 2]$ 1/day, $1/\gamma = 3$ days, $\rho \in [0, 0.01]$ 1/day, $\eta_w \in [0, 1]$, $\eta_c \in [0, 1]$, $\beta = 0.39$ 1/day, $p_S = 0$, $p_I = 1$ and $p_R = 0.5$. Contours of \mathcal{R}_0 are plotted for two testing strategies identified by a set of relative testing weights; (a) random testing where $W_S = W_I = W_R = 1$ and (b) targeted testing where $W_S = 0.3$ and $W_I = W_R = 1$. The black solid line in each panel represents the critical contour of $\mathcal{R}_0 = 1$. **[BMB: we still have some collisions in the y-axis tick labels]**

5 Discussion

Mathematical modeling of infectious disease outbreaks provides insights on how testing processes influence the epidemiological processes through isolation. Here, we develop a compartmental SIR-type model to study the potential effect of testing strategies, testing intensity, test sensitivity and specificity, test reporting time and isolation on epidemic dynamics. While targeted testing strategies **[DJDE: you need to be clearer: you mean targeting individuals who have contacted people who were infected. For example, here you could say something like “While targeting the contacts of confirmed cases...”]** are always more effective than random testing, as expected, we find that in some cases the direct effect of testing is that viral spread is greater for a slow test than for a fast test. This counter-intuitive effect can occur when people are cautious when awaiting a test result, and may not be robust to second-order effects of fast testing (such as better contact tracing).

We incorporated the compartment-specific relative testing weights, W_S , W_I and W_R , to model random testing and *targeted* testing strategies. Here, in the case of targeted testing and for the simplicity and illustration purposes, we assumed that infected and recovered individuals are tested at three times the *per capita* rate of susceptible individuals, thus $W_I/W_S = 3$ and $W_R = W_I$. Note that we have not specified a methodology to assign par-

particular relative testing weights corresponding to a particular targeted testing scenario. This part needs to be developed further in future work. Modeling different targeted testing strategies, equivalently test-specific testing weights in our framework, requires prior information of the conditional probabilities of getting tested for people in a given compartment. This can be implied when we would like to quantify and compare the effect of different levels of test focus **[DJDE: ??]** for infectious people on the basic reproduction number \mathcal{R}_0 , and conclude about the disease spread management. For example, when people are tested for "screening", the individuals with potential higher mobility, eg. people who are getting on flights, get more tested and thus the corresponding heavier testing weight is assigned than people awaiting for a surgery and are probably going to stay in a long-term care facility and consequently less mobile and more isolated to begin with. With our model, we would be able to compare the sensitivity of the disease dynamics, through \mathcal{R}_0 , with respect to testing processes.

The *per capita* testing intensity, ρ ; Proposition 1 part 2 and also Figure 2 indicates that increasing testing intensity ρ reduces \mathcal{R}_0 . This is sensible since as people are moved to test compartments, namely Z_p and Z_n for $Z \in \{S, I, R\}$, they may partially or fully self-isolate. Also individuals who have tested positive, namely Z_c compartment for $Z \in \{S, I, R\}$, are highly likely to self-isolate. The higher probability of being subject to isolation, the lower the force of infection (2) and the lower \mathcal{R}_0 will become. While our simulation supported this result, analytically it appeared to be hard to conclude this due to the complexity of \mathcal{R}_0 expression.

The potential advantage of slow test reporting, or favorable-delay-reporting; Individuals are highly likely to fully self-isolate when they are either awaiting the test results or they are reported, thus reducing or eliminating their potentially infectious contacts. Thus, the faster the test reporting rate, ω , the shorter these individuals stay in the "safe" awaiting-confirmed compartments, namely Z_n , Z_p and Z_c for $Z \in \{S, I, R\}$, and the more they get involved in the infection process. This advantage of slow test reporting is real, and neglected. We also compare to an individual-level advantage of fast tests: people who test positive may be even more careful. In our model analysis, Proposition 1 part 3 is describing this potential advantage. It states that returning test results more rapidly, i.e., increasing the rate ω , does not necessarily lower the reproduction number \mathcal{R}_0 ; whether increasing ω lowers \mathcal{R}_0 depends on the precise combination of model parameters including test reporting rate, testing strategies represented by compartment-specific testing weights, test sensitivity and specificity, and the level of isolation. Specifically, in the case of perfect isolation, i.e., when $\eta_w = 0$ and $\eta_c = 0$, \mathcal{R}_0 may increase as the test reporting process becomes faster. This can be seen from expression (S7). Another example of this favorable delay in ω could be when the test being employed produces many false negatives. Because many infected asymptomatic individuals will believe they are uninfected/uninfectious, thus may unknowingly spread the virus to many others. Again the delay in the test reporting rate keeps these individuals in the "safe" awaiting-confirmed compartments.

we are missing out on community-level advantages of fast testing: better assessment of the situation, identification of hot spots, contact-tracing, etc. The *per capita* testing intensity of CIVID-19 after about a year from the first case reported in December 2019, is still low ($(\rho \approx 0)$ in our model). In near future new test kits may be widely accessible, our model provide insights in this case. In particular, if a cheap test can identify on average

more infected individuals as an expensive test, then our model predicts that the cheap test will lower \mathcal{R}_0 more. In contrast, if an expensive test can identify on average more infected individuals, it will not necessarily lower \mathcal{R}_0 more than the cheap test. The use of tests cheaper than RT-PCR has been proposed as a potential strategy for containing the COVID-19 pandemic. While cheaper tests may be less sensitive and reliable than RT-PCR, they allow for broader and more intense testing. Using our Taylor approximation of \mathcal{R}_0 near $\rho = 0$, we examined what circumstances (i.e., model parameters) make the use of one test more favourable than another, and give a complete description through inequality S19. In general, we found that the expensive test tends to more effectively lower \mathcal{R}_0 when (a) individuals who test positive self-isolate much more than individuals who are waiting for their test result, (b) the time it takes to return tests is much shorter than the mean infectious period, and (c) the testing intensity is much greater for infected individuals than susceptible individuals.

[Ali: *not sure if we want these 2 following paragraphs, or shrink it into a few short sentences?*] In addition to the favorable-delay-reporting observation of our model (S1), the model enables us to quantify the amount of delay required in the test reporting process as a strategy to reduce \mathcal{R}_0 . To give a biological interpretation, we describe the qualitative trends predicted by the inequality (S18) [TODO: *Fady please confirm*][Fady: *yes, looks good to me*]. To summarise, returning test results more rapidly tends to be favourable (i.e., reduces \mathcal{R}_0) when (i) *Confirmed-positive* individuals, lower their contact much more than individuals who are waiting for their test results (i.e., $\eta_c \ll \eta_w$), (ii) the test is highly sensitive (i.e., p_I is close to 1), and (iii) the targeted testing strategy is used (i.e., $W_i \gg W_s$).

one point we can make when discussing the relative weight of testing in different compartments is to discuss pre-testing screening tools, such as surveys or questionnaires. If we have a quantitative description of how the testing intensities affect the dynamics, we can make statements like “our results suggest that employing a pre-testing screening tool can help target infected individuals more effectively. In particular, doubling the sensitivity of the pre-screening tool would *do something* to \mathcal{R}_0 .”

[DJDE: *Some of this reads like notes for discussion rather than text for the paper, so I'm not trying to edit.*]

5.1 Model and calculation of \mathcal{R}_0

The model in the form of a system of ordinary differential equations is

$$dS_u/dt = -\Lambda S_u - F_S S_u + \omega S_n, \quad (\text{S1a})$$

$$dS_n/dt = -\Lambda S_n + (1 - p_S) F_S S_u - \omega S_n, \quad (\text{S1b})$$

$$dS_p/dt = -\Lambda S_p + p_S F_S S_u - \omega S_p, \quad (\text{S1c})$$

$$dS_c/dt = -\Lambda S_c + \omega S_p, \quad (\text{S1d})$$

$$dI_u/dt = \Lambda S_u - F_I I_u + \omega I_n - \gamma I_u, \quad (\text{S1e})$$

$$dI_n/dt = \Lambda S_n + (1 - p_I) F_I I_u - \omega I_n - \gamma I_n, \quad (\text{S1f})$$

$$dI_p/dt = \Lambda S_p + p_I F_I I_u - \omega I_p - \gamma I_p, \quad (\text{S1g})$$

$$dI_c/dt = \Lambda S_c + \omega I_p - \gamma I_c, \quad (\text{S1h})$$

$$dR_u/dt = \gamma I_u - F_R R_u + \omega R_n, \quad (\text{S1i})$$

$$dR_n/dt = \gamma I_n + (1 - p_R) F_R R_u - \omega R_n, \quad (\text{S1j})$$

$$dR_p/dt = \gamma I_p + p_R F_R R_u - \omega R_p, \quad (\text{S1k})$$

$$dR_c/dt = \gamma I_c + \omega R_p, \quad (\text{S1l})$$

$$dN/dt = \omega(S_n + I_n + R_n), \quad (\text{S1m})$$

$$dP/dt = \omega(I_p + R_p), \quad (\text{S1n})$$

where parameters are specified in Table 1. The next generation matrix for this model is $G = FV^{-1}$, where matrix F represents the inflow of new infection to the infected compartments and matrix V represents the flow in the infected compartments when the population is totally susceptible. Matrices F and V are

$$F = \beta/N_0 \begin{bmatrix} S_u & \eta_w S_u & \eta_w S_u & \eta_c S_u \\ S_n & \eta_w S_n & \eta_w S_n & \eta_c S_n \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (\text{S2})$$

$$V = \begin{bmatrix} F_I + \gamma & -\omega & 0 & 0 \\ -(1 - p_I) F_I & \omega + \gamma & 0 & 0 \\ -p_I F_I & 0 & \omega + \gamma & 0 \\ 0 & 0 & -\omega & \gamma \end{bmatrix}, \text{ thus} \quad (\text{S3})$$

$$V^{-1} = \begin{bmatrix} \frac{\omega + \gamma}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & \frac{\omega}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & 0 & 0 \\ \frac{(1 - p_I) F_I}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & \frac{F_I + \gamma}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & 0 & 0 \\ \frac{p_I F_I}{\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I} & \frac{\omega p_I F_I}{(\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I)(\omega + \gamma)} & \frac{1}{\omega + \gamma} & 0 \\ \frac{\omega F_I p_I}{(\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I)\gamma} & \frac{\omega^2 F_I p_I}{(\omega \gamma + \gamma F_I + \gamma^2 + \omega F_I p_I)(\omega + \gamma)\gamma} & \frac{\omega}{(\omega + \gamma)\gamma} & \frac{1}{\gamma} \end{bmatrix}. \quad (\text{S4})$$

The particular form of F with two rows of zeros at the bottom, simplifies G as

$$G = \begin{bmatrix} G_{11} & G_{12} \\ 0 & 0 \end{bmatrix}, \text{ where } G_{11} = C \begin{bmatrix} A S_u & B S_u \\ A S_n & B S_n \end{bmatrix}, \quad (\text{S5})$$

where expressions A , B and C are specified in (5). The block matrix G_{12} does not influence \mathcal{R}_0 (defined as the spectral radius of G). All that matters here are the eigenvalues of G_{11} , which are 0 and \mathcal{R}_0 (4). Also note that, all S_u 's and S_n 's are evaluated at the DFE (3) in matrix F (S2) and the block matrix G_{11} (S5).

5.2 On Testing Rate and Numerical Singularity

In this work, we didn't do any numerical solutions for the trajectories in our analysis. However, if one tries to do so there would be a singularity issue to deal with. Specifically, the numerical singularity issue with the chosen σ (1) is that the population in S compartments appeared to blow up when the DFE is achieved. This is once the only untested people are susceptibles, the FOI will become $\Lambda = 0$, testing rate $F_s = \rho N_0 / S_u$. Thus, the first equation of the model (S1) will become $dS_u/dt = -\rho N_0 + \omega S_n$. Thus changes in S_u will be no longer dependent on S_u with a linear rate of leaving the S_u compartment. IN fact the testing rate, σ , should be formulated such that people from the untested compartments will not be tested if they are not there. One way to fix this issue, is to consider a maximum testing rate, τ (1/day). In general, we want to test at a rate of ρ across the whole population. This won't always be possible, so we impose a maximum rate of τ per testable person and redefine $\sigma = \frac{\tau \rho N_0}{\tau W + \rho N_0}$, with the assumption that $\tau \gg \rho$. This alteration in σ , does not change any results related to \mathcal{R}_0 , thus we only impose it in the simulation of the epidemic dynamic.

5.3 Taylor Approximation of \mathcal{R}_0 at $\rho = 0$

The basic reproduction number, \mathcal{R}_0 , close to $\rho = 0$ can be approximated linearly in ρ by using Taylor approximation. it follows

$$R_0 \approx \beta/\gamma + \frac{\beta\rho}{\omega(\omega + \gamma)\gamma^2 W_S} \left(\gamma(\eta_w - 1)(\gamma W_S + \omega W_I) + (\eta_c - 1)P_i W_i \omega^2 \right) + \mathcal{O}(\rho^2). \quad (\text{S6})$$

Also, to analyze the influence of ω on \mathcal{R}_0 , the approximation (S6) was used. Then it is straight forward to have

$$\partial R_0 / \partial \omega = \frac{-\beta\rho}{\gamma W_S \omega^2 (\gamma + \omega)^2} (a\omega^2 + b\omega + c), \quad (\text{S7})$$

where $a = (\eta_w - 1)W_I - (\eta_c - 1)P_i W_I = ((s - p_I)\eta_c + (p_I - 1))W_I$, $b = 2(\eta_w - 1)\gamma W_S$ and $c = (\eta_w - 1)\gamma^2 W_S$. Given that $0 \leq \eta_c \leq \eta_w \leq 1$, one can easily derive $b \leq 0$ and $c \leq 0$.

Note that in general, the necessary and sufficient condition for $a \geq 0$ is $(s - p_I)\eta_c \geq (1 - p_I)$, where $s = \frac{\eta_w}{\eta_c} \geq 1$.

In the case of "perfect isolation", i.e., when $\eta_w = 0$ and consequently $\eta_c = 0$, it is straightforward to see that $a \leq 0$, $b < 0$ and $c < 0$. Thus, $\partial \mathcal{R}_0 / \partial \omega \geq 0$. As an example, in the case of a very accurate testing regime, i.e., $P_i = 1$, $a \geq 0$ is achieved. If $a \geq 0$, the

quadratic expression in (S7), has Real roots. Assuming that $\omega_1 < 0$ and $\omega_2 > 0$ be the roots of the quadratic expression in $\partial\mathcal{R}_0/\partial\omega$. Thus, $\partial\mathcal{R}_0/\partial\omega > 0$ for $0 < \omega < \omega_2$ and $\partial\mathcal{R}_0/\partial\omega < 0$ for $\omega > \omega_2$.

5.4 rate of returning tests

The linearization of \mathcal{R}_0 around $\rho = 0$ is

$$\mathcal{R}_0 \approx \beta/\gamma + \frac{\beta\rho}{\omega(\omega + \gamma)\gamma^2 W_s} \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_i W_i \omega^2 \right). \quad (\text{S8})$$

So when $\rho \approx 0$ we have

$$\partial\mathcal{R}_0/\partial\omega \approx \frac{-\beta\rho}{\gamma W_s \omega^2 (\gamma + \omega)^2} (a\omega^2 + b\omega + c).$$

where $a = (\eta_w - 1)W_i - (\eta_c - 1)P_i W_i$, $b = 2(\eta_w - 1)\gamma W_s$ and $c = (\eta_w - 1)\gamma^2 W_s$.

Perhaps counter-intuitively, the equation above does not predict that \mathcal{R}_0 is monotone decreasing with respect to ω . In other words; our model does not predict that returning test results more rapidly *always* lower \mathcal{R}_0 . In order to gain insight into this intriguing behavior, we examine the zeroes of $\frac{\partial\mathcal{R}_0}{\partial\omega}(\omega)$.

Defining the following quantity, Q , will help us write the roots of $\partial\mathcal{R}_0/\partial\omega$ neatly.

$$Q = \frac{W_i}{W_s} \left(1 - \frac{n_t - 1}{n_w - 1} P_i \right) \quad (\text{S9})$$

$$(\text{S10})$$

With that in mind, we can write the roots of $\partial\mathcal{R}_0/\partial\omega$ as

$$\omega_1 = \frac{\gamma}{-\sqrt{1 - Q} - 1} \quad (\text{S11})$$

$$\omega_2 = \frac{\gamma}{\sqrt{1 - Q} - 1}. \quad (\text{S12})$$

Note that the zeroes are real if and only if $Q < 1$. Note that have $\eta_c < \eta_w$, so if $P_i \approx 1$, we will have $Q < 0 < 1$. Thus, if we assume near-perfect test sensitivity, ω_1 and ω_2 will be real.

Assuming ω_1, ω_2 are real, it is easy to confirm that $\omega_1 < 0$ by looking at the denominator. To see that $\omega_2 > 0$, recall that $Q < 0$, so $\sqrt{1 - Q} > 1$ and so $\sqrt{1 - Q} - 1 > 0$. Knowing that $\omega_1 < 0$, the only root of interest (i.e., biologically relevant quantity) is ω_2 .

We can prove that $\partial\mathcal{R}_0/\partial\omega > 0$ when $\omega \in (0, \omega_2)$ and $\partial\mathcal{R}_0/\partial\omega < 0$ when $\omega \in (\omega_2, \infty)$ by computing the limits of $\partial\mathcal{R}_0/\partial\omega$ at 0 and ∞ respectively. So it follows that \mathcal{R}_0 has a global maximum with respect to ω at $\omega = \omega_2$.

Now we want to characterize the parameter regions on which $\partial\mathcal{R}_0/\partial\omega < 0$ (i.e., the conditions under which returning test results more rapidly is favorable). By the previous analysis, this is equivalent to solving for $\omega > \omega_2$. So

[**TODO:** *Fady, can you clean this up*][**Fady:** *Done (I think)*]

$$\omega > \omega_2$$

$$\omega > \frac{\gamma}{\sqrt{1-Q}-1}$$

$$\sqrt{1-Q} > \frac{\gamma}{\omega} + 1 \quad (\text{S13})$$

$$1-Q > \left(\frac{\gamma}{\omega} + 1\right)^2 \quad (\text{S14})$$

$$(\text{S15})$$

362 Substituting in Q from (S9) we have

$$1 - \frac{W_i}{W_s} \left(1 - \frac{n_t - 1}{n_\omega - 1} P_i\right) > \left(\frac{\gamma}{\omega} + 1\right)^2 \quad (\text{S16})$$

$$- \frac{W_i}{W_s} \left(1 - \frac{n_t - 1}{n_\omega - 1} P_i\right) > \left(\frac{\gamma}{\omega} + 1\right)^2 + 1 \quad (\text{S17})$$

$$\frac{W_i}{W_s} \left(\frac{1 - n_t}{1 - n_\omega} P_i - 1\right) > \left(\frac{\gamma}{\omega} + 1\right)^2 + 1. \quad (\text{S18})$$

363 Since all steps in deriving (S18) are reversible, (S18) gives a necessary and sufficient
364 condition for $\omega > \omega_2$, which characterizes when returning tests more rapidly would cause a
365 decrease in \mathcal{R}_0 .

366 5.5 Expensive vs. cheap tests

367 The use of tests cheaper than RT-PCR has been proposed as a potential strategy for con-
368 taining the COVID-19 pandemic. While cheaper tests may be less sensitive and reliable
369 than RT-PCR, they allow for broader and more intense testing. In the analysis below, we
370 compare the \mathcal{R}_0 predicted by our model depending on the testing strategy.

Consider a test that allows us to test at rate ρ_1 and has sensitivity $P_{i,1}$, and another test that allows us to test at ρ_2 and has sensitivity $P_{i,2}$. Suppose that $\rho_1 > \rho_2$. Recall that the linearization of \mathcal{R}_0 around $\rho \approx 0$ is given by

$$\mathcal{R}_0 \approx \beta/\gamma + \frac{\beta\rho}{\omega(\omega + \gamma)\gamma^2 W_s} \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_i W_i \omega^2 \right).$$

Treating \mathcal{R}_0 as a function of ρ and P_i , we can reduce the inequality

$$\mathcal{R}_0(\rho_2, P_{i,2}) < \mathcal{R}_0(\rho_1, P_{i,1})$$

371 into

$$\begin{aligned} & \rho_1 \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_{i,1} W_i \omega^2 \right) - \rho_2 \left(\gamma(\eta_w - 1)(\gamma W_s + \omega W_i) + (\eta_c - 1)P_{i,2} W_i \omega^2 \right) > 0 \\ & \vdots \\ & \frac{\rho_2 P_{i,2} - \rho_1 P_{i,1}}{\rho_1 - \rho_2} > \frac{1 - \eta_w}{1 - \eta_c} \cdot \frac{\gamma(\gamma W_s + \omega W_i)}{\omega^2 W_i} \end{aligned} \quad (\text{S19})$$

Note that the RHS is positive, thus a necessary condition for the inequality above to hold is that $\rho_2 P_{i,2} > \rho_1 P_{i,1}$, equivalently

$$\frac{P_{i,2}}{P_{i,1}} > \frac{\rho_1}{\rho_2}. \quad (\text{S20})$$

To state an example of this, if test A is three times as expensive as test B (and hence one can test three times as many people with test B), using test A rather than B will be favorable only if test A is at least 3 times more sensitive than test B . Note that this is a necessary but not sufficient condition, so even if test A is three times more sensitive, it is still possible for test B to be more effective.

Eq. (S19) tells us precisely when a test corresponding to $\rho_2, P_{i,2}$ will yield a lower \mathcal{R}_0 than a test corresponding to $\rho_1, P_{i,1}$, where $\rho_1 > \rho_2$. Some of the qualitative *trends* that favor test 2 (the higher-sensitivity test) include

- individuals who test positive self-isolate much more than individuals who are waiting for their test result.
- the time it takes to return tests is much shorter than the mean infectious period.
- the testing intensity is much greater for infected individuals than susceptible individuals.

5.6 Literature Review

5.6.1 Explicit models of TTI (trace/test/isolate) based on network or agent-based models

(?) [Ali: *It seems to me that this is just a statistical model to estimate the parent-offspring of an infected index, not sure if it fits into agent-based group!*] Used simulation on a branching process model to assess the forward and backward contact tracing efficiency. Assuming a negative-binomial branching process with a mean R , reproduction number, and overdispersion parameter k , the mean total number of generation G_3 and averted G_3 are estimated. The effectiveness of TTI is defined as the ratio of averted to the mean.

(?) developed a network-based transmission model for SARS-CoV-2 on the Diamond Princess outbreak to characterize transmission dynamics and to estimate the epidemiological impact of outbreak control and prevention measures.

(?) [seems similar to MacPan model!]

(?) Was discussed in the Math 747 SEIR Asymptomatic and symptomatic I_1, I_2 . Used linear chain trick Stringency index as a control force lowering β .

(?) Effect of school closures on mortality. Reproduce Report 9 results by spatial agent based CovidSim.

5.6.2 Models of repeated random testing of isolated populations

? (1) Model, assumptions: They developed a function, namely expected exposure $E(C, \tau)$, to approximate trade-offs between the frequency of testing, n , the sensitivity of testing,

q, and the delay between testing and results, d. This function is explicitly derived and was connected the effective reproduction number $R = R_0 S$, where S is the proportion of population susceptible. assumption that transmission rates are a step function: individuals who have COVID go from non-infectious to fully infectious instantaneously, and remain fully infectious until they are no longer able to transmit disease. Test sensitivity takes the same form over the course of infection. More sophisticated models could allow varying infectiousness and varying sensitivity over time, as in (?).

(?) Used a Deterministic SEIR model, incorporated TTI, applicable to a university setting. They assumed a fairly high reproductive number that is not reduced through social distancing measures. They found that community-introduction of SARS-CoV-2 infection onto campus can be relatively controlled with effective testing, isolation, contact tracing and quarantine.

(?) used an age-structured compartmental model of COVID-19 transmission in the population of Ontario, Canada. We compared a base case with limited testing, isolation and quarantine to different scenarios.

5.6.3 Other maybe-related works

(?) developed a SLIAR compartmental model to study the spread of an epidemic, specifically COVID-19, in a population. The model incorporates an Erlang distribution of times of sojourn in incubating, symptomatically and asymptotically infectious compartments. Basic reproduction number is derived. Also, sensitivity analysis with respect to the underlying parameters for the following two outputs was carried out; (i) the number of observable cases during the course of the epidemic and at the peak, and (ii) the timing of the peak of the outbreak. Sensitivity analysis is performed using the R package multisensi.

(?) novel with-in-a-minute breath testing with 80% accuracy.