

# STATISTICAL PATTERN RECOGNITION

## ASSIGNMENT 5

Ali Gholami

Department of Computer Engineering & Information Technology  
Amirkabir University of Technology

<https://aligholamee.github.io>  
[aligholami7596@gmail.com](mailto:aligholami7596@gmail.com)

### Abstract

**Keywords.** *Dimensionality Reduction, Principal Component Analysis, Fisher Linear Discriminant Analysis, Feature Subset Selection, Sequential Feature Selection, Data Visualization & Representation.*

## 1 Sequential Feature Selection

Given the following objective function, use **SFS**, **SBS** and **Plus-2 Minus-1 Selection** to select 3 features:

$$J(x) = 5x_1 + 7x_2 + 4x_3 + 9x_4 + 3x_5 - 2x_1x_2 + 2x_1x_2x_3 - 2x_2x_3 - 4x_1x_2x_3x_4 + 3x_1x_3x_5$$

### Solution

#### SFS

In this method, we start feature selection from an empty set. We add features one by one and compute the value of the objective function with respect to each of the features being added. The feature with the largest objective function will be selected. The iteration goes on until all features are covered. We then select ideal features (subset with  $k$  features and maximum objective). The actual algorithm is as following:

1. Start with the empty set  $Y_0 = \emptyset$
2. Select the next best feature  $x^+ = \operatorname{argmax}[J(Y_k + x)]$
3. Update  $Y_{k+1} = Y_k + x^+$ ;  $k = k + 1$
4. Go to 2

Below is the demonstration of iterations taken to completely explore the search space. The first iteration is:

- $J(x_1) = 5$
- $J(x_2) = 7$
- $J(x_3) = 4$
- $J(x_4) = 9$
- $J(x_5) = 3$

According to the heuristic nature of sequential subset selection, we'll choose  $x_4$  as the first best feature. We'll then generate subsets containing combination of features with  $x_4$ :

- $J(x_4x_1) = 14$
- $J(x_4x_2) = 16$
- $J(x_4x_3) = 13$
- $J(x_4x_5) = 12$

Thus, features  $x_4$  and  $x_2$  are selected until now. We'll drive the 3 sized subsets:

- $J(x_4x_2x_1) = 19$
- $J(x_4x_2x_3) = 18$
- $J(x_4x_2x_5) = 22$

Three best features selected by the algorithm are  $x_4$ ,  $x_2$  and  $x_5$ .

## SBS

This method initiates the feature selection procedure using a complete subset of features. It then removes each feature and evaluates the objective function. The feature that causes the lowest decrease in the objective function will be remove (useless feature!). We'll stop when we reach a satisfying 3 sized feature subset. The algorithm is formally working as follows:

1. Start with the full set  $Y_0 = X$
2. Remove the worst feature  $x^- = \operatorname{argmax}[J(Y_k - x)]$
3. Update  $Y_{k+1} = Y_k - x^-$ ;  $k = k + 1$
4. Go to 2

Applying this algorithm on the given objective function yields the following results:

- $J(x_1x_2x_3x_4x_5) = 25$

And the results of removing each of the features:

- $J(x_1x_2x_3x_4) = 19$
- $J(x_1x_2x_3x_5) = 20$
- $J(x_1x_2x_4x_5) = 22$
- $J(x_1x_3x_4x_5) = 24$
- $J(x_2x_3x_4x_5) = 21$

It is obvious that  $x_2$  is the most useless feature among these. We'll remove  $x_2$  and obtain the feature subset with 4 features:  $x_1, x_3, x_4$  and  $x_5$ .

- $J(x_1x_3x_4x_5) = 24$

We can obtain the following subsets:

- $J(x_1x_3x_4) = 18$
- $J(x_1x_3x_5) = 15$
- $J(x_1x_4x_5) = 17$
- $J(x_3x_4x_5) = 16$

$x_5$  will be removed since it has the lowest effect on the greatness of evaluation. The proper feature subset includes:  $x_1, x_3$  and  $x_4$ .

## 2 PCA & FLDA

In this problem, dimensionality reduction with a two-feature two-class dataset is explored. Consider the following dataset and the test sample:  $x = [0.851.15]^T$

- **Class 1:**  $[[0.8, 1.2], [0.9, 1.4], [1.2, 1.4], [1.1, 1.5]]$
  - **Class 2:**  $[[0.8, 1.1], [0.6, 1], [0.65, 1.1], [0.75, 0.9]]$
- Demonstrate the preprocessing steps (need to show step-by-step details). Calculate the **mean** of each class ( $m_1$  and  $m_2$ ). Calculate the **covariance** matrix of each class.
  - Using **Fisher's Linear Discriminant** to find a projection vector ( $w$ ) which optimally separates the projections of these two classes.
  - Is the vector derived from *FLD* along the same direction as the  $m_1 - m_2$ ? Plot both of them on the same figure.
  - Using **Principal Component Analysis** to reduce the dimension to 1 and plot the principal component on the same figure.

- (e) Comment on the differences between **FLD** and **PCA** and  $m_1 - m_2$ . Make up a scenario where **FLD** will be aligned, perpendicular to  $m_1 - m_2$ , if possible at all.
- (f) Project the test sample  $x$  onto  $w$  derived from **FLD** and determine its label.
- (g) Project the test sample  $x$  onto the principal axis from **PCA** and determine its label.

## Solution

(a) Mean of each class  $k$  can be calculated using (2.1).

$$m_k = \frac{1}{|n_k|} \sum_{i=1}^{n_k} x_i \quad (2.1)$$

Thus, for each class we'll have to following results:

- $m_1 = \frac{1}{4}[0.8 + 0.9 + 1.2 + 1.1 \quad 1.2 + 1.4 + 1.4 + 1.5]^T = [1 \quad 1.3]$
- $m_2 = \frac{1}{4}[0.8 + 0.6 + 0.65 + 0.75 \quad 1.1 + 1 + 1.1 + 0.9]^T = [0.7 \quad 1]$

Covariance matrix for each class  $k$  can be calculated using (2.2).

$$\Sigma_k = \frac{1}{n_k - 1} \sum_{i=1}^{n_k} (x - m_k)(x - m_k)^T \quad (2.2)$$

Thus, the results for each class will be:

- $\Sigma_1 = \frac{1}{3} \begin{bmatrix} -0.2 & -0.1 & 0.2 & 0.1 \\ -0.1 & 0.1 & 0.1 & 0.2 \end{bmatrix} \begin{bmatrix} -0.2 & -0.1 \\ -0.1 & 0.1 \\ 0.2 & 0.1 \\ 0.1 & 0.2 \end{bmatrix} = \begin{bmatrix} 0.03 & 0.01 \\ 0.01 & 0.02 \end{bmatrix}$
- $\Sigma_2 = \frac{1}{3} \begin{bmatrix} 0.1 & -0.1 & -0.05 & 0.05 \\ 0.1 & 0 & 0.1 & -0.1 \end{bmatrix} \begin{bmatrix} 0.1 & 0.1 \\ -0.1 & 0 \\ -0.05 & 0.1 \\ 0.05 & -0.1 \end{bmatrix} = \begin{bmatrix} 0.008 & 0 \\ 0 & 0.01 \end{bmatrix}$

(b) In order to find a proper projection vector ( $w$ ), we shall compute the **within class scatter** matrix. Using the definition of  $S_w$ :

$$S_w = S_1 + S_2 \quad (2.3)$$

which is equal to the addition of scatter matrices. Scatter matrix of class  $k$  can be obtained from its covariance matrix using (2.4).

$$S_k = (|n_k| - 1)\Sigma_k \quad (2.4)$$

Replacing the results from the first part into (2.4) yields the following results:

- $S_1 = 3 \begin{bmatrix} 0.03 & 0.01 \\ 0.01 & 0.02 \end{bmatrix} = \begin{bmatrix} 0.09 & 0.03 \\ 0.03 & 0.06 \end{bmatrix}$
- $S_2 = 3 \begin{bmatrix} 0.008 & 0 \\ 0 & 0.01 \end{bmatrix} = \begin{bmatrix} 0.024 & 0 \\ 0 & 0.03 \end{bmatrix}$

Thus, the within class scatter matrix can be written as following:

$$S_w = \begin{bmatrix} 0.117 & 0.03 \\ 0.03 & 0.09 \end{bmatrix}$$