



Advanced Optimization

Homework 1

Autumn 1400
Due date: Aban 12th



Analytical Questions

1. Consider the function $f(x_1, x_2) = (x_1 + x_2^2)^2$. At the point $x^T = (1, 0)$ we consider the search direction $p^T = (-1, 1)$. Show that p is a descent direction and find all minimizers of the following problem at each iteration t .

$$\min_{\alpha > 0} f(x^t + \alpha p^t)$$

2. In the update rule of Newton method,

$$x_{k+1} = x_k - \alpha_k (\nabla^2 f(x_k))^{-1} \nabla f(x_k)$$

one way to obtain the step-size is by solving the following sub-problem,

$$\alpha_k = \arg \min_{\alpha \geq 0} f(x_k - \alpha_k (\nabla^2 f(x_k))^{-1} \nabla f(x_k)).$$

For the objective function $f(x) = \frac{1}{2}x^T Q x - x^T b$ where $Q = Q^T \succ 0$, show if with this approach the algorithm would be able to reach the optimal point x^* where $\nabla f(x^*) = 0$, in just one iteration or not.

3. For minimizing $f(x) = \|x\|^\beta$ while $\beta > 1$ with pure Newton method, show that for what value of β the algorithm converges to the optimum point? What happens if $\beta \leq 1$?

Hint : $(I + BA)^{-1}B = B(I + AB)^{-1}$

Computer Questions

1. For the function,

$$f(x) = 100(x_2 - x_1^2)^2 + (1 - x_1)^2$$

- Compute the gradient $\nabla f(x)$ and Hessian $\nabla^2 f(x)$.
- Start from $x_0 = (-4, 10)^T$, minimize these functions with steepest descent and Newton methods while using backtracking line search. Make comparison between the results. Is there a way to use the benefits of these two methods at the same time?
- Report function value, distance to optimal point and step-size in 3 separate plots. Explain the results.
- Try different strategies for initial step-size.

2. Minimize the below continuous function,

$$f(x) = x^3 - 60x^2 + 900x + 100$$

with Simulated Annealing while considering following terms:

- Variable x is represented as a string of 5 bits.
- The neighborhood defined as randomly flipping just one bit.
- The initial point is 10011. ($x = 19, f(x) = 2399$)
- Analyze two scenarios:
 - initial temperature T_0 equals to 500
 - initial temperature T_0 equals to 100
- Cooling coefficient is equal to 0.9.

Report the results thoroughly.