To get started, do the following:


1. Download the script $tut6_spline.m$ from Canvas.

To get started, do the following:

- 1. Download the script $tut6_spline.m$ from Canvas.
- 2. Set the random seed to be rng(5). Run the script. You should see the following figure:


To get started, do the following:

- 1. Download the script $tut6_spline.m$ from Canvas.
- 2. Set the random seed to be rng(5). Run the script. You should see the following figure:



To get started, do the following:

- 1. Download the script $tut6_spline.m$ from Canvas.
- 2. Set the random seed to be rng(5). Run the script. You should see the following figure:



3. Goal: Compute the coefficients required to plot the missing $S_1(x)$

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

Recall from lecture the following conditions for our cubic spline:

1. S(x) interpolates our data points

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

- 1. S(x) interpolates our data points
- 2. S(x) is continuous acrosss the data points x_j

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

- 1. S(x) interpolates our data points
- 2. S(x) is continuous acrosss the data points x_j
- 3. S'(x) is continuous acrosss the data points x_j

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

- 1. S(x) interpolates our data points
- 2. S(x) is continuous acrosss the data points x_j
- 3. S'(x) is continuous acrosss the data points x_j
- 4. S''(x) is continuous acrosss the data points x_j

Write down (in words) the set of conditions we want our Cubic Spline S(x) to satisfy, and number them (you should have 5). You can already assume we know S(x) is piece-wise cubic

- 1. S(x) interpolates our data points
- 2. S(x) is continuous acrosss the data points x_j
- 3. S'(x) is continuous acrosss the data points x_j
- 4. S''(x) is continuous acrosss the data points x_i
- 5. End point Conditions (e.g Natural, Clamped, Not-a-knot,....)

Based on the conditions you wrote down, write down the set of equations that need to be solved for the coefficients of $S_0(x), S_1(x), S_2(x)$ in our problem. Use the following notation:

Based on the conditions you wrote down, write down the set of equations that need to be solved for the coefficients of $S_0(x), S_1(x), S_2(x)$ in our problem. Use the following notation:

$$S_j(x) = f_j + b_j(x - x_j) + c_j(x - x_j)^2 + d_j(x - x_j)^3$$

Based on the conditions you wrote down, write down the set of equations that need to be solved for the coefficients of $S_0(x), S_1(x), S_2(x)$ in our problem. Use the following notation:

$$S_j(x) = f_j + b_j(x - x_j) + c_j(x - x_j)^2 + d_j(x - x_j)^3$$

You may also use the notation $h_j = x_{j+1} - x_j$, as in the notes. Which boundary conditions do you need to enforce? You may ask or look it up!

Based on the conditions you wrote down, write down the set of equations that need to be solved for the coefficients of $S_0(x), S_1(x), S_2(x)$ in our problem. Use the following notation:

$$S_j(x) = f_j + b_j(x - x_j) + c_j(x - x_j)^2 + d_j(x - x_j)^3$$

You may also use the notation $h_j = x_{j+1} - x_j$, as in the notes. Which boundary conditions do you need to enforce? You may ask or look it up!

• Next to each equation, write the number that corresponds to the condition being enforced.

Note that the condition $S(x_j)=f_j$ is already enforced by our notation. We have 3 unknowns per $S_j(x)$, therefore 9 total unknowns. Our 9 equations are:

Note that the condition $S(x_j) = f_j$ is already enforced by our notation. We have 3 unknowns per $S_j(x)$, therefore 9 total unknowns. Our 9 equations are:

$$f_{0} + b_{0}h_{0} + c_{0}h_{0}^{2} + d_{0}h_{0}^{3} = f_{1} \quad \textcircled{1}$$

$$f_{1} + b_{1}h_{1} + c_{1}h_{1}^{2} + d_{1}h_{1}^{3} = f_{2} \quad \textcircled{1}$$

$$f_{2} + b_{2}h_{2} + c_{2}h_{2}^{2} + d_{2}h_{2}^{3} = f_{3} \quad \textcircled{1}$$

$$b_{0} + 2c_{0}h_{0} + 3d_{0}h_{0}^{2} = b_{1} \quad \textcircled{3}$$

$$b_{1} + 2c_{1}h_{1} + 3d_{1}h_{1}^{2} = b_{2} \quad \textcircled{3}$$

$$2c_{0} + 6d_{0}h_{0} = 2c_{1} \quad \textcircled{4}$$

$$2c_{1} + 6d_{1}h_{1} = 2c_{2} \quad \textcircled{4}$$

$$6d_{0} = 6d_{1} \quad \textcircled{5}$$

$$6d_{1} = 6d_{2} \quad \textcircled{5}$$

Note that the condition $S(x_j) = f_j$ is already enforced by our notation. We have 3 unknowns per $S_j(x)$, therefore 9 total unknowns. Our 9 equations are:

$$f_{0} + b_{0}h_{0} + c_{0}h_{0}^{2} + d_{0}h_{0}^{3} = f_{1} \quad \textcircled{1}$$

$$f_{1} + b_{1}h_{1} + c_{1}h_{1}^{2} + d_{1}h_{1}^{3} = f_{2} \quad \textcircled{1}$$

$$f_{2} + b_{2}h_{2} + c_{2}h_{2}^{2} + d_{2}h_{2}^{3} = f_{3} \quad \textcircled{1}$$

$$b_{0} + 2c_{0}h_{0} + 3d_{0}h_{0}^{2} = b_{1} \quad \textcircled{3}$$

$$b_{1} + 2c_{1}h_{1} + 3d_{1}h_{1}^{2} = b_{2} \quad \textcircled{3}$$

$$2c_{0} + 6d_{0}h_{0} = 2c_{1} \quad \textcircled{4}$$

$$2c_{1} + 6d_{1}h_{1} = 2c_{2} \quad \textcircled{4}$$

$$6d_{0} = 6d_{1} \quad \textcircled{5}$$

$$6d_{1} = 6d_{2} \quad \textcircled{5}$$

The last two equations impose Not-a-knot end point conditions!

In our plot, only b_1, c_1, d_1 are unknown. Write down the 3 equations needed to solve for these coefficients. Use the values given to you on the plot to write down numerical values for all known quantities. Note: The spacing between x values is 1/3.

In our plot, only b_1, c_1, d_1 are unknown. Write down the 3 equations needed to solve for these coefficients. Use the values given to you on the plot to write down numerical values for all known quantities. Note: The spacing between x values is 1/3.

Based on the given information, we can write down the following equations:

$$S_1(x_2) = 1.2463 + \frac{1}{3}b_1 + \frac{1}{9}c_1 + \frac{1}{27}d_1 = 0.28287$$

$$S_1'(x_1) = b_1 = -1.0362$$

$$S_1''(x_2) = 2c_1 + 2d_1 = 6.3656$$

Using the second equation, we can fid b_1 , and we are left with a 2×2 system for c_1 and d_1 :

Using the second equation, we can fid b_1 , and we are left with a 2×2 system for c_1 and d_1 :

$$\frac{1}{9}c_1 + \frac{1}{27}d_1 = -0.6180$$
$$2c_1 + 2d_1 = 6.3656$$

Using the second equation, we can fid b_1 , and we are left with a 2×2 system for c_1 and d_1 :

$$\frac{1}{9}c_1 + \frac{1}{27}d_1 = -0.6180$$
$$2c_1 + 2d_1 = 6.3656$$

Which has the solution:

$$c_1 = -9.9344, \quad d_1 = 13.1175$$

Cubic Splines - Check our answer

Once we have values for b_1, c_1, d_1 , fill in the vector c1 in the code so that it has the form

$$c1 = [d_1 c_1 b_1 f_1]$$

Run the code again and see if you correctly computed $S_1(x)$