



# Hypoglycemia (Adults)

Blood sugar  $< 3.9$  mmol/l + hypoglycaemic symptoms: **Check Conscious Level**

**Conscious**

**Unconscious:**

If patient **oriented & able to swallow** Give:

If patient **disoriented & unable to cooperate** Give:

**Call Ambulance**  
**Check ABCDE** Give:

Fast acting **oral glucose 15g**  
Followed by **long-acting carbohydrates**

IM/SC  
**GLUCAGON 1mg**

IM/SC: **GLUCAGON 1mg**

OR

IV: **50% Dextrose 50ml**

SE: nausea and vomiting → put patient in recovery position

**N.B: Glucagon Dosing (Pediatrics)**

- $< 8$  years OR  $< 25$  kg: **0.5 mg IM/SC**
- $\geq 8$  years OR  $\geq 25$  kg: **1 mg IM/SC**

- **Remeasure blood glucose** after 15 minutes & repeat treatment until blood sugar  $> 4$  mmol/l
- If patient on **Sulfonylureas: Refer** to Hospital in spite of correcting blood
- If **child is unconscious or unsafe to take oral carbohydrates**, IV dextrose should be administered: **Initial bolus - 2 mL/kg of dextrose 10%** (0.2 g dextrose/kg body weight), If glucose fails to increase after 15 to 20 minutes, repeat bolus. Higher concentrations of dextrose are not recommended as an initial bolus