# Counselling/Preventive

A 50-year-old diabetic woman admitted to the medical ward with chest pain. The chest X-ray is suggestive of empyema. The radiologist was consulted and advised to do CT guided aspiration. Who should perform the informed consent process?
A. The radiologist
B. The head nurse
C. The internal medicine resident
D. Any member of the treating team

Answer: A
An intemal medicine resident is teaching the medical students how to examine a diabetic patient. He forgot to close the curtain while exposing the patient's abdomen. Which one of the ethical principles is violated?
A. Respect patients dignity
B. Respect patients privacy
C. Respect patients autonomy
D. Respect patients confidentiality

Answer: B
A 70-year-old man is diagnosed with metastatic cancer. His family asked the oncologist not to tell their father of the diagnosis. Which of the following is the most appropriate action?
A. Consult the hospitals legal advisor
B. Inform the patient about his diagnosis
C. Consult the hospitals ethics committee
D. Do not inform the patient about his diagnosis

Answer: B
A 78-year man with post chemotherapy and radiotherapy end stage cancer is suffering from unbearable pain. His family asked the oncologist to increase the pain killer doses to relive their father's pain. The oncologist agreed after explaining to the family that the higher doses may hasten their father's death. Which of the following principles is used to make the decision?
A. Principle of totality
B. Principle of subsidiarity
C. Principle of double effect
D. Principle of informed choice

Answer: C
A 70-year-old woman admitted for a hip replacement operation. The surgeon was interrupted by the patient when he was trying to explain the surgery and its possible complications. The patient expressed her wish to not knowing the details. The surgeon respected her wish and informed the anesthesiologist who insisted to explain to the patient all details. Which of the following actions should be taken by the surgeon?
A. Consult the hospitals ethics committee
B. Let the anesthesiologist deal with the patient
C. Inform the patient about the anesthesiologist decision
D. Reschedule the operation and find another anesthesiologist

Answer: A

A 40-year-old diabetic man presents to the Emergency Room with charcot knee. He was stabilized and transferred to the Surgical Ward. The surgical team explained to the patient the need for amputation. The patient refused. Which of the following is the most appropriate next action?
A. Assess the patient mental capacity
B. Refer the patient to another hospital
C. Consult the hospitals clinical ethics committee
D. Ask the patient to sign discharge against medical advice form

Answer: A
A 20-year-old man was admitted for a simple tonsillectomy. The operation took longer time than usual because of a bleeding resulted from a surgical error. The surgery was successful and the patient fully recovered with no complications. Next day the patient asked the surgeon why his operation took longer time than usual. Which of the following is the most appropriate action?
A. Consult the hospital's ethics committee
B. Reassure the patient and tell him what happened
C. Ask another surgeon from the team to tell the patient
D. Reassure the patient but do not tell him what happened

Answer: B
A 30-year-old woman presented to the Emergency Room with abdominal pain. The signs and symptoms are suggestive of acute appendicitis. During the surgery, the appendix found in normal state. Appendectomy was done as it is the common practice is to remove the appendix even if it is not inflamed. The surgeon does not remember that he informed the patient about this decision during the consent procedure. What should the surgeon do when the patient recovered next day?
A. Tell the patient that it is part of the standard procedure
B. Ask another surgeon from the team to tell the patient
C. Consult the hospital's ethics committee
D. Do not tell the patient what happened

Answer: A
A 90-year-old man with advanced metastatic lung cancer was admitted to the Surgical Intensive Care Unit with deteriorating pulmonary functions and labelled a do not resuscitate case by the intensivist. The thoracic surgeon would like to do a de-bulking operation to enable the patient to breath but not sure, if the patient could survive the operation, Which of the following is the most appropriate action?
A. Explain to the patient and ask him to sign high risk consent form
B. Do not operate since the patient is likely not going to improve
C. Consult the hospitals ethics committee
D. Discuss the case with the intensivist

Answer: B

which of the following is the percentage prevalence of smoking among Saudi adult men?
A. 11 %
B. 21 %
C. 37 %
D. 51 %

Answer: B
A 42-year-old man presented to the Primary Care Clinic for the routine checkup. He has been smoking 1- pack of cigarettes a day for 22 years. He has no intention of quitting and feels fine. Which of the following is the most appropriate cessation approach?

A. Set a "quit date" with the patient
B. Prescribe nicotine replacement therapy
C. Give clear, personalized advice to quit
D. Refer him to classes for smoking cessation

Answer: C
A 45-year-old smoker came to the clinic for his diabetes follow-up. During discussion, he acknowledged that smoking is not good for his health. He plans to quit this year. According to the Stages of Change Model, at what stage of change is this patient?
A. Precontemplation
B. Contemplation
C. Preparation
D. Maintenance

Answer: B
Tobacco products consists of thousands of chemical compounds including nicotine. Which of the following is the most outcome related to nicotine substance?
A. COPD
B. Addiction
C. Lung cancer
D. Heart diseases

Answer: B
Which of the following is the most approved evidence-based intervention to be used in smoking cessation?
A. Hypnosis
B. Varenicline
C. Acupuncture
D. BICOM bioresonance device

Answer: B
Which of the following is the most effective preventive measure to improve post-myocardial infarction prognosis?
A. Lowering lipid
B. Regular exercise
C. Smoking cessation
D. Maintaining an ideal body weight

Answer: C
Which of the following sets indicates the 5 A's approach during smoking cessation counselling?
A. Ask, advise, assess, assist, arrange follow-up
B. Ask, assist, advise, advocate, arrange follow-up
C. Assess, advocate, ask, advise, arrange follow-up
D. Advocate, ask, assess, advise, arrange follow-up

Answer: A

What is the minimum number of days per week recommended to practice muscle-strengthening physical activity?
A. 2
B. 3
C. 4
D. 5

Answer: A
Which of the following is the most appropriate intervention for primary prevention of type 2 diabetes?
A. Screening
B. Pharmacotherapy
C. Physical activity
D. Bariatric surgery

Answer: C
Which of the following is the recommended minimum number of minutes for an effective small session of aerobic physical activity?
A. 5
B. 10
C. 15
D. 20

Answer: B
Which of the following is the most appropriate characteristic for a new test as a confirmatory test for a disease diagnosis?
A. Specific
B. Sensitive
C. Non-Invasive
D. Easy to administer

Answer: A
which of the following is the recommended age in years to start screening for dyslipidemia in a low-risk man?
A. 18
B. 25
C. 35
D. 50

Answer: C
A 55-year-old woman has just completed her screening mammogram and the results were normal. She has no symptoms and no family history of breast cancer. When she should have another mammogram?
A. 6 months
B. 1 year
C. 2 years
D. 3 years

Answer: C
What is the recommended age in years to start screening for high blood pressure in adults with no risk factors?
A. 18
B. 25
C. 35
D. 50

Answer: A
Which of the following is a recommended age in years for a low-risk adult man to start screening for colon cancer?

A. 30
B. 40
C. 50
D. 60

Answer: C
A screening test for liver cirrhosis was administered to 800 people with proven liver cirrhosis and to 800 people without liver cirrhosis. The test results were positive for 200 of the proven cases and 100 of the normal persons. What is the sensitivity of the screening test?
A. 25 %
B. 33 %
C. 67 %
D. 88 %

Answer: A
A new screening test for diabetes has a sensitivity of 90% and specificity of 80%. Which of the following is the best interpretation?
A. 80% of non-diabetics had positive result
B. 80% of diabetic patients had positive result
C. The test was positive for 90% of diabetic patients
D. For each 100 positive test results, 90% were diabetics

Answer: C
An 82-year-old woman is admitted at noon with pneumonia to the General Medical Ward. At night, she is found disoriented and restless by the nursing staff.
Blood pressure 152/88 mmHg
Heart rate 99 /min
Respiratory rate 22 /min
Temperature 38.2 °C
Oxygen saturation 94 %
Which one of the following is the best initial step the nurse should do?
A. Call her family to come and watch her at bedside
B. Restrain the patient physically in her bed
C. Orient the patient to her environment, start nasal 02 and raise the rails of her bed
D. Call the duty doctor for Lorazepam order of 5 mg IV Stat

Answer: C
A 77-year-old woman is admitted to Neurology Ward with right hemispheric intracranial bleeding. 5 days after admission the physiotherapist reported verbally to the nurse that the patient left buttock is getting red with skin loss. The nurse forgot to tell the resident and no action was taken. 2 days later, the patient spiked a fever with hypotension. On examination, the resident found purulent discharge from the pressure sore. Which of the following is the best approach to prevent this incident?
A. Better nutritional support
B. Better communication
C. More nursing training
D. Extra nursing staff

Answer: B
A 92-year-old man was admitted after fall at home. There was no fracture. Foley catheter was inserted due to immobility from pain. His appetite was good. Acetaminophen with codeine was prescribed for pain. 4 days later, the patient became confused with agitation. The nurse noted cloudy urine in the bag. Blood pressure 168/92 mmHg

Heart rate 105 /min
Respiratory rate 18 /min
Temperature 38.2 °C
Oxygen saturation 98 %
What is the best action to prevent this complication?
A. Avoid Codeine in this age group
B. Request frequent turning of the patient by family
C. Review the need for urinary catheter daily
D. Add nutritional supplement twice daily

Answer: C

A 45-year-old woman underwent open laparotomy for a ruptured appendicitis. 3 days postoperatively, she developed fever again with severe abdominal pain and vomiting. On examination, her abdomen was rigid with decreased bowel sounds. CT scan showed foreign body in right iliac fossa. Patient was taken urgently back to Operating Room and was found to have a missed infected sponge in her peritoneal cavity (see lab results).
Test Result Normal Values
WBC 27 4.5-10.5 x 109/L
Neutrophils 95 40-60%
Creatinine 87 44-115 mol/L
Which of the following is the most appropriate action by the surgeon?
A. Report the surgical assistant mistake to the administration
B. Do not inform the patient about the finding
C. Call his malpractice lawyer and report the incident
D. Inform the patient with apology about the mistake

Answer: D
An inter was attending Operation Room for laparoscopic nephrectomy for suspicious renal mass. The urologist asked to prep for the right kidney approach. However, the intern remember from his history and exam that it was the left kidney. Which of the following is the most appropriate action by the intern?
A. Keep quiet since he might be wrong
B. Leave OR and check the electronic note
C. Call the chief intern to ask for guidance
D. Inform the surgeon about his concern

Answer: D
While doing ward round the consultant surgeon asked the resident to write Pipercillin-Tazobactam for their patient. The pharmacist reported that the patient has a questionable allergy to penicillin. The consultant insisted to write the order. Which of the following is the most appropriate action by the resident?
A. Write the order and let the consultant sign it
B. Write the order then call chief surgeon to complain
C. Gently refuse to write the order
D. Write the order and keep quiet fearing retaliation

Answer: C

Which of the following is considered a secondary prevention for diabetes?
A. Screening
B. Chronic diseases management
C. Education about healthy lifestyle

D. Immunization against infectious diseases

Answer: A
In cardiovascular disease prevention, which of the following is considered primordial prevention?
A. Prevention of coronary heart disease risk factors
B. Initiation of therapy to reduce recurrent coronary heart disease events
C. Decrease cardiac mortality in patients with established coronary heart disease
D. Modification of risk factors in order to prevent or delay the onset of coronary heart disease

Answer: A
Which of the following factors is directly related to increased risk of stroke?
A. Obesity
B. Diabetes
C. Hypertension
D. Hypercholesterolemia

Answer: C
Which of the following is the strongest risk factor for diabetes in adults?
A. Stress
B. Pregnancy
C. Abdominal obesity
D. High body mass index

Answer: C
Which of the following interventions is the best to use with pregnant women who smoke?
A. Bupropion
B. Varenicline
C. Behavioral therapy
D. Nicotine replacement therapy

Answer: C
What is the latent period (incubation period) for lung cancer caused by cigarette smoking (in years)?
A. =5
B. =10
C. =15
D. =20

Answer: D
After how many years the excess risk of myocardial infarction will decline significantly for individual who stop smoking?
A. 1
B. 2
C. 3
D. 4

Answer: A
How many times the risk of myocardial infarction is increased in smokers compared to nonsmokers?
A. 2
B. 3
C. 4
D. 5

Answer: A

On average, how many years of life the average smokers are expected to lose?
A. 5
B. 10
C. 15
D. 20

Answer: B
How many times the risk of lung cancer is increased in smokers compared to nonsmokers?
A. 5
B. 10
C. 15
D. 20

Answer: C
in a screening program in a population with a low prevalence of a particular disease, what most of the positive results would likely to be?
A. True positive
B. False positive
C. True negative
D. False negative

Answer: B
Natural history of the disease has various stages. In which of the following stages a screening program would be most useful?
A. Pre-disease stage
B. Clinical disease stage
C. Pre-clinical disease stage
D. Recovery, disability or death stage

Answer: C
Which of the following criteria is supporting the use of a screening test for community screening program?
A. The disease is rare
B. The test is very specific
C. The test is quick and easy
D. Natural history of the disease is not yet clear

Answer: C
Which of the following screening test parameter is most likely to be affected by the disease prevalence?
A. Specificity
B. Sensitivity
C. Likelihood ratios
D. Predictive values

Answer: D

Which of the following cancers is recommended to be screened for in asymptomatic adults?
A. Skin cancer
B. Colon cancer
C. Thyroid cancer
D. Bladder cancer

Answer: B

What is the recommended age to start screening for osteoporosis with bone measurement testing in women?
A 50-54
B. 55-59
C. 60-64
D. 65-69

Answer: D

which of the following provides the highest level of evidence for establishing effect of exercise on the quality of life?
A. Cohort study
B. Clinical trials
C. Case control study
D. Observational study

Answer: B
Which of the following is the most appropriate measurement to estimate the impact of obesity prevention on the prevalence of type 2 diabetes?
A. Odds ratio
B. Relative risk
C. Attributable risk
D. Cross product ratio

Answer: C
Which of the following is the most important characteristic to consider when choosing a screening test for cancer?
A. Cheap
B. Highly specific
C. Highly sensitive
D. Quick to perform

Answer: C
which of the following is most likely to increase the risk of cardiovascular disease?
A. Sedentary lifestyle
B. Having a BMI of 28
C. Having a HbA1c of 6%
D. Consuming only 5 servings/day of vegetables

Answer: A
Which of the following is recommended to improve the general health of a person?
A. Monitoring dietary protein intake
B. Performing regular physical activity
C.Switching from cigarette use to waterpipe
D. Consuming at least 2 servings of fruits and vegetables per day

Answer: B
Which of the following fulfils the minimum recommended time of moderate physical activity for healthy adults?
A. 20 minutes for 6 times per week
B. 30 minutes for 5 times per week

C. 40 minutes for 3 times per week
D. 50 minutes for 2 times per week

Answer: B
Which of the following is the most significant risk factor for developing breast cancer among women?
A. Age
B. Late menopause
C. Early menarche
D. Delaying childbirth until after 30

Answer: A

A 75-year-old woman presented with significant height loss. Which of the following is the most appropriate investigation?
A. Bone mineral density
B. Measure serum 25-OH-D
C. Fracture risk assessment
D. lateral thoracic and lumbar spine X-ray

Answer: D
Which of the following number of sleep hours is associated with more weight gain?
A. <5
B. 6
C. 7
D. >7

Answer: A
For which of the following cancers "population screening" is recommended?
A. Oral cancer
B. Ovarian cancer
C. Pancreatic cancer
D. Colorectal cancer

Answer: D
Which of the following is the treatment goal for diabetic patients with coronary artery disease?
A. Triglycende < 150
B. Blood pressure < 140/90
C. Low density lipoprotein < 70
D. High density lipoprotein > 70

Answer: C
Which level of prevention we apply when detecting a pregnant woman infected with hepatitis B?
A. Primordial
B. Primary
C. Secondary
D. Tertiary

Answer: C
Which of the following best explains the increase of diseases prevalence when incidence is constant?
A. Loss at follow-up
B. Poor case ascertainment
C. Improved quality of care

D. Increased diagnostic accuracy

Answer: C
Which of the following is the major preventable cause for dyslipidemia and coronary heart disease in women?
A. Diet
B. Smoking
C. Low sleep quality
D. Physical Inactivity

Answer: B
A 65-year-old man with a history of myocardial infarction (see lab results).
Blood pressure 140/90 mmHg
BMI 25 kq/m2
Test Result Normal Values
Cholesterol (LDL) 3.7 <3.36 mmol/L Cholesterol (HDL) 1.1 > 1.03 mmol/L
Apo A-1 1.37 0.88-1.8 g/L (Male) 0.98-2.1 g/L (Female)
Apo B-100 1.16 0.55-1.51 g/L (Male)
0.44-1.48 g/L (Female)
Which of the following is the most powerful risk prediction value for cardiovascular disease?
A. Hypertension
B. Body mass index
C. LDL cholesterol
D. Apo B/Apo A1 Ratio

Answer: D

# Psychiatry

11. A 78-year-old man presents with a 2-year history of slow and progressive deterioration in his memory. The patient is able to perform some activities of daily living, including dressing and bathing. His personality has changed from that of a kind and caring person into someone who has periods of agitation and aggression. Neurologic and musculoskeletal examination cannot be carried out.
Which of the following is the most likely diagnosis?
A. Mixed dementia
B. Alzheimer's disease
C. Multi-infarct dementia
D. Major depressive disorder

Answer: B

12. A 46-year-old woman with schizophrenia presents to the clinic with a 1-month history of abnormal orofacial movements. She has been prescribed amisulpride 600mg for many years and procyclidine 5 mg daily.
Which of the following is the most likely cause?
A. Tardive dyskinesia
B. Catatonic phenomenon
C. Cerebrovascular accident

Answer: A

13. Which of the following is the peak of nicotine withdrawal symptoms after smoking cessation?
A. 1 day
B. 2 to 4 days
C. 5 to 7 days
D. 8 to 10 days

Answer: B

14. Which of the following is an alternative treatment for severe depression?
A. Psychotherapy
B. Tricyclic antidepressants
C. Electroconvulsive therapy
D. Electroencephalogram therapy

Answer: C

15. Which of the following is the antidote for an opiate overdose?
A. Naloxone
B. Flumazenil
C. Lorazepam
D. Buprenorphine

Answer: A

16. Which of the following is the most likely side effect of bupropion when prescribed for smoking cessation?
A. Arrhythmia
B. Xerostomia
C. Headache
D. Seizure

Answer: D

17. A 24-year-old inactive woman presents to the clinic with a 5-month history of amenorrhea. On examination, she has a body mass index of 20, stable for the last 5 years (see lab results).
Pregnancy test: Negative.
Which of the following is the most likely diagnosis?
A. Eating disorder
B. Illegal drug use
C. Pituitary neoplasm
D. Chronic depression

Answer: A

18. Which of the following factors is the most important in determining the success of a smoking cessation program?
A. Patient's desire to quit smoking
B. Behavior modification
C. Pharmacologic agents
D. Medical counseling

Answer: A

19. A 45-year-old man newly diagnosed with type 2 diabetes presents to the clinic for further management. He is on time for the appointment, but because of an emergency case, the

physician arrives late to the clinic. When the physician walks in, the patient appears extremely angry.

Which of the following is the most appropriate response by the physician?

A. Acknowledge the patient's anger
B. Explore the reasons for the anger
C. Help the patient understand their anger
D. Empathize with newly diagnosed diabetic patient

Answer: A

20. A 56-year-old man presents to the clinic for evaluation. The physician has difficulty in obtaining accurate information regarding the reason for attendance.

Which of the following is the most appropriate next step in consultation?

A. Use medical terms that the patient understands
B. Control the interview with direct questions
C. Involve the patient in the treatment plan
D. Use open-ended questions

Answer: D

21. A 76-year-old man admitted with a diagnosis of pneumonia has fluctuating levels of attention, changes in awareness, and a marked disturbance of the sleep/wake cycle.

Which of the following is the most appropriate initial management?

A. Isolation in a quiet, dark room
B. Allow a relative to stay with the patient
C. Haloperidol regularly until symptoms stop
D. Risperidone regularly until symptoms abate

Answer: B

22. A 27-year-old woman delivered a healthy baby 3 months ago. She is currently undergoing psychotherapy for postpartum depression.

Which of the following is the most appropriate to be added to optimize treatment?

A. Prescribe very small doses of imipramine
B. Encourage continuation of breastfeeding
C. Include family members in therapy
D. Prescribe a calcium supplement

Answer: C

23. A 23-year-old man complains of having an unusually prominent jaw. He has been seen by 3 plastic surgeons, but each one has refused to do anything. On examination, he appears depressed and has no visible protrusion, the jaw and face are normal. He has a past medical history of rhinoplasty and blepharoplasty.

Which of the following is the most likely diagnosis?

A. Dysthymia
B. Hypochondriasis
C. Somatization disorder
D. Body dysmorphic disorder

Answer: D

24. A 41-year-old man presents to the clinic requesting a complete examination to exclude colon cancer. He recently had some investigations done and provided a list of test results. He complains that this fear is interfering severely with his work and social activities.

Which of the following is the most likely diagnosis?
A. Hypochondriasis
B. Factitious disorder
C. Somatization disorder
D. Obsessive-compulsive disorder

Answer: A

25. A 16-year-old boy learns about the death of his father and within hours, his speech becomes disorganized, he is walking outside of his house with no clothes on, and he claims that his father is inside the house and telling him to do these things in his honor. The symptoms resolved after 3 days, and the son returned to full pre-morbid level of function but is obviously confused over the events of the past few days.

Which of the following is the most likely diagnosis?

A. Schizophreniform disorder
B. Schizoaffective disorder
C. Brief psychotic disorder

Answer: C

26. Which of the following is a prediction of the feeling of hopelessness?

A. Impulse control problems
B. Gender identity disorder
C. Suicidal behavior
D. Panic disorder

Answer: C

27. A 20-year-old woman presents to the clinic with a history of abdominal pain, nausea, vomiting, localized weakness to the right leg, and has a history of irregular menses. In the past, she has sought treatment for head, back, joint, and chest pain. Factitious disorder and malingering have been excluded as diagnoses.

Which of the following is the most likely diagnosis?
A. Hypochondriasis
B. Conversion disorder
C. Somatization disorder
D. Somatoform pain disorder

Answer: C

Explanation: somatization disorder is characterized by a history of multiple unexplained physical complaints across various organ systems, while somatic symptom disorder (formerly somatoform pain disorder) is characterized by persistent, distressing somatic symptoms that lead to significant impairment, with a focus on the meaning or impact of the symptoms.

28. Which of the following investigations should be recommended for an elderly patient with cognitive deficit?
A. Brief IQ test
B. Hearing loss evaluation
C. Wechsler memory scale
D. Involuntary movement test

Answer: B

1. An 8-year-old boy recently moved to another city. Soon after moving, school started, but the boy hesitated to make friends with anyone and had a hard time paying attention in class. Which of the following is the most likely diagnosis?
A. Dysthymia
B. Hypomania
C. Depression
D. Adjustment disorder

Answer: D
2. A 48-year-old man presents for a routine visit. While reporting his medical history, he reveals a significant alcohol intake. The physician wants to assess the drinking further using the cut- annoyed-guilty-eye (CAGE) screening questionnaire.

Which of the following is a question in the CAGE screening questionnaire?
A. "Do you eat more after you drink?"
B. "Do you think you are an alcoholic?"
C. "Have you ever felt guilty about your drinking?"
D. "How many drinks do you consume?"

Answer: C

3. A 46-year-old woman presents to the clinic with an 8-month history of irritability, excessive worry, and difficulty falling asleep and concentrating. Which of the following is the most likely diagnosis?
A. Mood disorder not otherwise specified
B. Major depressive disorder with anxiety
C. Anxiety disorder with depression
D. Generalized anxiety disorder

Answer: D

4. A 42-year-old woman is concerned about losing her job because of tiredness and would like to take antidepressants because of having a difficult time falling asleep and getting up in the morning. Her sleep disturbance is related to the inability to relax after getting into bed. She repeatedly checks to make sure that the door is locked, the oven is off, and the children's school supplies are organized. A similar routine occurs in the morning. She states that her behavior is out of control but if is not done, she will not be able to sleep or leave the house. Which of the following disorders is the most likely diagnosis?
A. Obsessive-compulsive
B. Generalized anxiety
C. Somatization
D. Delusional

Answer: A

5. A 35-year-old healthy man presents to the clinic with a 3-month history of feeling depressed. He is distressed at work and burnt-out. Four months earlier, he was promoted to a managerial position and was having an interpersonal conflict with 2 of the employees. Which of the following is the most appropriate next step in management?
A. Selective serotonin reuptake inhibitors
B. TCA tricyclic antidepressant
C. Supportive psychotherapy
D. Phenothiazine

Answer: C

6. An 18-year-old man presents to the Emergency Department with a 20-minute history of lightheadedness, headaches, and nausea. He complains of tingling around the mouth and in the fingertips. On examination, he is anxious, tremulous, sweating, and breathing heavily. Which of the following is the most appropriate first action?
A. Patient rebreathe into a paper bag
B. Conduct an amobarbital interview
C. Blood alcohol concentration
D. CT Brain scan

Answer: A

7. Which of the following atypical antipsychotic agents is associated with the most weight gain?
A. Ziprasidone
B. Aripiprazole
C. Olanzapine
D. Quetiapine

Answer: C

8. Which of the following is the best treatment for generalized anxiety disorder?
A. Buspirone
B. Bupropion
C. Alprazolam
D. Escitalopram

Answer: D

9. For which of the following childhood psychiatric disorders, clozapine is the most appropriate medication?
A. Bipolar illness
B. Schizophrenia
C. Substance abuse
D. Major depression

Answer: B

10. Which of the following antidepressants has high efficacy in childhood and adolescent depression?
A. Lithium
B. Clozapine
C. Fluoxetine
D. Imipramine

Answer: C

11. A 30-year-old man presents to the clinic with a 3-month history of aliens talking to him and putting thoughts into his mind. Which of the following is the most appropriate management?
A. Behavioral therapy
B. Antipsychotic therapy
C. Bereavement counseling
D. Antidepressant medications

Answer: B

12. A 55-year-old man presents to the clinic with a history of difficulty maintaining erections and impotence that occurs during stressful periods of life. He is currently under stress at work. On examination, he has normal physical findings. Which of the following is the first step in management?

A. Prescribe sildenafil
B. Refer to an endocrinologist
C. Refer to a urologist for evaluation
D. Encourage use of relaxation strategies

Answer: D

13. A 38-year-old woman presents with a 3-month history of recurrent intense fear accompanied by irritability, excessive worry, palpitations, and chest tightness. She is avoiding social gatherings fearing of having the same symptoms. Physical examination findings are normal. Which of the following medications is the most appropriate first line of treatment?

A. Selective serotonin reuptake inhibitor
B. Monoamine oxidase inhibitor
C. Tricyclic antidepressants
D. Benzodiazepine

Answer: A

14. A 5-year-old boy is brought to the clinic by his mother with a history of impaired language development, impaired intelligence, compulsive repetitive behavior, and a preoccupation with inanimate objects. Which of the following is the most likely diagnosis?

A. Autism
B. Dyslexia
C. Conductive hearing loss
D. Manic depressive disorder

Answer: A

15. A mother of a 3-month-old baby presents to the clinic and tells the physician that she is "going crazy." She states that since the baby has been born, she has continuously visualized a snake crawling into the baby's crib at night. She states that she knows it is not likely to happen but cannot get the thought out of her mind. She checks the baby's crib up to 50 times per night to ensure that there is no snake. Which of the following is the most likely diagnosis?

A. Delusions
B. Obsessions
C. Hallucinations
D. Postpartum psychosis

Answer: B

16. A 50-year-old woman presents to the clinic with a history of anxiety. Two months ago, during a meeting at her job, she became very anxious, sweaty, and short of breath. She had to leave the meeting to calm down. Since then, she has avoided meeting rooms out of fear that she will have a recurrence of her symptoms. Which of the following is the most likely diagnosis?

A. Panic disorder
B. Specific anxiety disorder
C. Generalized anxiety disorder
D. Post-traumatic stress disorder

Answer: A

17. A 24-year-old man presents to the clinic with a history of gradual fear of automobiles. He has always been anxious about cars, but now can no longer look at them without experiencing shortness of breath, sweating, and fear. Which of the following disorders is the most likely diagnosis?
A. Panic
B. Specific anxiety
C. Generalized anxiety
D. Post-traumatic stress

Answer: B

18. A 21-year-old woman presents to the clinic with a history of red dry skin on her hands. She washes her hands up to 70 times per day because she always feels dirty. Washing her hands relieves her anxiety, but as she touches anything, she wants to wash again. Which of the following disorders is the most likely diagnosis?
A. Obsessive-compulsive
B. Dysmorphic
C. Phobic
D. Panic

Answer: A

19. A 30-year-old man with a history of drug abuse presents to the clinic weekly with a variety of health complaints, which vary each time, and are greatly exaggerated. When he is unaware of being observed he appears asymptomatic. Which of the following is the most likely diagnosis?
A. Depression
B. Malingering
C. Somatic delusion
D. Conversion reaction

Answer: B

20. A 19-year-old woman presents to the clinic for a routine physical check-up. On examination, she has fine hair all over the body. She feels fat and subsequently diets. BMI 16.8 kg/m2 Which of the following is the most likely diagnosis?
A. Bulimia
B. Anorexia nervosa
C. Body dysmorphic disorder
D. Generalized anxiety disorder

Answer: B

21. A 48-year-old woman calls the clinic expressing fear of leaving the house and has been inside for 2 months. She becomes anxious at the thought of walking out the front door and is now unable to enter the room containing the front door. Which of the following is the most likely diagnosis?
A. Depression
B. Malingering
C. Agoraphobia
D. Substance abuse

Answer:C

22. A 27-year-old man presents to the clinic with a history of sweating, palpitations, nausea, and a "panicky feeling" when required to deliver a speech at work. He has no other difficulties at work. Which of the following is the most likely diagnosis?
A. Malingering

B. Agoraphobia
C. Performance anxiety
D. Generalized anxiety disorder

Answer: C

23. A 56-year-old woman presents to the clinic with a history of constant counting of floor tiles, cars, items on a dinner plate, and words in a conversation. She is upset by the counting but becomes anxious and agitated when she tries to stop herself from counting. Which of the following is the most likely diagnosis?
A. Delusion
B. Obsession
C. Compulsion
D. Hallucination

Answer: C

24. A 61-year-old man presents with a history of constantly thinking about aliens landing in his backyard, especially when away from home. Despite knowing that aliens do not exist, he feels overwhelmed with the idea that they have landed. He fears that he is "going insane." Which of the following is the most likely diagnosis?
A. Delusion
B. Obsession
C. Compulsion
D. Hallucination

Answer: B

25. A 38-year-old man presents to the clinic with a history of hearing voices coming from the toaster and refrigerator, warning that all the food is poisoned. Which of the following is the most likely diagnosis?
A. Delusional thinking
B. Visual hallucination
C. Disorganized thinking
D. Auditory hallucination

Answer: D

26. A 27-year-old man presents to the clinic for a routine check-up. On examination, he kept looking to the right frequently. When questioned, he said, "My mother is there, but no one can see her except me." He claims that she comes and goes, even though the family indicates she died when the patient was a child. Which of the following is the most likely diagnosis?
A. Delusion
B. Visual hallucination
C. Auditory hallucination
D. Episode of disorganized thinking

Answer: B

27. A 17-year-old man with a history of intravenous drug abuse presents to the clinic for clinical evaluation. Which of the following is the most appropriate next step in investigation?
A. Methicillin-resistant Staphylococcus aureus
B. Human immunodeficiency virus
C. Streptococcus viridans
D. Viral hepatitis B

Answer: B

28. A 17-year-old high school athlete presents to the clinic for an annual physical check-up. On examination, he has gained 20 Kg in the past 4 months and has very well-developed musculature of the upper body, acne over the face and back and foul breath.
Heart rate 94 /min
Temperature 37.6° C
Blood pressure 138/89 mmHg
Respiratory rate 14 /min

Which of the following should the patient be screened for?
A. Depression
B. Cocaine use
C. Thyroid tumor
D. Anabolic steroid use

Answer: D

29. A 52-year-old man presents to the clinic for a routine physical check-up. He has a history of pulling out hair when under stress. On examination, he has diffuse hair thinning along with the absence of eyelashes. Which of the following is the most likely diagnosis?
A. Post-traumatic stress disorder
B. Cicatricial syndrome
C. Trichotillomania
D. Alopecia areata

Answer: C

1. A 15-year-old girl presents to the clinic with history of tetany, swollen glands, and eroded dental enamel (see lab result).

Electrolytes test Result Normal Value
Potassium 2.8 3.5-5.1 mmol/L

Which of the following is the most likely diagnosis?
A. Bulimia nervosa
B. Substance abuse
C. Anorexia nervosa
D. Alcohol withdrawal

Answer: A

2. A 40-year-old drug addict man presents to the Emergency Department with 1-hour history of unconsciousness. On examination, he has pinpoint pupils and a respiratory rate of 8/ minute.
Which of the following is the most appropriate treatment?
A. Atropine
B. Naloxone
C. Furosemide
D. N-acetylcysteine

Answer: B

3. A 53-year-old woman with history of depression presents to the Emergency Department with history of nausea, vomiting, epigastric pain, tinnitus, and shortness of breath 6 hours after consuming 50 tablets of aspirin (see lab result).
Test Result Normal Value
Plasma salicylate concentration 54 15-30 mg/dL
Which of the following is the most appropriate initial management?

A. Hemodialysis
B. Gastric lavage
C. Activated charcoal
D. Urine alkalization

Answer: D

4. A 34-year-old healthy man presents to the clinic with a 2-year history of worsening sad mood, poor appetite, insomnia, fatigue, low self-esteem, and feelings of hopelessness.
Which of the following is the most likely diagnosis?
A. Organic mood disorder
B. Depressive disorder
C. Minor depression
D. Dysthymia

Answer: D

5. A 9-year-old girl is brought to the clinic by her mother with a bald patch on her scalp. Her mother noted that she plucks her hair while studying and under stress.
Which of the following is the most appropriate treatment?
A. Olanzapine
B. Bupropion
C. Fluoxetine
D. Lithium

Answer: C

6. A 20-year-old man presents to the clinic with poor personal hygiene, echolalia, echopraxia, muttering to self, sitting in an odd posture for hours, and insomnia for the past 2 months. On examination, his orientation and memory are normal.
Which of the following is the most appropriate treatment?
A. Oxcarbazepine
B. Amisulpride
C. Venlafaxine
D. Lithium

Answer: B

7. A 45-year-old woman presents to the Primary Health Care Clinic with symptoms of anxiety due to difficulties in her relationship with her co-workers. Her examination was normal and the doctor has decided that there is no need for any investigation.
Which of the following would be the most effective approach for counselling?
A. Instructing how to deal with others
B. Empathy during consultation
C. Asking humorous questions
D. Standing up for her rights

Answer: B

8. A 55-year-old smoker presents to the Primary Health Care Clinic for the result of the biopsy of a lesion in his right lung. When informed that he has cancer in his right lung, he asked with anxiety, "Is it a mistake?"
Which of the following is the most appropriate description for the patient's reaction?
A. Anger
B. Denial
C. Bargaining
D. Depression

Answer: B

9. A 30-year-old man presents to the clinic with a 10-month history of restlessness, difficulty concentrating, and disturbance in his sleep pattern. He is diagnosed with generalized anxiety disorder.
Which of the following is the most appropriate treatment?
A. Alprazolam
B. Bupropion
C. Haloperidol
D. Naltrexone

Answer: A

Explanation: Benzodiazepines are the 2nd line treatment. Bupropion is atypical antidepressant.
10. An 18-year-old man presents to the Emergency Department with a 1-week history of euphoria, increased blood pressure, tachycardia, nausea, weight loss, psychomotor agitation, chills, sweating, and visual hallucinations.
Which of the following is the most likely diagnosis?
A. Amphetamine intoxication
B. Cannabis intoxication
C. Acute schizophrenia
D. Cocaine withdrawal

Answer: A

11. A 3-year-old child is brought to the clinic for his yearly physical examination. His history revealed various behavioral issues, including poor interaction with peers and family, delayed language development, repetitive movements and difficulty of accepting change. The child stiffens physically when touched, does not respond to the examiner and does not make eye contact.
Which of the following is the most likely diagnosis?
A. Attention deficit hyperactivity disorder
B. Asperger's syndrome
C. Autistic disorder
D. Down syndrome

Answer: C

12. A 23-year-old woman is admitted to the Psychiatric Unit after causing superficial lacerations to her wrists. The reason is that she felt that her therapist has abandoned her. She has a depressed mood that varies by the hour. On admission, she claimed that she has heard a voice speaking to her, although she currently denies it. She caused a disagreement between the psychiatric resident and the nursing staff, who have conflicting views of her behavior in the unit.
Which of the following personality disorders is the most likely diagnosis?
A. Antisocial
B. Borderline
C. Obsessive
D. Schizoid

Answer: B

13. A 40-year-old man presents to the Emergency Department with irritability, suspiciousness, overactivity and poor hygiene. Emergency Department doctors regarded his behavior as abnormal and called for a psychiatrist.
Which of the following is the most likely diagnosis?
A. Acute confusional state

B. Alzheimer's Disease
C. Schizophrenia
D. Psychoses

Answer: D

14. A 58-year-old man presents to the clinic with a history of disturbed recent memory as a cardinal feature of his disturbed state of mind along with behavioral changes.
Which of the following is the most likely diagnosis?
A. Multiple Sclerosis
B. Alzheimer's Disease
C. Huntington's chorea
D. Multi-infarct dementia

Answer: B

15. A 30-year-old man presents to the clinic complaining of reduced energy, lack of enjoyment, reduced sleep, reduced appetite, suicidal ideas and lack of self-confidence. Examination and lab results are normal.
Which of the following is the most likely diagnosis?
A. Anxiety disorders
B. Depressive disorders
C. Drug-induced psychoses
D. Dissociative conversion disorder

Answer: B

16. A 28-year-old was brought to the Emergency Department complaining that his neighbors are trying to harm him and are planning to kill him. On questioning, he was very sure about this and said that he is still hearing their voices talking with each other, though none of them were present.
Which of the following is the most likely diagnosis?
A. Cognitive and perceptive disturbance
B. Depersonalization and derealization
C. Obsessive-compulsive disorder
D. Hallucinations and delusions

Answer: D

17. A 26-year-old woman presents to the Psychiatric Outpatient Clinic with a history of psychomotor excitation and irritability. She has grandiose delusions and auditory hallucinations. On a mental state examination, she exhibited a lack of insight and flight of ideas.
Which of the following disorders is the most likely diagnosis?
A. Neurosis
B. Psychosis
C. Personality
D. Dissociative conversion

Answer: B

18. A 32-year-old man presents to the clinic with a 2-year history of getting orgasm too readily during the sexual act. He complains that during sexual intercourse he is having palpitations and worrying thoughts in his mind all the time. On examination, he has normal findings.

Which of the following is the most likely diagnosis?
A. Anxiety disorder

B. Depressive disorder
C. Bipolar affective disorder
D. Primary erectile dysfunction

Answer: A

19. A 16-year-old girl presents for a follow-up visit complaining that she is unable to eat because of an intense desire to reduce her weight. Sometimes she develops an impulse to eat excessively and she then follows the impulse. She then intermittently self-induces vomiting. On examination, she looked weak.

Which of the following is the most likely diagnosis?
A. Bulimia nervosa
B. Anorexia nervosa
C. Personality disorder
D. Obsessive-compulsive disorder

Answer: B

20. A 20-year-old student presents to the Psychiatric Outpatient Clinic complaining that she feels embarrassed, dizzy, and tremulous whenever she tries to answer verbally in the classroom. Which of the following is the best treatment?
A. Small doses of alprazolam
B. Behavioral therapy
C. Relaxation training
D. Bupropion

Answer: B

21. A 21-year-old man with depression presents for a follow-up visit. During the consultation, the psychiatrist asked, "Do you think that you are mentally ill?"
Which of the following findings is the psychiatrist wants to assess?
A. Insight
B. Judgment
C. Compliance
D. Guilt feeling

Answer: A

22. A 23-year-old college student was brought to the Psychiatric Outpatient Clinic with a 3-month history of persecutory delusions, hearing voices commenting on his actions, and disorganized behavior without disturbance of mood, then returning to normal with no medication. Which of the following is the most likely diagnosis?
A. Schizophreniform disorder
B. Brief psychotic disorder
C. Delusional disorder
D. Schizophrenia

Answer: A

23. A 34-year-old man presents to the clinic for routine visit. During the consultation, he speaks loudly and rapidly without completely expressing one thought before beginning another. Speech patterns are rhyming and the patient is easily distracted by cues in the immediate surroundings. Which of the following is the most likely description for these findings?
A. Flight of ideas
B. Perseveration

C. Thought insertion
D. Loosening of associations

Answer: A

24. A 22-year-old man presents to the clinic with a history of approaching strangers and asking them the same inappropriate question without stopping when instructed to do so. The patient seemed unaware of the fact that this was an inappropriate behavior.
Which of the following is the most likely description for this behavior?
A. Flight of ideas
B. Perseveration
C. Thought insertion
D. Loosening of associations

Answer: B

25. A mother brings her 2-week-old infant to the Well-Baby Clinic for a check-up. During the visit, the physician found the child to be normal but the mother gives inappropriate answers to the questions. She appeared disoriented in time and date. She states that the child is deformed and unlikely to live long.

Which of the following is the most likely diagnosis?
A. Postnatal non-bonding
B. Postpartum psychosis
C. Neurosis
D. Fatigue

Answer: B

26. A 34-year-old woman presents to the clinic for postpartum care. She has experienced tearfulness and irritability during the 1 week following the delivery of her child. This episode resolved spontaneously and her mood has been normal since then.
Which of the following is the most likely diagnosis?
A. Maternity blues
B. Postnatal depression
C. Postpartum psychosis
D. Postnatal non-bonding

Answer: A

27. Which of the following conditions is most often associated with manic behavior?
A. Paranoid schizophrenia
B. Bipolar affective disorder
C. Grandiose mood disorder
D. Generalized anxiety disorder

Answer: B

28. A 21-year-old woman presents to the clinic with a history of changes in the voice that she hears. Until now, it had been merely making critical comments about everything she did. For the past 2 weeks, however, she thinks the voices seemed to have been able to "suck thoughts out" of her and send them to others.
Which of the following is the most likely diagnosis?
A. Mania
B. Agoraphobia
C. Schizophrenia
D. Mood disorder

Answer: C

29. A 26-year-old man presents to the clinic with a history of unusual behavior. He keeps a blanket over the television believing that the government uses the actors to spy through the TV. He believes that the instructions to cover the TV came from God who spoke through the light bulb.
Which of the following is the most likely diagnosis?

A. Schizophrenia
B. Agoraphobia
C. Dementia
D. Mania

Answer: A

30. A 25-year-old healthy man complains of episodes of chest tightness, palpitations, dizziness and breathlessness. These episodes resolved after about 5 minutes. He remains normal between episodes and the interval may extend for several days before a new episode occurs. He has had constant anxiety about having another attack and feels sad about the situation. The patient cannot identify a triggering mechanism.
Which of the following is the most likely diagnosis?

A. Generalized anxiety disorder
B. Major depressive disorder
C. Phobic reaction
D. Panic disorder

Answer: D

A 30-year-old man gives a history of pancreatic carcinoma and reports having lost all his prescriptions. He says that despite having recently completed chemotherapy he has developed new symptoms and is afraid of a recurrence. There are no medical records for him in the hospitals. When confronted with this information, the patient ran away from the clinic.

Which of the following is the most likely diagnosis?

A. Conversion disorder

B. Factitious disorder

C. Hypochondriasis

D. Malingering

Answer: B

A 40-year-old woman presents to the clinic with a 2-month history of a sleep disturbance, loss of interest, low energy, poor memory, and suicidal ideation. On examination, she looks weak and inactive. Laboratory investigations are normal.

Which of the following is the most likely diagnosis?

A. Dysthymia

B. Bipolar disorder

C. Major depression

D. Minor depression

Answer: C

A 19-year-old healthy man is brought to the clinic for evaluation of progressive erratic behaviour. Over the past 4 months, he has believed that people on television are reading his thoughts. He expressed the presence of supernatural powers and that he has begun to act out in response to these ideas. He has not caused injuries, but the trends are becoming more violent. On examination, he is normal.

Which of the following is the most likely diagnosis?

A. Schizophrenia

B. Conversion disorder

C. Rapid cycling depression

D. Manic phase of bipolar disorder

Answer: A

Which of the following is the most appropriate management plan for major depression?

A. Select any drug to start, as all have equal efficacy

B. Initiate therapy with one medication for 2 weeks

C. Stop medication with symptomatic improvement

D. Change medications if the patient doesn't respond

Answer: A

Which of the following systems is stimulated by stressful situations?

A. Urinary

B. Respiratory

C. Sympathetic

D. Parasympathetic

Answer: D

Stress leads to an increase in cortisol and catecholamine secretion that can alter the levels of lymphocyte production that fight disease.

Which of the following is the most likely explanation for these effects?

A. Interpretation of the seeing

B. Psychoneuroimmunology

C. Vaccination effect

D. Personal effects

Answer: B

Which of the following is the main difference between delirium and dementia?

A. Disturbance of consciousness

B. Multiple cognitive deficits

C. Memory impairment

D. Aphasia

Answer: A

A 72-year-old man experiences inability to sustain attention, disorganized thinking, hallucinations, and disorientation commencing 12 hours after an aortic femoral popliteal bypass. The symptoms fluctuated during the next 2 days.

Which of the following is the most likely diagnosis?

A. Multi-infarct dementia

B. Depression

C. Delirium

D. Mania

Answer: D

A 70-year-old man with Alzheimer's dementia becomes agitated and aggressive after developing hallucinations and delusions. A complete evaluation does not reveal any environmental, medical, or psychiatric cause, and behavioral interventions failed to improve the agitation.

Which of the following is the most appropriate next step in management?

A. Thioridazine

B. Haloperidol

C. Olanzapine

D. Buspirone

Answer: C

A 78-year-old man is brought to the Outpatient Clinic with a 2-year history of slow and progressive deterioration in his memory. He is able to perform some activities of daily living, including dressing and bathing, as well as cooking. He has personality changes, once a kind

and caring parent; he now displays periods of both agitation and aggression. Neurologic and musculoskeletal examinations cannot be carried out.

Which of the following is the best management?

A. Cost-effective laboratory investigation

B. Prescribe Risperidone for the patient

C. Admission to a chronic care facility

D. Refer to a geriatrician

Answer: A

A 30-year-old man presents to the Outpatient Clinic with manifestations of opioid withdrawal. After full investigations, he was diagnosed as a heroin addict. As part of the rehabilitation process, he was prescribed a substitute for the addictive drug.

Which of the following drugs should be prescribed?

A. Naloxone

B. Papaverine

C. Methadone

D. Dextromethorphan

Answer: C

Which of the following is described as "False belief not in accordance with a person's intelligence or culture"?

A. Illusion

B. Delusion

C. Hallucination

D. Somatization

Answer: B

A 65-year-old man with congestive heart failure and osteoarthritis moved into a nursing home 3 months ago. During the past 4 weeks, the patient has lost 3.6 kg, lost interest in all social activities, and has been crying almost every day. The mood is worse in the morning. At times, especially when the mood is impaired, the patient's short-term memory appears to be lost.

Which of the following is the most likely diagnosis?

A. Multi-infarct dementia

B. Alzheimer's disease

C. Hypothyroidism

D. Depression

Answer: D

A 19-year-old girl presents with a 12 kg weight loss over the last 6 months. She feels fat with an intense fear of gaining weight. Over the last 4 months, the patient has had amenorrhea. On examination, she has lanugo hair distribution.

Height 167 cm
Weight 42 kg

Which of the following is the most likely diagnosis?

A. Bulimia

B. Anorexia nervosa

C. Masked depression

D. Generalized anxiety disorder

Answer: B

A 29-year-old woman presents to the clinic with recurrent attacks of palpitations, chest pain, difficulty in breathing, and a fear of dying. She also trembles, experiences numbness and tingling sensations in the hands and feet. The past medical history is not significant. On examination, she looks tense, and the hands are cold with fine tremors.

Heart rate 108 /min, regular
Temperature 37.2° C
Blood pressure 120/82 mmHg
Respiratory rate 14 /min

Which of the following is the most likely diagnosis?

A. Pheochromocytoma

B. Hyperthyroidism

C. Panic disorder

D. Agoraphobia

Answer: C

A 29-year-old man presents with a 2-year history of intense fear before giving classes as a teacher in a secondary school. He claims that it is only a matter of time before a major teaching mistake is made.

Which of the following phobias is the most likely diagnosis?

A. Social

B. Mixed

C. Specific

D. Agoraphobia

Answer: A

A 55-year-old man presents to the clinic with a 2-month history of anxiety, irritability, impotence, difficulty in sleeping, and a weight loss of 10 kg. There is no evidence of marital disharmony or any external causes for the patient's anxiety. The physical examination was normal.

Which of the following is the most likely diagnosis?

A. Social phobia

B. Major depression

C. Secondary depression

D. Generalized anxiety disorder

Answer: B

Which of the following is the most likely a characteristic of a major depressive illness?

A. Hypomania

B. Late morning awakening

C. Hallucination and flight of ideas

D. Decreased eye contact in conversation

Answer: D

A 52-year-old woman presents with a 5-day history of insomnia and crying because of the death of a sibling. The patient has not slept for the last 2 days.

Which of the following is the most appropriate short course of therapy?

A. Fluoxetine

B. Imipramine

C. Lorazepam

D. Chlorpromazine

Answer: C

Which of the following factors indicates a poor prognosis for schizophrenia?

A. Acute onset of psychosis

B. Onset during adolescence

C. Family history of schizophrenia

D. Anxiety during psychotic episodes

Answer: B

A 72-year-old man who lost his spouse 2 months ago after a prolonged illness presents with a sad mood, decreased appetite, and sleep disturbance.

Which of the following is the most likely diagnosis?

A. Dysthymia

B. Bereavement

C. Major depression

D. Minor depression

Answer: B

Which of the following drug mechanisms of action is most appropriate to treat obsessional behavioral disorder?

A. Blocks serotonin production

B. Increases the availability of serotonin

C. Increases the production of serotonin

D. Decreases the availability of serotonin

Answer: B

Which of the following medications causes insomnia, anxiety, and restlessness?

A. Tricyclic antidepressant

B. Tetracyclic Antidepressant

C. Monoamine oxidase inhibitor

D. Selective serotonin-reuptake inhibitor

Answer: D

A 76-year-old man newly diagnosed with depression started on 30 mg amitriptyline each night. Shortly after the beginning of treatment, he complains of dizziness every morning.

Which of the following is the most appropriate next step in management?

A. Change to a selective serotonin reuptake inhibitor

B. Advise to take the medication with food

C. Advice to take the medication in the morning

D. Change amitriptyline dose to 10 mg 3 times a day

Answer: D

Which of the following is a potential side effect of amitriptyline?

A. Dystonia

B. Weight gain

C. Hypersalivation

D. Hyperpigmentation

Answer: B

A 12-year-old obese boy presents to the Primary Care Clinic with a history of being teased by children at school and, as a result, has considered "taking a lot of pills to just go to sleep and not wake up."

Which of the following is the most appropriate action?

A. Weight loss advice

B. Reassuring the patient

C. Immediate psychiatry referral

D. Prescription of a fast-acting antidepressant

Answer: C

A 76-year-old man presents to the Psychiatry Clinic for follow-up accompanied by his son. He has progressive memory loss, difficulty in finding words, wandering, and agitation, as well as urinary and fecal incontinence for the last 3 years. On examination, he was conscious but not oriented in time and place; his short memory was severely impaired with an intact long memory (see report).

Brain CT scan:

Atrophied hippocampal and temporal lobes.

Which of the following chromosomes harbors the mutated gene is responsible for this presentation?

A. 13

B. 15

C. 21

D. X

Answer: C

Which of the following clinical signs caused by a lesion in Wernicke's area?

A. Can understand but unable to say words easily

B. Cannot understand spoken or written words

C. Loss of ability to recognize faces

D. Loss of short-term memory

Answer: B

A 55-year-old woman presents to the Primary Health Care Centre complaining of loss of interest and pleasure in daily activities for 3 weeks. She also complains that she has difficulty in concentrating and sleeping. Further, she has also expressed that she feels that she is worthless in the world and therefore wants to end her life.

Which of the following biochemical substance deficiency is the most likely cause?

A. Aspartate

B. Serotonin

C. Glutamate

D. Acetylcholine

Answer: B

A 55-year-old woman presents to the Primary Health Care Centre complaining of loss of interest and pleasure in daily activities for 3 weeks. She also complains that she has difficulty in concentrating and sleeping. Further, she has also expressed that she feels that she is worthless in the world and therefore wants to end her life.

Which of the following reuptake inhibitors is the best treatment?

A. DOPA

B. GABA

C. Serotonin

D. Acetylcholine

Answer: C

Which of the following drugs used in Alzheimer's disease is most likely to cause hepatotoxicity?

A. Tacrine

B. Donepezil

C. Rivastigmine

D. Galantamine

Answer: A

Which of the following antipsychotic medications is most likely to cause weight gain?

A. Clozapine

B. Olanzapine

C. Risperidone

D. Ziprasidone

Answer: B

A 35-year-old depressed housewife was started with antidepressants. After 2 weeks of taking the medication, the patient started complaining of constipation.

Which of the following groups of antidepressants is the most likely cause?

A. Tricyclic antidepressants

B. Monoamine oxidase inhibitors

C. Selective serotonin reuptake inhibitors

D. Selective nor-adrenaline and serotonin inhibitors

Answer: A

Which of the following conditions is the most likely cause of dementia?

A. Spinocerebellar ataxia

B. Vitamin B12 deficiency

C. Idiopathic parkinsonism

D. Amyotrophic lateral sclerosis

Answer: C

A 15-year-old girl who had a serious disagreement with her friend. She did not want to meet her friend or talk to her anymore. If the girl saw her friend coming towards her or sitting in a room, she would move away and not go in that direction.

Which of the following defense mechanisms is the most likely explanation for this behavior?

A. Denial

B. Acting out

C. Avoidance

D. Repression

Answer: C

A 65-year-old man with depression presents to the clinic for a follow-up visit. The physician did his consultation using a bio-psycho-social model of treatment.

Which of the following is a characteristic of this treatment model?

A. Evidence-based medicine

B. Traditional allopathy

C. Geriatric medicine

D. Holistic medicine

Answer: D

A 36-year-old diabetic woman presents to the clinic for a follow-up visit, and after an initial assessment, the physician wished to perform a physical examination. He first asked her permission, which was granted.

Which of the following ethical measures was the physician adopting?

A. Obtain informed consent

B. Respect for the patient

C. Fulfill his responsibility

D. Show his efficiency

Answer: A

A 40-year-old diabetic man was treated appropriately with small doses of insulin. On the next follow-up visit, he bought a very expensive watch as a gesture of gratitude to his physician.

Which of the following is the most appropriate response?

A. Accept and advise not to repeat

B. Refuse and give a warning

C. Politely refuse the gift

D. Accept from the rich

Answer: C

A 36-year-old newly diagnosed hypertensive man presents for follow-up visit. After a thorough assessment, the physician discussed with him his diagnosis, the etiology of his illness, the probable side effects of the prescribed medicine, and the prognosis of his illness and finally the restrictions the patient might experience due to the illness.

Which of the following professional skills is the physician used?
A. Developing rapport
B. Improving communication
C. Providing informational care
D. Developing bonding with the patient

Answer: C

Which of the following is the most useful communication technique when taking a patient history?
A. Humor
B. Silent listening
C. Specific questions

D. Open-ended questions

Answer: D

Which of the following is an example of the open-ended questions in consultation?
A. "Tell me about the pain."
B. "Tell me about the pain in your chest."
C. "Point to the area of pain in your chest."
D. "Have you seen a doctor in these 76 months?"

Answer: A

A 6-year-old boy is brought to the clinic by his mother. She noticed that the child has been eating clay and paper for the last 4 months.

Which of the following is the most appropriate initial treatment?
A. Fluoxetine
B. Imipramine
C. Clonazepam
D. Behavioral therapy

Answer: D

A 34-year-old woman presents to the clinic with a 6-month history of extreme fear of going to the park, the zoo, and sporting events. This has worsened to the point where she has stopped going out of the house.

Which of the following is the most likely diagnosis?
A. Conversion disorder
B. Schizophrenia
C. Social phobia
D. Agoraphobia

Answer: D

A 45-year-old man presents in an anxious, tearful state. He complains of chronic, upper abdominal pain and is certain that he has stomach cancer, but no one believes him. He has seen 6 different physicians in the past year and has had a GI series and barium enema performed, an ultrasound of the abdomen twice, and a CT scan. The patient was thinking of next going to a Herbal Clinic.

Which of the following is the most likely diagnosis?
A. Somatoform pain disorder
B. Somatization disorder
C. Conversion disorder
D. Hypochondriasis

Answer: D

A 43-year-old woman delivered a full-term infant. On examination, he has epicanthic folds, delayed milestones, flat facial profile, simian palmar creases, and an atrial septal defect. Which of the following is the most likely diagnosis?
A. Trisomy 12
B. Trisomy 21
C. Di-George syndrome
D. Robertsonian translocation

Answer: B

A 16-year-old girl is brought by her father to the Emergency Department with complaints of tachypnea, light-headedness, paraesthesia, and palpitations. Her father gives the history that she has recently failed her math test

Which of the following is the most likely diagnosis?
A. Oppositional defiant disorder
B. Hyperventilation syndrome
C. Munchausen syndrome
D. Anxiety disorder

Answer: B

A 27-year-old man presents to the clinic with a 1-month history of low mood and lack of sleep. It was associated with weight gain, isolation, and concentration impairment. His wife mentioned that he bought a very expensive car and thinking that he is a wealthy man 3 months ago. He is very talkative and difficult to interrupt. Which of the following medications is the most effective for long-term management?
A. Lithium
B. Bupropion
C. Olanzapine
D. Mirtazapine

Answer: A

Which of the following is the diagnostic criteria for Attention deficit hyperactivity disorder in children, according to the International Classification of Diseases-10?
A. 1 inattentive, 3 hyperactive, and 1 impulsive symptoms
B. 1 inattentive, 3 hyperactive, and 3 impulsive symptoms
C. 3 inattentive, 3 hyperactive, and 1 impulsive symptoms
D. 6 inattentive, 3 hyperactive, and 1 impulsive symptoms

Answer: D

A 23-year-old woman with frequent panic attacks presents to clinic complains that during her last attack 3 days ago she felt that she was disconnected from the world, as if the environment is strange.

Which of the following would best describe her feelings?
A. Illusion
B. Derealization
C. Dull perception
D. Depersonalization

Answer: B

A 30-year-old man has a personality disorder, which is characterized by being aggressive and violent.

Which of the following is most likely associated with violence?
A. Low levels of serotonin
B. High levels of serotonin
C. Low levels of endorphins
D. High levels of endorphins

Answer: A

Which of the following is associated with dyslexia?

A. Abnormal gait
B. Brainstem defects
C. Verbal language difficulties
D. Defect in visual or hearing acuity

Answer: C

Which of the following is the most likely clinical presentation of delusional patients?
A. Disturbance in perception
B. Rare in manic disorders
C. Found only in schizophrenia
D. Disturbance of thought content

Answer: D

A 7-year-old boy is brought by his mother because he was unable to sit in school without interrupting the class, wandering around the classroom or poking his neighbors. On examination, he has normal growth parameters.

Which of the following is the best management?
A. Bupropion
B. Paroxetine
C. Guanfacine
D. Dextroamphetamine

Answer: D

A 30-year-old woman presents with a complaint of lower back pain for 1 year. During the last 2 years, she had presented with recurrent abdominal pain, bloating, headache, and migratory joint pain. Her menstrual periods are irregular and heavy. She was terminated twice from the job due to frequent absence from work. She is upset that proper diagnosis or treatment was not given to her. She had multiple laboratory tests done during the last 2 years and all were normal. The physical examination is unremarkable.

Which of the following is the most likely diagnosis?
A. Malingering
B. Hypochondriasis
C. Factitious disorder
D. Somatization disorder

Answer: D

A 23-year-old man with bronchial asthma presents with a 5-year history of severe anxiety in social gatherings. He often has sweating and palpitation when he leads discussions at work. He prefers to sit alone at home. He notices that his presenting complaint is provoked by salbutamol. His physical examination is unremarkable.

Which of the following is the most appropriate therapy?
A. Sertraline
B. Bupropion
C. Alprazolam
D. Propranolol

Answer: A

A 28-year-old woman with bipolar disorder, treated for 6 months, presents for a follow-up visit. On examination, she has 8 kg weight gain, generalized weakness, and hair loss.

Which of the following is the most likely cause?
A. Hypothalamic disorder
B. Hypothyroidism
C. Verapamil
D.Lithium

Answer: D

A 56-year-old man with schizophrenia is started on neuroleptic medications to treat his psychotic symptoms.

Which of the following side effects is the patient at greatest risk?
A. Seizures
B. Akathisia
C. Hyperthermia
D. Myocardial infarction

Answer: B

Which of the following is the most likely benefit of using antipsychotic therapy in schizophrenic patients?
A. Improve apathy
B. Reduce delusions
C. Improve thinking ability
D. Increase social interaction

Answer: B

A 23-year-old man is brought to the psychiatrist clinic because for the first time he started to hear some voices talking to him with unknown sources. On examination, he was apathetic and emotionally withdrawn.

Which of the following is the best management?
A. Olanzapine
B. Fluoxetine
C. Cognitive behavioral therapy
D. Antipsychotic and cognitive behavioral therapy

Answer: D

A 45-year-old man with recurrent episodes of schizophrenia is on maintenance therapy to maintain suppression of symptoms, prevent relapse, improve quality of life, and support engagement in psychosocial therapy.

Which of the following is the best outcome following the maintenance therapy?
A. Complete resolution
B. 5% will achieve remission
C. 70% will have a satisfactory quality of life
D. 33% obtain long-lasting symptom reduction

Answer: D

Which of the following selective serotonin reuptake inhibitors is the most toxic in overdose?
A. Sertraline
B. Citalopram
C. Paroxetine
D. Fluoxetine

Answer: B

A 23-year-old woman is newly diagnosed with paranoid schizophrenia based on the presence of both negative and positive symptoms of schizophrenia.

Which of the following is the most effective medication?
A. Clozapine
B. Trazodone
C. Fluphenazine
D. Chlorpromazine

Answer: A

Which of the following is the lowest fatal dose of tricyclic antidepressant?
A. 500 mg
B. 1 gm
C. 2 gm
D. 3 gm

Answer: D