Burns

Check **ABCD & Exposure:** expose the body to look for burns

- Assess **affected area** (rule of nine)
- Assess **depth of burn** superficial (1st degree), partial (2nd degree) or full thickness (3rd degree)



Percentual relativo da área da superfície do corpo (% ASC) afetada pelo crescimento

Parte do corpo	Idade 0 ano	1 ano	5 anos	10 anos	15 anos
a = 1/2 da cabeça	9 1/2	8 1/2	6 1/2	5 1/2	4 1/2
b = 1/2 da coxa	2 3/4	3 1/4	4	4 1/4	4 1/2
c = 1/2 da perna inferior	2 1/2	2 1/2	2 3/4	3	3 1/4

Call ambulance and **refer** to burn center in:

- Partial thickness **> 10% of TBSA.**
- Burn **in face, hands, genitalia, perineum or major joints**
- **Third degree burns** in any age group.
- **Electrical, chemical and inhalational burns**
- **Burns associated with trauma** (e.g. fracture)

Management:

- **Irrigate** with running **cool water**
- **Adults with > 15% of TBSA require IV fluids (Parkland formula)**
- **Leave Blisters intact** to decrease risk of infection
- **Dressing** with non-adhesive gauze padding
- **T.Toxoid** (if last shot >5 years)
- **Analgesia** (paracetamol, NSAIDs)
- Examine dressing after **48 hours (2nd and 3rd degree)**
- **Change Dressing for 2nd degree burn after 3-5 days** in absence of infection