


A Strong linker contrast



median $|\Delta \log q|$
0.812

WC concord.
5/5

Selectivity
4/4


A mean metal REPEAT q/e: Zn +1.06

B mean metal REPEAT q/e: Zn +1.01

robust node | REPEAT W1 all/hetero = 0.110/0.211 e

Large linker contrast with process agreement.

B Strong metal contrast



median $|\Delta \log q|$
0.307

WC concord.
5/5

Selectivity
4/4


A mean metal REPEAT q/e: Zn +1.30

B mean metal REPEAT q/e: Cu +0.78

robust coordination | REPEAT W1 all/hetero = 0.082/0.116 e

Large metal-identity contrast with process agreement.

C Cu–Zn pressure exception



median $|\Delta \log q|$
0.082

WC concord.
5/5

Selectivity
4/4


A mean metal REPEAT q/e: Zn +1.33

B mean metal REPEAT q/e: Cu +0.99

method-sensitive: 8 sites | REPEAT W1 = 0.052/0.038 e

Pressure-exception case within the Cu/Zn set.

D Near-null linker contrast



median $|\Delta \log q|$
0.012

WC concord.
5/5

Selectivity
3/4


A mean metal REPEAT q/e: Cu +1.04

B mean metal REPEAT q/e: Cu +1.03

near-null | REPEAT W1 all/hetero = 0.015/0.007 e

Matched comparison with minimal adsorption separation.

E Process-discordant contrast



median $|\Delta \log q|$
0.064

WC concord.
1/5

Selectivity
1/4


A mean metal REPEAT q/e: Zn +1.04

B mean metal REPEAT q/e: Zn +1.04

descriptive context | REPEAT W1 = 0.048/0.064 e

Process translation breaks alignment.

F Exploratory functional motif



median $|\Delta \log q|$
0.060

WC concord.
4/5

Selectivity
4/4

A mean metal REPEAT q/e: V +1.34

B mean metal REPEAT q/e: V +1.35

exploratory | REPEAT W1 all/hetero = 0.019/0.041 e

Exploratory only; not class-level evidence.