

Aggregatable Subvector Commitments for Stateless Cryptocurrencies

Alin Tomescu¹
[@alinush407](#)

Justin Drake²
[@drakejustin](#)

Ittai Abraham¹
[@ittaia](#)

Dankrad Feist²
[@dankrad](#)

Vitalik Buterin²
[@VitalikButerin](#)


Dmitry Khovratovich²
[@Khovr](#)

¹VMware Research, ²Ethereum Foundation

September 14th, 2020


Motivation

Miners rely on **state** to validate transactions and blocks.



Motivation

Miners rely on **state** to validate transactions and blocks.




Motivation

Miners rely on **state** to validate transactions and blocks.



Motivation

Miners rely on **state** to validate transactions and blocks.



Validation state can be **very large**:

Motivation

Miners rely on **state** to validate transactions and blocks.




Validation state can be **very large**:

- Hundreds of GBs in Ethereum

Motivation

Miners rely on **state** to validate transactions and blocks.



Validation state can be **very large**:

- Hundreds of GBs in Ethereum
- GBs in Bitcoin

Motivation

Miners rely on **state** to validate transactions and blocks.




Validation state can be **very large**:

- Hundreds of GBs in Ethereum
- GBs in Bitcoin

This poses scalability challenges:

Motivation

Miners rely on **state** to validate transactions and blocks.



Validation state can be **very large**:


- Hundreds of GBs in Ethereum
- GBs in Bitcoin

This poses scalability challenges:

- Consensus via sharding

Motivation

Miners rely on **state** to validate transactions and blocks.



Validation state can be **very large**:

- Hundreds of GBs in Ethereum
- GBs in Bitcoin

This poses scalability challenges:

- Consensus via sharding
- Barrier to entry for P2P nodes

Motivation

Miners rely on **state** to validate transactions and blocks.



Validation state can be **very large**:

- Hundreds of GBs in Ethereum
- GBs in Bitcoin

This poses scalability challenges:

- Consensus via sharding
- Barrier to entry for P2P nodes
- DoS attacks


Previous Work Outsources State via *Vector Commitments (VCs)*

[CPZ18] was first to proposed stateless validation using VCs!




Previous Work Outsources State via *Vector Commitments (VCs)*

[CPZ18] was first to proposed stateless validation using VCs!




Previous Work Outsources State via *Vector Commitments (VCs)*

[CPZ18] was first to proposed stateless validation using VCs!




Previous Work Outsources State via *Vector Commitments* (VCs)

[CPZ18] was first to proposed stateless validation using VCs!




Previous Work Outsources State via *Vector Commitments* (VCs)

[CPZ18] was first to proposed stateless validation using VCs!



Previous Work Outsources State via *Vector Commitments* (VCs)

[CPZ18] was first to proposed stateless validation using VCs!



Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters
 - Keeps trusted setup simple

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters
 - Keeps trusted setup simple
 - Keeps parameters *updatable*

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters
 - Keeps trusted setup simple
 - Keeps parameters *updatable*
- Subtleties of VC-based stateless cryptocurrencies

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters
 - Keeps trusted setup simple
 - Keeps parameters *updatable*
- Subtleties of VC-based stateless cryptocurrencies
 - Keeping track of transaction counters

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters
 - Keeps trusted setup simple
 - Keeps parameters *updatable*
- Subtleties of VC-based stateless cryptocurrencies
 - Keeping track of transaction counters
 - Verifiable update keys

Our contributions: **Efficient** VC from KZG polynomial commitments [KZG10]

- Constant-sized, aggregatable and updatable proofs
- Constant-sized **update keys**
- Quasilinear-time proof pre-computation, via Feist-Khovratovich (FK) technique [FK20]
- **Previous work** is either: non-aggregatable, or has linear-sized update keys, or is based on less efficient, hidden-order groups

Additionally,

- New security definition for KZG batch proofs, proven secure under n -SBDH
- *Publicly-derivable* parameters from “powers-of-tau” parameters
 - Keeps trusted setup simple
 - Keeps parameters *updatable*
- Subtleties of VC-based stateless cryptocurrencies
 - Keeping track of transaction counters
 - Verifiable update keys
 - DoS attacks on new user registration

Thank you!

Paper is too long? **Read our blogpost!**

<https://alinush.github.io/2020/05/06/aggregatable-subvector-commitments-for-stateless-cryptocurrencies.html>

Appendix

Outline

Appendix

Previous Work

Background

Kate-Zaverucha-Goldberg (KZG) Polynomial Commitments

VCs from KZG Commitments to Lagrange Polynomials

Our Techniques

Updating Proofs (Case $i = j$)

Updating Proofs (Case $i \neq j$)

Aggregating Proofs into Subvector Proofs

Extras

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Table 1: Asymptotic comparison to previous (aS)VCs. n is the size of the vector \vec{v} .

(aS)VC scheme	Public parameters	Proof size	Update key size	Digest update	Aggr. b proofs	Prove all
---------------	-------------------	------------	-----------------	---------------	------------------	-----------

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Table 1: Asymptotic comparison to previous (aS)VCs. n is the size of the vector \vec{v} .

(aS)VC scheme	Public parameters	Proof size	Update key size	Digest update	Aggr. b proofs	Prove all
Merkle trees [Mer88]	1	$\log n$	\times	\times	\times	n

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Table 1: Asymptotic comparison to previous (aS)VCs. n is the size of the vector \vec{v} .

(aS)VC scheme	Public parameters	Proof size	Update key size	Digest update	Aggr. b proofs	Prove all
Merkle trees [Mer88]	1	$\log n$	\times	\times	\times	n
CDHK [CDHK15]	n	1	n	✓	\times	n^2
CPZ [CPZ18]	n	$\log n$	$\log n$	✓	\times	$n \log n$
TCZ [TCZ ⁺ 20, Tom20]	n	$\log n$	$\log n$	✓	\times	$n \log n$
Pointproofs [GRWZ20]	n	1	n	✓	$b_{\mathbb{G}}$	n^2

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Table 1: Asymptotic comparison to previous (aS)VCs. n is the size of the vector \vec{v} .

(aS)VC scheme	Public parameters	Proof size	Update key size	Digest update	Aggr. b proofs	Prove all
Merkle trees [Mer88]	1	$\log n$	\times	\times	\times	n
CDHK [CDHK15]	n	1	n	✓	\times	n^2
CPZ [CPZ18]	n	$\log n$	$\log n$	✓	\times	$n \log n$
TCZ [TCZ ⁺ 20, Tom20]	n	$\log n$	$\log n$	✓	\times	$n \log n$
Pointproofs [GRWZ20]	n	1	n	✓	$b_{\mathbb{G}}$	n^2
BBF [BBF19]	1	$1_{\mathbb{G}_?}$	\times	\times	$b \log n_{\mathbb{G}_?}$	$n \log n_{\mathbb{G}_?}$
CFG ₁ [CFG ⁺ 20]	1	$1_{\mathbb{G}_?}$	\times	\times	$b \log b \log n_{\mathbb{G}_?}$	$n \log^2 n_{\mathbb{G}_?}$

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Table 1: Asymptotic comparison to previous (aS)VCs. n is the size of the vector \vec{v} .

(aS)VC scheme	Public parameters	Proof size	Update key size	Digest update	Aggr. b proofs	Prove all
Merkle trees [Mer88]	1	$\log n$	\times	\times	\times	n
CDHK [CDHK15]	n	1	n	✓	\times	n^2
CPZ [CPZ18]	n	$\log n$	$\log n$	✓	\times	$n \log n$
TCZ [TCZ ⁺ 20, Tom20]	n	$\log n$	$\log n$	✓	\times	$n \log n$
Pointproofs [GRWZ20]	n	1	n	✓	$b_{\mathbb{G}}$	n^2
BBF [BBF19]	1	$1_{\mathbb{G}_?}$	\times	\times	$b \log n_{\mathbb{G}_?}$	$n \log n_{\mathbb{G}_?}$
CFG ₁ [CFG ⁺ 20]	1	$1_{\mathbb{G}_?}$	\times	\times	$b \log b \log n_{\mathbb{G}_?}$	$n \log^2 n_{\mathbb{G}_?}$
CFG ₂ [CF13, LM19, CFG ⁺ 20]	1	$1_{\mathbb{G}_?}$	$1_{\mathbb{G}_?}$	✓	$b \log^2 b_{\mathbb{G}_?}$	$n \log^2 n_{\mathbb{G}_?}$

Aggregatable Subvector Commitments (aSVCs) versus Previous Work

Table 1: Asymptotic comparison to previous (aS)VCs. n is the size of the vector \vec{v} .

(aS)VC scheme	Public parameters	Proof size	Update key size	Digest update	Aggr. b proofs	Prove all
Merkle trees [Mer88]	1	$\log n$	\times	\times	\times	n
CDHK [CDHK15]	n	1	n	✓	\times	n^2
CPZ [CPZ18]	n	$\log n$	$\log n$	✓	\times	$n \log n$
TCZ [TCZ ⁺ 20, Tom20]	n	$\log n$	$\log n$	✓	\times	$n \log n$
Pointproofs [GRWZ20]	n	1	n	✓	$b_{\mathbb{G}}$	n^2
BBF [BBF19]	1	$1_{\mathbb{G}_?}$	\times	\times	$b \log n_{\mathbb{G}_?}$	$n \log n_{\mathbb{G}_?}$
CFG ₁ [CFG ⁺ 20]	1	$1_{\mathbb{G}_?}$	\times	\times	$b \log b \log n_{\mathbb{G}_?}$	$n \log^2 n_{\mathbb{G}_?}$
CFG ₂ [CF13, LM19, CFG ⁺ 20]	1	$1_{\mathbb{G}_?}$	$1_{\mathbb{G}_?}$	✓	$b \log^2 b_{\mathbb{G}_?}$	$n \log^2 n_{\mathbb{G}_?}$
Our aSVC	n	1	1	✓	$b \lg^2 b_{\mathbb{F}} + b_{\mathbb{G}}$	$n \log n$

Some limitations

Unlike schemes based on hidden-order groups [CF13, LM19, BBF19, CFG⁺20], we have:

Some limitations

Unlike schemes based on hidden-order groups [CF13, LM19, BBF19, CFG⁺20], we have:

- Trusted setup

Some limitations

Unlike schemes based on hidden-order groups [CF13, LM19, BBF19, CFG⁺20], we have:

- Trusted setup
- No *incremental* aggregation & no dis-aggregation [CFG⁺20]

Some limitations

Unlike schemes based on hidden-order groups [CF13, LM19, BBF19, CFG⁺20], we have:

- Trusted setup
- No *incremental* aggregation & no dis-aggregation [CFG⁺20]
- No space-time trade-off for proof pre-computation [BBF19, CFG⁺20]

Outline

Appendix

Previous Work

Background

Kate-Zaverucha-Goldberg (KZG) Polynomial Commitments

VCS from KZG Commitments to Lagrange Polynomials

Our Techniques

Updating Proofs (Case $i = j$)

Updating Proofs (Case $i \neq j$)

Aggregating Proofs into Subvector Proofs

Extras

KZG Constant-sized Polynomial Commitments [KZG10]

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

$$c(\phi) = g^{\phi(\tau)} \tag{1}$$

(2)

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

$$c(\phi) = g^{\phi(\tau)} \tag{1}$$

$$= (g^{\tau^n})^{\phi_n} (g^{\tau^{n-1}})^{\phi_{n-1}} \dots (g^{\tau})^{\phi_1} (g)^{\phi_0} \tag{2}$$

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

$$c(\phi) = g^{\phi(\tau)} \quad (1)$$

$$= (g^{\tau^n})^{\phi_n} (g^{\tau^{n-1}})^{\phi_{n-1}} \dots (g^{\tau})^{\phi_1} (g)^{\phi_0} \quad (2)$$

Homomorphism: For all field elements a, b and polynomials $\phi(X), \psi(X)$, we have:

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

$$c(\phi) = g^{\phi(\tau)} \quad (1)$$

$$= (g^{\tau^n})^{\phi_n} (g^{\tau^{n-1}})^{\phi_{n-1}} \dots (g^{\tau})^{\phi_1} (g)^{\phi_0} \quad (2)$$

Homomorphism: For all field elements a, b and polynomials $\phi(X), \psi(X)$, we have:

$$c(a \cdot \phi + b \cdot \psi) = g^{a \cdot \phi(\tau) + b \cdot \psi(\tau)} \quad (3)$$

(4)

(5)

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

$$c(\phi) = g^{\phi(\tau)} \quad (1)$$

$$= (g^{\tau^n})^{\phi_n} (g^{\tau^{n-1}})^{\phi_{n-1}} \dots (g^{\tau})^{\phi_1} (g)^{\phi_0} \quad (2)$$

Homomorphism: For all field elements a, b and polynomials $\phi(X), \psi(X)$, we have:

$$c(a \cdot \phi + b \cdot \psi) = g^{a \cdot \phi(\tau) + b \cdot \psi(\tau)} \quad (3)$$

$$= (g^{\phi(\tau)})^a (g^{\psi(\tau)})^b \quad (4)$$

$$(5)$$

KZG Constant-sized Polynomial Commitments [KZG10]

Fix n -SDH **public parameters** $(g^{\tau^i})_{0 \leq i \leq n}$ such that **trapdoor** $\tau \in \mathbb{F}_p$ is unknown.

Committing: Given $\phi \in \mathbb{F}_p[X]$, where $\phi(X) = \sum_{i=0}^n \phi_i X^i = \langle \phi_0, \phi_1, \dots, \phi_n \rangle$ of degree $\leq n$:

$$c(\phi) = g^{\phi(\tau)} \quad (1)$$

$$= (g^{\tau^n})^{\phi_n} (g^{\tau^{n-1}})^{\phi_{n-1}} \dots (g^{\tau})^{\phi_1} (g)^{\phi_0} \quad (2)$$

Homomorphism: For all field elements a, b and polynomials $\phi(X), \psi(X)$, we have:

$$c(a \cdot \phi + b \cdot \psi) = g^{a \cdot \phi(\tau) + b \cdot \psi(\tau)} \quad (3)$$

$$= (g^{\phi(\tau)})^a (g^{\psi(\tau)})^b \quad (4)$$

$$= c(\phi)^a c(\psi)^b \quad (5)$$

VCs from Lagrange Basis Polynomials [CDHK15]

VCs from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

Vectors from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \tag{6}$$

(7)

VCs from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (6)$$

$$L_i(X) = \prod_{\substack{j \in [0, n) \\ j \neq i}} \frac{X - j}{i - j} \quad (7)$$

VCs from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (6)$$

$$L_i(X) = \prod_{\substack{j \in [0, n) \\ j \neq i}} \frac{X - j}{i - j} \quad (7)$$

Applying the KZG homomorphism to Equation (6):

VCS from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (6)$$

$$L_i(X) = \prod_{\substack{j \in [0, n) \\ j \neq i}} \frac{X - j}{i - j} \quad (7)$$

Applying the KZG homomorphism to Equation (6):

$$c(\phi) = \prod_{i=0}^{n-1} c(L_i)^{v_i} \quad (8)$$

VCs from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (6)$$

$$L_i(X) = \prod_{\substack{j \in [0, n) \\ j \neq i}} \frac{X - j}{i - j} \quad (7)$$

Applying the KZG homomorphism to Equation (6):

$$c(\phi) = \prod_{i=0}^{n-1} c(L_i)^{v_i} \quad (8)$$

Note: Public parameters include commitments $c(L_i)$.

VCs from Lagrange Basis Polynomials [CDHK15]

Represent vector \vec{v} with a polynomial ϕ s.t. $\phi(i) = v_i$:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (6)$$

$$L_i(X) = \prod_{\substack{j \in [0, n) \\ j \neq i}} \frac{X - j}{i - j} \quad (7)$$

Applying the KZG homomorphism to Equation (6):

$$c(\phi) = \prod_{i=0}^{n-1} c(L_i)^{v_i} \quad (8)$$

Note: Public parameters include commitments $c(L_i)$. Can derive from g^{T^i} 's.

Updating the Digest = Updating Polynomial & Updating Commitment

Updating the Digest = Updating Polynomial & Updating Commitment

Assume v_i changed to $v_i + \delta_i$.

Updating the Digest = Updating Polynomial & Updating Commitment

Assume v_i changed to $v_i + \delta_i$. Old polynomial was:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (9)$$

Updating the Digest = Updating Polynomial & Updating Commitment

Assume v_i changed to $v_i + \delta_i$. Old polynomial was:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (9)$$

Updated polynomial will be:

$$\phi'(X) = \phi(X) + \delta_i L_i(X) \quad (10)$$

Updating the Digest = Updating Polynomial & Updating Commitment

Assume v_i changed to $v_i + \delta_i$. Old polynomial was:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (9)$$

Updated polynomial will be:

$$\phi'(X) = \phi(X) + \delta_i L_i(X) \quad (10)$$

Updated commitment will be:

$$c(\phi') = c(\phi) \cdot c(L_i)^{\delta_i} \quad (11)$$

Updating the Digest = Updating Polynomial & Updating Commitment

Assume v_i changed to $v_i + \delta_i$. Old polynomial was:

$$\phi(X) = \sum_{i=0}^{n-1} v_i \cdot L_i(X) \quad (9)$$

Updated polynomial will be:

$$\phi'(X) = \phi(X) + \delta_i L_i(X) \quad (10)$$

Updated commitment will be:

$$c(\phi') = c(\phi) \cdot c(L_i)^{\delta_i} \quad (11)$$

Thus, for our purposes, each upk_i will include $c(L_i)$.

Proof π_i for v_i Refresher

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i$

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i \Leftrightarrow \phi \bmod (X - i) = v_i$.

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i \Leftrightarrow \phi \bmod (X - i) = v_i$.

$$q_i(X) = \frac{\phi(X) - v_i}{X - i} \tag{12}$$

(13)

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i \Leftrightarrow \phi \bmod (X - i) = v_i$.

$$q_i(X) = \frac{\phi(X) - v_i}{X - i} \quad (12)$$

$$\pi_i = c(q_i) = g^{q_i(\tau)} \quad (13)$$

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i \Leftrightarrow \phi \bmod (X - i) = v_i$.

$$q_i(X) = \frac{\phi(X) - v_i}{X - i} \quad (12)$$

$$\pi_i = c(q_i) = g^{q_i(\tau)} \quad (13)$$

To verify, use pairings:

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i \Leftrightarrow \phi \bmod (X - i) = v_i$.

$$q_i(X) = \frac{\phi(X) - v_i}{X - i} \quad (12)$$

$$\pi_i = c(q_i) = g^{q_i(\tau)} \quad (13)$$

To verify, use pairings:

$$e(c(\phi) / g^{v_i}, g) = e(\pi_i, g^\tau / g^i) \Leftrightarrow \quad (14)$$

$$(15)$$

Proof π_i for v_i Refresher

A proof π_i for v_i must convince that $\phi(i) = v_i \Leftrightarrow \phi \bmod (X - i) = v_i$.

$$q_i(X) = \frac{\phi(X) - v_i}{X - i} \quad (12)$$

$$\pi_i = c(q_i) = g^{q_i(\tau)} \quad (13)$$

To verify, use pairings:

$$e(c(\phi) / g^{v_i}, g) = e(\pi_i, g^\tau / g^i) \Leftrightarrow \quad (14)$$

$$\phi(\tau) - v_i = q_i(\tau)(\tau - i) \quad (15)$$

Outline

Appendix

Previous Work

Background

Kate-Zaverucha-Goldberg (KZG) Polynomial Commitments

VCS from KZG Commitments to Lagrange Polynomials

Our Techniques

Updating Proofs (Case $i = j$)

Updating Proofs (Case $i \neq j$)

Aggregating Proofs into Subvector Proofs

Extras

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

(17)

(18)

(19)

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$(18)$$

$$(19)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_i(L_i(X) - 1)}{X - i} \quad (18)$$

$$(19)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_i (L_i(X) - 1)}{X - i} \quad (18)$$

$$q'_i(X) = q_i(X) + \delta_i \left(\frac{L_i(X) - 1}{X - i} \right) \quad (19)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_i (L_i(X) - 1)}{X - i} \quad (18)$$

$$q'_i(X) = q_i(X) + \delta_i \left(\frac{L_i(X) - 1}{X - i} \right) \quad (19)$$

Applying KZG homomorphism, it follows that:

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_i(L_i(X) - 1)}{X - i} \quad (18)$$

$$q'_i(X) = q_i(X) + \delta_i \left(\frac{L_i(X) - 1}{X - i} \right) \quad (19)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_i(X) - 1}{X - i}\right)^{\delta_i} \quad (20)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_i (L_i(X) - 1)}{X - i} \quad (18)$$

$$q'_i(X) = q_i(X) + \delta_i \left(\frac{L_i(X) - 1}{X - i} \right) \quad (19)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_i(X) - 1}{X - i}\right)^{\delta_i} \quad (20)$$

Thus, each upk_i must include $c\left(\frac{L_i(X)-1}{X-i}\right)$.

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change (i, δ_i)

We know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - (v_i + \delta_i)}{X - i} \quad (16)$$

$$= \frac{(\phi(X) + \delta_i L_i(X)) - v_i - \delta_i}{X - i} \quad (17)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_i (L_i(X) - 1)}{X - i} \quad (18)$$

$$q'_i(X) = q_i(X) + \delta_i \left(\frac{L_i(X) - 1}{X - i} \right) \quad (19)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_i(X) - 1}{X - i}\right)^{\delta_i} \quad (20)$$

Thus, each upk_i must include $c\left(\frac{L_i(X)-1}{X-i}\right)$. Can derive these from g^{τ^i} 's!

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$(22)$$

$$(23)$$

$$(24)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$(23)$$

$$(24)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$(24)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$q'_i(X) = q_i(X) + \delta_j \left(\frac{L_j(X)}{X - i} \right) \quad (24)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$q'_i(X) = q_i(X) + \delta_j \left(\frac{L_j(X)}{X - i} \right) \quad (24)$$

Applying KZG homomorphism, it follows that:

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$q'_i(X) = q_i(X) + \delta_j \left(\frac{L_j(X)}{X - i} \right) \quad (24)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_j(X)}{X - i}\right)^{\delta_j} \quad (25)$$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$q'_i(X) = q_i(X) + \delta_j \left(\frac{L_j(X)}{X - i} \right) \quad (24)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_j(X)}{X - i}\right)^{\delta_j} \quad (25)$$

Problem: To update any π_i after a change to j , need $c\left(\frac{L_j(X)}{X - i}\right), \forall i \neq j$

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$q'_i(X) = q_i(X) + \delta_j \left(\frac{L_j(X)}{X - i} \right) \quad (24)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_j(X)}{X - i}\right)^{\delta_j} \quad (25)$$

Problem: To update any π_i after a change to j , need $c\left(\frac{L_j(X)}{X - i}\right), \forall i \neq j \Rightarrow O(n)$ -sized upk_j .

New Technique: Updating Proof $\pi_i = c(q_i)$ After Change $(j, \delta_j), j \neq i$

Once again, we know $\pi'_i = c(q'_i)$, where:

$$q'_i(X) = \frac{\phi'(X) - v_i}{X - i} \quad (21)$$

$$= \frac{(\phi(X) + \delta_j L_j(X)) - v_i}{X - i} \quad (22)$$

$$= \frac{\phi(X) - v_i}{X - i} - \frac{\delta_j L_j(X)}{X - i} \quad (23)$$

$$q'_i(X) = q_i(X) + \delta_j \left(\frac{L_j(X)}{X - i} \right) \quad (24)$$

Applying KZG homomorphism, it follows that:

$$\pi'_i = c(q'_i) = c(q_i) \cdot c\left(\frac{L_j(X)}{X - i}\right)^{\delta_j} \quad (25)$$

Problem: To update any π_i after a change to j , need $c\left(\frac{L_j(X)}{X - i}\right), \forall i \neq j \Rightarrow O(n)$ -sized upk_j .

Solution: Compute $c\left(\frac{L_j(X)}{X - i}\right)$ in $O(1)$ time from information in upk_i and upk_j .

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$.

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

$$\frac{A(X)}{(X-i)(X-j)} = \frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \quad (27)$$

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

$$\frac{A(X)}{(X-i)(X-j)} = \frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \quad (27)$$

Now, replacing Equation (27) into Equation (26):

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \left(\frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \right) \quad (28)$$

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

$$\frac{A(X)}{(X-i)(X-j)} = \frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \quad (27)$$

Now, replacing Equation (27) into Equation (26):

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \left(\frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \right) \quad (28)$$

As a result:

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

$$\frac{A(X)}{(X-i)(X-j)} = \frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \quad (27)$$

Now, replacing Equation (27) into Equation (26):

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \left(\frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \right) \quad (28)$$

As a result:

$$c\left(\frac{L_j(X)}{X-i}\right) = \left(c\left(\frac{A(X)}{X-i}\right)^{\frac{1}{i-j}} \cdot c\left(\frac{A(X)}{X-j}\right)^{\frac{1}{j-i}} \right)^{\frac{1}{A'(j)}} \quad (29)$$

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

$$\frac{A(X)}{(X-i)(X-j)} = \frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \quad (27)$$

Now, replacing Equation (27) into Equation (26):

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \left(\frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \right) \quad (28)$$

As a result:

$$c\left(\frac{L_j(X)}{X-i}\right) = \left(c\left(\frac{A(X)}{X-i}\right)^{\frac{1}{i-j}} \cdot c\left(\frac{A(X)}{X-j}\right)^{\frac{1}{j-i}} \right)^{\frac{1}{A'(j)}} \quad (29)$$

Thus, each *upk*_{*j*} must include $c\left(\frac{A(X)}{X-i}\right)$ and $A'(i)$.

New Technique: Compute $c\left(\frac{L_j(X)}{X-i}\right)$ in $O(1)$ Time

Let $A(X) = \prod_{i \in [0, n)} (X - i)$. Then:

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \frac{A(X)}{(X-j)(X-i)} \quad (26)$$

Next, use **partial fraction decomposition** to rewrite:

$$\frac{A(X)}{(X-i)(X-j)} = \frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \quad (27)$$

Now, replacing Equation (27) into Equation (26):

$$\frac{L_j(X)}{X-i} = \frac{1}{A'(j)} \cdot \left(\frac{1}{i-j} \cdot \frac{A(X)}{X-i} + \frac{1}{j-i} \cdot \frac{A(X)}{X-j} \right) \quad (28)$$

As a result:

$$c\left(\frac{L_j(X)}{X-i}\right) = \left(c\left(\frac{A(X)}{X-i}\right)^{\frac{1}{i-j}} \cdot c\left(\frac{A(X)}{X-j}\right)^{\frac{1}{j-i}} \right)^{\frac{1}{A'(j)}} \quad (29)$$

Thus, each **upk_i** must include $c\left(\frac{A(X)}{X-i}\right)$ and $A'(i)$. Can derive from g^{τ^i} 's!

Aggregating Proofs

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

High-level ideas, thanks to Drake and Buterin:

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

High-level ideas, thanks to Drake and Buterin:

- Each π_i is a commitment to quotient q_i of division $\frac{\phi(X)}{X-i}$

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

High-level ideas, thanks to Drake and Buterin:

- Each π_i is a commitment to quotient q_i of division $\frac{\phi(X)}{X-i}$
- π_I is a commitment to quotient q_I of division $\frac{\phi(X)}{\prod_{i \in I}(X-i)}$ (see [KZG10])

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

High-level ideas, thanks to Drake and Buterin:

- Each π_i is a commitment to quotient q_i of division $\frac{\phi(X)}{X-i}$
- π_I is a commitment to quotient q_I of division $\frac{\phi(X)}{\prod_{i \in I} (X-i)}$ (see [KZG10])
- Compute c_i 's such that $\frac{1}{\prod_{i \in I} (X-i)} = \sum_{i \in I} c_i \frac{1}{X-i}$

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

High-level ideas, thanks to Drake and Buterin:

- Each π_i is a commitment to quotient q_i of division $\frac{\phi(X)}{X-i}$
- π_I is a commitment to quotient q_I of division $\frac{\phi(X)}{\prod_{i \in I} (X-i)}$ (see [KZG10])
- Compute c_i 's such that $\frac{1}{\prod_{i \in I} (X-i)} = \sum_{i \in I} c_i \frac{1}{X-i}$
- Then, $q_I(X) = \sum_{i \in I} c_i \cdot q_i(X)$

Aggregating Proofs

Given many $(\pi_i)_{i \in I}$, can aggregate into succinct **subvector proof** π_I .

High-level ideas, thanks to Drake and Buterin:

- Each π_i is a commitment to quotient q_i of division $\frac{\phi(X)}{X-i}$
- π_I is a commitment to quotient q_I of division $\frac{\phi(X)}{\prod_{i \in I} (X-i)}$ (see [KZG10])
- Compute c_i 's such that $\frac{1}{\prod_{i \in I} (X-i)} = \sum_{i \in I} c_i \frac{1}{X-i}$
- Then, $q_I(X) = \sum_{i \in I} c_i \cdot q_i(X)$
- Thus, $\pi_I = \prod_{i \in I} \pi_i^{c_i}$

Outline

Appendix

Previous Work

Background

Kate-Zaverucha-Goldberg (KZG) Polynomial Commitments

VCS from KZG Commitments to Lagrange Polynomials

Our Techniques




Updating Proofs (Case $i = j$)

Updating Proofs (Case $i \neq j$)

Aggregating Proofs into Subvector Proofs




Extras

References i

-  Dan Boneh, Benedikt Bünz, and Ben Fisch.
Batching Techniques for Accumulators with Applications to IOPs and Stateless Blockchains.
In *CRYPTO'19*, 2019.
-  Jan Camenisch, Maria Dubovitskaya, Kristiyan Haralambiev, and Markulf Kohlweiss.
Composable and Modular Anonymous Credentials: Definitions and Practical Constructions.
In *ASIACRYPT'15*, 2015.
-  Dario Catalano and Dario Fiore.
Vector Commitments and Their Applications.
In *PKC'13*, 2013.

References ii

-  Matteo Campanelli, Dario Fiore, Nicola Greco, Dimitris Kolonelos, and Luca Nizzardo.
Vector Commitment Techniques and Applications to Verifiable Decentralized Storage, 2020.
<https://eprint.iacr.org/2020/149>.
-  Alexander Chepur, Charalampos Papamanthou, and Yupeng Zhang.
Edrax: A Cryptocurrency with Stateless Transaction Validation, 2018.
<https://eprint.iacr.org/2018/968>.
-  Dankrad Feist and Dmitry Khovratovich.
Fast amortized Kate proofs, 2020.
<https://github.com/khovratovich/Kate>.

-  Sergey Gorbunov, Leonid Reyzin, Hoeteck Wee, and Zhenfei Zhang.
Pointproofs: Aggregating Proofs for Multiple Vector Commitments, 2020.
<https://eprint.iacr.org/2020/419>.
-  Aniket Kate, Gregory M. Zaverucha, and Ian Goldberg.
Constant-Size Commitments to Polynomials and Their Applications.
In *ASIACRYPT'10*, 2010.
-  Russell W. F. Lai and Giulio Malavolta.
Subvector Commitments with Application to Succinct Arguments.
In *CRYPTO'19*, 2019.



Ralph C. Merkle.

A Digital Signature Based on a Conventional Encryption Function.

In Carl Pomerance, editor, *CRYPTO '87*, pages 369–378, Berlin, Heidelberg, 1988.
Springer Berlin Heidelberg.



Alin Tomescu, Robert Chen, Yiming Zheng, Ittai Abraham, Benny Pinkas, Guy Golan Gueta, and Srinivas Devadas.

Towards Scalable Threshold Cryptosystems.

In *IEEE S&P'20*, May 2020.



Alin Tomescu.

How to Keep a Secret and Share a Public Key (Using Polynomial Commitments).

PhD thesis, Massachusetts Institute of Technology, Cambridge, MA, USA, 2020.