






AN INTRODUCTION TO
ALGEBRAIC NUMBER THEORY

Alistair Pattison




November 2, 2023










I wonder if there
are more solutions?



I wonder if there
are more solutions?

When is recess?



GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$z^2 = x^2 + y^2$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \end{aligned} \quad \text{over } \mathbb{Z}[i]$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \quad \text{over } \mathbb{Z}[i] \end{aligned}$$

x and y relatively prime $\implies \alpha$ and β relatively prime

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \quad \text{over } \mathbb{Z}[i] \end{aligned}$$

x and y relatively prime $\implies \alpha$ and β relatively prime

$$z^2 = \alpha\beta \implies \alpha = u\gamma^2$$

$$\gamma \in \mathbb{Z}[i], u \in \{\pm 1, \pm i\}$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \end{aligned} \quad \text{over } \mathbb{Z}[i]$$

$$\alpha = \gamma^2$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \end{aligned} \quad \text{over } \mathbb{Z}[i]$$

$$\begin{aligned} \alpha &= \gamma^2 \\ &= (a + bi)^2 \end{aligned}$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \end{aligned} \quad \text{over } \mathbb{Z}[i]$$

$$\begin{aligned} \alpha &= \gamma^2 \\ &= (a + bi)^2 \\ &= \underbrace{(a^2 - b^2)}_x + \underbrace{2ab}_y i \end{aligned}$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES

Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$\begin{aligned} z^2 &= x^2 + y^2 \\ &= \underbrace{(x + iy)}_{\alpha} \underbrace{(x - iy)}_{\beta} \end{aligned} \quad \text{over } \mathbb{Z}[i]$$

$$\begin{aligned} \alpha &= \gamma^2 & x &= a^2 - b^2 \\ &= (a + bi)^2 & y &= 2ab \\ &= \underbrace{(a^2 - b^2)}_x + \underbrace{2ab}_y i & z &= a^2 + b^2 \end{aligned}$$

GENERATING PRIMITIVE PYTHAGOREAN TRIPLES


Goal: find all relatively prime $x, y, z \in \mathbb{Z}$ such that

$$z^2 = x^2 + y^2$$

a	b	
2	1	$3^2 + 4^2 = 5^2$
3	2	$5^2 + 12^2 = 13^2$
4	3	$7^2 + 24^2 = 25^2$
4	2	$12^2 + 16^2 = 20^2$
4	1	$15^2 + 8^2 = 17^2$

ALGEBRAIC NUMBER THEORY

ALGEBRAIC NUMBER THEORY



Using tools from algebra like
rings and field extensions

ALGEBRAIC NUMBER THEORY

```
graph TD; A[ALGEBRAIC] --- B[NUMBER THEORY]; B --- C[Generating insight about the integers and the primes];
```

The diagram consists of two main parts. At the top, the words 'ALGEBRAIC' and 'NUMBER THEORY' are joined by a horizontal line. 'ALGEBRAIC' is in a black serif font, while 'NUMBER THEORY' is in a green serif font and is enclosed in a light green rectangular box. From the bottom of 'ALGEBRAIC', a thin black line extends diagonally down and to the left. From the bottom of 'NUMBER THEORY', a thin black line extends diagonally down and to the right. These lines point towards two separate text blocks at the bottom of the slide.


Using tools from algebra like
rings and field extensions

Generating insight about
the integers and the primes

ALGEBRAIC NUMBER THEORY

Using tools from algebra like
rings and field extensions

Generating insight about
the integers and the primes



OUTLINE

1. MATH 342 IN 3:42

OUTLINE

1. MATH 342 IN 3:42
2. NUMBER FIELDS AND NUMBER RINGS


OUTLINE

1. MATH 342 IN 3:42
2. NUMBER FIELDS AND NUMBER RINGS
3. THE IDEAL CLASS GROUP

MATH 342 IN 3:42



PLAYING FROM ALBUM
Math (Alistair's Version)




Algebra (5 Minute Version) (Al 
Alistair Pattison

0:00 5:00




RING HIERARCHY



DEFINITION (COMMUTATIVE RING)

RING HIERARCHY



DEFINITION (COMMUTATIVE RING)

"Things like the integers"

RING HIERARCHY




DEFINITION (COMMUTATIVE RING)

"Things like the integers"

- Addition/subtraction: $a + b, a - b \in R$

RING HIERARCHY




DEFINITION (COMMUTATIVE RING)

"Things like the integers"

- Addition/subtraction: $a + b, a - b \in R$
- Multiplication: $ab = ba \in R$

RING HIERARCHY




DEFINITION (COMMUTATIVE RING)

"Things like the integers"

- Addition/subtraction: $a + b, a - b \in R$
- Multiplication: $ab = ba \in R$
- Distributive property: $a(b + c) = ab + bc$

RING HIERARCHY




DEFINITION (COMMUTATIVE RING)

"Things like the integers"

- Addition/subtraction: $a + b, a - b \in R$
- Multiplication: $ab = ba \in R$
- Distributive property: $a(b + c) = ab + bc$
- **Not** division: $a/b \notin R$


RING HIERARCHY



DEFINITION (IDEAL)

An (additive) subgroup, I , such that $ra \in I$ for all $r \in R$, $a \in I$.

RING HIERARCHY




DEFINITION (IDEAL)

An (additive) subgroup, I , such that $ra \in I$ for all $r \in R$, $a \in I$.

- **Example:** Even integers ($8 \cdot 7$ is even)

RING HIERARCHY




DEFINITION (IDEAL)

An (additive) subgroup, I , such that $ra \in I$ for all $r \in R$, $a \in I$.

- **Example:** Even integers ($8 \cdot 7$ is even)
- **Non-example:** Odd integers ($7 \cdot 8$ is not odd)

RING HIERARCHY



DEFINITION (IDEAL)

An (additive) subgroup, I , such that $ra \in I$ for all $r \in R$, $a \in I$.

- **Example:** Even integers ($8 \cdot 7$ is even)
- **Non-example:** Odd integers ($7 \cdot 8$ is not odd)

DEFINITION (PRIME IDEAL)

An ideal, P , such that $ab \in P$ implies $a \in P$ or $b \in P$.

RING HIERARCHY



DEFINITION (IDEAL)

An (additive) subgroup, I , such that $ra \in I$ for all $r \in R$, $a \in I$.


- **Example:** Even integers ($8 \cdot 7$ is even)
- **Non-example:** Odd integers ($7 \cdot 8$ is not odd)

DEFINITION (PRIME IDEAL)


An ideal, P , such that $ab \in P$ implies $a \in P$ or $b \in P$.

- This is a generalization of Euclid's Lemma

RING HIERARCHY




RING HIERARCHY



DEFINITION

A commutative ring is an *integral domain* if $ab = 0$ implies $a = 0$ or $b = 0$. (No zero divisors.)

RING HIERARCHY




DEFINITION

A commutative ring is an *integral domain* if $ab = 0$ implies $a = 0$ or $b = 0$. (No zero divisors.)

- **Example:** \mathbb{Z} , $\mathbb{Z}[i]$

RING HIERARCHY



DEFINITION


A commutative ring is an *integral domain* if $ab = 0$ implies $a = 0$ or $b = 0$. (No zero divisors.)

- **Example:** \mathbb{Z} , $\mathbb{Z}[i]$
- **Non-example:** in $\mathbb{Z}/8\mathbb{Z}$, we have $4 \cdot 2 = 0$

RING HIERARCHY




RING HIERARCHY



DEFINITION (INTEGRALLY CLOSED DOMAIN)

A ring R is *integrally closed* if for all $\alpha/\beta \in \text{Frac } R$ that are integral over R , then $\beta \mid \alpha$, i.e., $\alpha/\beta \in R$.

RING HIERARCHY




DEFINITION (INTEGRALLY CLOSED DOMAIN)

A ring R is *integrally closed* if for all $\alpha/\beta \in \text{Frac } R$ that are integral over R , then $\beta \mid \alpha$, i.e., $\alpha/\beta \in R$.

- $\text{Frac } \mathbb{Z} = \mathbb{Q}$; $\text{Frac } \mathbb{C}[x]$ is the field of rational functions, $\mathbb{C}(x)$

RING HIERARCHY




DEFINITION (INTEGRALLY CLOSED DOMAIN)

A ring R is *integrally closed* if for all $\alpha/\beta \in \text{Frac } R$ that are integral over R , then $\beta \mid \alpha$, i.e., $\alpha/\beta \in R$.

- $\text{Frac } \mathbb{Z} = \mathbb{Q}$; $\text{Frac } \mathbb{C}[x]$ is the field of rational functions, $\mathbb{C}(x)$
- If $f(p/q) = 0$ with monic $f \in \mathbb{Z}[x]$, then $q \mid p$

RING HIERARCHY




DEFINITION (INTEGRALLY CLOSED DOMAIN)


A ring R is *integrally closed* if for all $\alpha/\beta \in \text{Frac } R$ that are integral over R , then $\beta \mid \alpha$, i.e., $\alpha/\beta \in R$.

- $\text{Frac } \mathbb{Z} = \mathbb{Q}$; $\text{Frac } \mathbb{C}[x]$ is the field of rational functions, $\mathbb{C}(x)$
- If $f(p/q) = 0$ with monic $f \in \mathbb{Z}[x]$, then $q \mid p$
- **Example:** \mathbb{Z}

RING HIERARCHY



RING HIERARCHY




DEFINITION (UFD)

A commutative ring R is a *unique factorization domain* if every element factors uniquely into irreducible elements.


- **Example:** \mathbb{Z}
- **Non-example:** $\mathbb{Z}[i\sqrt{5}]$:

$$6 = 2 \cdot 3 = (1 + i\sqrt{5})(1 - i\sqrt{5})$$

RING HIERARCHY




RING HIERARCHY



DEFINITION (PID)

A commutative ring R is a *principal ideal domain* if every ideal is generated by a single element.

RING HIERARCHY




DEFINITION (PID)

A commutative ring R is a *principal ideal domain* if every ideal is generated by a single element.

- **Example:** \mathbb{Z} , $\mathbb{Q}[x]$, $\mathbb{Z}[i]$

RING HIERARCHY




DEFINITION (PID)


A commutative ring R is a *principal ideal domain* if every ideal is generated by a single element.

- **Example:** \mathbb{Z} , $\mathbb{Q}[x]$, $\mathbb{Z}[i]$
- **Non-example:** $\mathbb{Z}[x]$ because of $(2, x)$

RING HIERARCHY




RING HIERARCHY



DEFINITION (FIELD)

"Things like the rationals" or "rings where you can divide".

RING HIERARCHY




DEFINITION (FIELD)

"Things like the rationals" or "rings where you can divide".

- **Example:** \mathbb{Q} , \mathbb{C} , \mathbb{R}

RING HIERARCHY



DEFINITION (FIELD)

"Things like the rationals" or "rings where you can divide".

- **Example:** \mathbb{Q} , \mathbb{C} , \mathbb{R}
- **Non-example:** \mathbb{Z} , $\mathbb{Z}[x]$

FIELD EXTENSIONS

$$f(x) = x^2 + 5$$

FIELD EXTENSIONS

$$f(x) = x^2 + 5$$

- Over \mathbb{Q} , f is irreducible

FIELD EXTENSIONS

$$\begin{aligned}f(x) &= x^2 + 5 \\&= (x - i\sqrt{5})(x + i\sqrt{5})\end{aligned}$$

- Over \mathbb{Q} , f is irreducible
- Over \mathbb{C} , f factors

$$e \quad \pi \quad \sqrt{7} \quad 1 + 2\sqrt{17}$$

FIELD EXTENSIONS

$$\begin{aligned}f(x) &= x^2 + 5 \\ &= (x - i\sqrt{5})(x + i\sqrt{5})\end{aligned}$$

- Over \mathbb{Q} , f is irreducible
- Over \mathbb{C} , f factors

$$e \quad \pi \quad \sqrt{7} \quad 1 + 2\sqrt{17}$$

- But f is equally happy living in

$$\mathbb{Q}(i\sqrt{5}) = \{a + bi\sqrt{5} : a, b \in \mathbb{Q}\}$$

NUMBER FIELDS AND NUMBER RINGS

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

THEOREM

Any number field can be written in the form

$$K = \mathbb{Q}[\alpha] = \text{span}_{\mathbb{Q}}\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$$

where n is the degree of the minimal polynomial of α .

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

THEOREM

Any number field can be written in the form

$$K = \mathbb{Q}[\alpha] = \text{span}_{\mathbb{Q}}\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$$

where n is the degree of the minimal polynomial of α .

Examples

- $\mathbb{Q}[i\sqrt{5}] = \{a + ib\sqrt{5} : a, b \in \mathbb{Q}\}$

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

THEOREM

Any number field can be written in the form

$$K = \mathbb{Q}[\alpha] = \text{span}_{\mathbb{Q}}\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$$

where n is the degree of the minimal polynomial of α .

Examples

- $\mathbb{Q}[i\sqrt{5}] = \{a + ib\sqrt{5} : a, b \in \mathbb{Q}\}$
- $\mathbb{Q}[\omega], \omega = e^{2\pi i/p}$ (cyclotomic fields)

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

THEOREM

Any number field can be written in the form

$$K = \mathbb{Q}[\alpha] = \text{span}_{\mathbb{Q}}\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$$

where n is the degree of the minimal polynomial of α .

Examples

- $\mathbb{Q}[i\sqrt{5}] = \{a + ib\sqrt{5} : a, b \in \mathbb{Q}\}$
- $\mathbb{Q}[\omega]$, $\omega = e^{2\pi i/p}$ (cyclotomic fields)
- $\mathbb{Q}[\sqrt{m}]$, m squarefree (quadratic fields)

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

THEOREM

Any number field can be written in the form

$$K = \mathbb{Q}[\alpha] = \text{span}_{\mathbb{Q}}\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$$

where n is the degree of the minimal polynomial of α .

Examples

- $\mathbb{Q}[i\sqrt{5}] = \{a + ib\sqrt{5} : a, b \in \mathbb{Q}\}$
- $\mathbb{Q}[\omega]$, $\omega = e^{2\pi i/p}$ (cyclotomic fields)
- $\mathbb{Q}[\sqrt{m}]$, m squarefree (quadratic fields)

NUMBER FIELDS

DEFINITION (NUMBER FIELD)

A *number field* $K \subset \mathbb{C}$ is a finite extension of \mathbb{Q} .

THEOREM

Any number field can be written in the form

$$K = \mathbb{Q}[\alpha] = \text{span}_{\mathbb{Q}}\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$$

where n is the degree of the minimal polynomial of α .

Examples

- $\mathbb{Q}[i\sqrt{5}] = \{a + ib\sqrt{5} : a, b \in \mathbb{Q}\}$
- $\mathbb{Q}[\omega]$, $\omega = e^{2\pi i/p}$ (cyclotomic fields)
- $\mathbb{Q}[\sqrt{m}]$, m squarefree (quadratic fields)

Non-examples

- $\mathbb{Q}[\pi]$ because π is transcendental

NUMBER RINGS

NUMBER RINGS

DEFINITION (ALGEBRAIC INTEGER)

An *algebraic integer* is a complex number α that is the root of some monic polynomial $f \in \mathbb{Z}[x]$.

NUMBER RINGS

DEFINITION (ALGEBRAIC INTEGER)

An *algebraic integer* is a complex number α that is the root of some monic polynomial $f \in \mathbb{Z}[x]$.

We use \mathbb{A} to denote the set of all algebraic integers.

- Any integer is an algebraic integer

NUMBER RINGS

DEFINITION (ALGEBRAIC INTEGER)

An *algebraic integer* is a complex number α that is the root of some monic polynomial $f \in \mathbb{Z}[x]$.

We use \mathbb{A} to denote the set of all algebraic integers.

- Any integer is an algebraic integer
- $i\sqrt{5} \in \mathbb{A}$ because of $f(x) = x^2$

NUMBER RINGS

DEFINITION (ALGEBRAIC INTEGER)

An *algebraic integer* is a complex number α that is the root of some monic polynomial $f \in \mathbb{Z}[x]$.

We use \mathbb{A} to denote the set of all algebraic integers.

- Any integer is an algebraic integer
- $i\sqrt{5} \in \mathbb{A}$ because of $f(x) = x^2$
- $2 + \sqrt[3]{17} \in \mathbb{A}$ because of $f(x) = x^3 - 6x^2 + 12x - 25$

NUMBER RINGS

DEFINITION (ALGEBRAIC INTEGER)

An *algebraic integer* is a complex number α that is the root of some monic polynomial $f \in \mathbb{Z}[x]$.

We use \mathbb{A} to denote the set of all algebraic integers.

- Any integer is an algebraic integer
- $i\sqrt{5} \in \mathbb{A}$ because of $f(x) = x^2$
- $2 + \sqrt[3]{17} \in \mathbb{A}$ because of $f(x) = x^3 - 6x^2 + 12x - 25$
- \mathbb{A} is a subring of \mathbb{C}

NUMBER RINGS

DEFINITION (NUMBER RING)

The *number ring* of a number field $K = \mathbb{Q}(\alpha)$ is the set

$$\mathcal{O}_K = \mathbb{A} \cap K.$$

NUMBER RINGS

DEFINITION (NUMBER RING)

The *number ring* of a number field $K = \mathbb{Q}(\alpha)$ is the set

$$\mathcal{O}_K = \mathbb{A} \cap K.$$

- $\mathcal{O}_{\mathbb{Q}} = \mathbb{A} \cap \mathbb{Q} = \mathbb{Z}$

NUMBER RINGS

DEFINITION (NUMBER RING)

The *number ring* of a number field $K = \mathbb{Q}(\alpha)$ is the set

$$\mathcal{O}_K = \mathbb{A} \cap K.$$

- $\mathcal{O}_{\mathbb{Q}} = \mathbb{A} \cap \mathbb{Q} = \mathbb{Z}$
- $\mathcal{O}_{\mathbb{Q}[i]} = \mathbb{Z}[i]$

NUMBER RINGS

DEFINITION (NUMBER RING)

The *number ring* of a number field $K = \mathbb{Q}(\alpha)$ is the set

$$\mathcal{O}_K = \mathbb{A} \cap K.$$

- $\mathcal{O}_{\mathbb{Q}} = \mathbb{A} \cap \mathbb{Q} = \mathbb{Z}$
- $\mathcal{O}_{\mathbb{Q}[i]} = \mathbb{Z}[i]$
- $\mathcal{O}_{\mathbb{Q}[\zeta]} = \mathbb{Z}[\zeta]$ for primitive roots ζ

NUMBER RINGS

DEFINITION (NUMBER RING)

The *number ring* of a number field $K = \mathbb{Q}(\alpha)$ is the set

$$\mathcal{O}_K = \mathbb{A} \cap K.$$

- $\mathcal{O}_{\mathbb{Q}} = \mathbb{A} \cap \mathbb{Q} = \mathbb{Z}$
- $\mathcal{O}_{\mathbb{Q}[i]} = \mathbb{Z}[i]$
- $\mathcal{O}_{\mathbb{Q}[\zeta]} = \mathbb{Z}[\zeta]$ for primitive roots ζ
- $\mathcal{O}_{\mathbb{Q}[\sqrt{5}]} = \mathbb{Z}[1, \frac{1}{2} + \frac{1}{2}\sqrt{5}]$

DEDEKIND DOMAINS

THEOREM

Number rings are Dedekind domains.

DEFINITION (DEDEKIND DOMAIN)

A *Dedekind domain* is an integrally closed domain R such that

1. every ideal is finitely generated and
2. every nonzero prime ideal is maximal.

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.
- In $R = \mathbb{Z}[i\sqrt{5}]$,

$$\begin{aligned} 6 &= 2 \cdot 3 \\ &= (1 + i\sqrt{5})(1 - i\sqrt{5}) \end{aligned}$$

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.
- In $R = \mathbb{Z}[i\sqrt{5}]$,

$$\begin{aligned}6 &= 2 \cdot 3 \\ &= (1 + i\sqrt{5})(1 - i\sqrt{5})\end{aligned}$$

$$(6) = (2, 1 + i\sqrt{5})^2 (3, 1 + i\sqrt{5}) (3, 1 - i\sqrt{5})$$

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.
- In $R = \mathbb{Z}[i\sqrt{5}]$,

$$\begin{aligned}6 &= 2 \cdot 3 \\ &= (1 + i\sqrt{5})(1 - i\sqrt{5})\end{aligned}$$

$$(6) = (2, 1 + i\sqrt{5})^2 (3, 1 + i\sqrt{5}) (3, 1 - i\sqrt{5})$$

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.
- In $R = \mathbb{Z}[i\sqrt{5}]$,

$$\begin{aligned}6 &= 2 \cdot 3 \\ &= (1 + i\sqrt{5})(1 - i\sqrt{5})\end{aligned}$$

$$(6) = (2, 1 + i\sqrt{5})^2 (3, 1 + i\sqrt{5}) (3, 1 - i\sqrt{5})$$

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.
- In $R = \mathbb{Z}[i\sqrt{5}]$,

$$\begin{aligned} 6 &= 2 \cdot 3 \\ &= (1 + i\sqrt{5})(1 - i\sqrt{5}) \end{aligned}$$

$$(6) = (2, 1 + i\sqrt{5})^2 (3, 1 + i\sqrt{5}) (3, 1 - i\sqrt{5})$$

DEDEKIND DOMAINS

FACTORING IDEALS

THEOREM

Every ideal of a Dedekind domain R uniquely factors into prime ideals.

- Allows replacing unique factorization of elements with unique factorization of ideals.
- In $R = \mathbb{Z}[i\sqrt{5}]$,

$$\begin{aligned} 6 &= 2 \cdot 3 \\ &= (1 + i\sqrt{5})(1 - i\sqrt{5}) \end{aligned}$$

$$(6) = (2, 1 + i\sqrt{5})^2 (3, 1 + i\sqrt{5}) (3, 1 - i\sqrt{5})$$

DEDEKIND DOMAINS

IN THE CLASS HIERARCHY

THEOREM


A UFD is a PID iff it's a Dedekind domain.

DEDEKIND DOMAINS

IN THE CLASS HIERARCHY

THEOREM

A UFD is a PID iff it's a Dedekind domain.




DEDEKIND DOMAINS

IN THE CLASS HIERARCHY

THEOREM

A UFD is a PID iff it's a Dedekind domain.



- **DD, not UFD**
 $\mathbb{Z}[i\sqrt{5}]$


DEDEKIND DOMAINS

IN THE CLASS HIERARCHY

THEOREM

A UFD is a PID iff it's a Dedekind domain.

Integrally Closed Domains



- **DD, not UFD**

$$\mathbb{Z}[i\sqrt{5}]$$

- **UFD, not DD**

$$\mathbb{R}[x_1, x_2, \dots], \mathbb{Q}[x, y]$$

THE IDEAL CLASS GROUP

THE IDEAL CLASS GROUP

DEFINITION (IDEAL CLASS GROUP)

Let $K = \mathbb{Q}[\alpha]$ be a number field. The *class group* of K is the set of ideals of \mathcal{O}_K , modulo the equivalence relation

$$I \sim J \text{ iff } \alpha I = \beta J \text{ for some nonzero } \alpha, \beta \in R.$$

THE IDEAL CLASS GROUP

DEFINITION (IDEAL CLASS GROUP)

Let $K = \mathbb{Q}[\alpha]$ be a number field. The *class group* of K is the set of ideals of \mathcal{O}_K , modulo the equivalence relation

$$I \sim J \text{ iff } \alpha I = \beta J \text{ for some nonzero } \alpha, \beta \in R.$$

- Back to $K = \mathbb{Q}[i\sqrt{5}]$ and $\mathcal{O}_K = \mathbb{Z}[i\sqrt{5}]$

$$(2) \sim (3)$$

$$2(3) = (6) = 3(2)$$

THE IDEAL CLASS GROUP

DEFINITION (IDEAL CLASS GROUP)

Let $K = \mathbb{Q}[\alpha]$ be a number field. The *class group* of K is the set of ideals of \mathcal{O}_K , modulo the equivalence relation

$$I \sim J \text{ iff } \alpha I = \beta J \text{ for some nonzero } \alpha, \beta \in R.$$

- Back to $K = \mathbb{Q}[i\sqrt{5}]$ and $\mathcal{O}_K = \mathbb{Z}[i\sqrt{5}]$

$$(2) \sim (3)$$

$$2(3) = (6) = 3(2)$$

$$(2) \not\sim (2, 1 + i\sqrt{5})$$

THE IDEAL CLASS GROUP

DEFINITION (IDEAL CLASS GROUP)

Let $K = \mathbb{Q}[\alpha]$ be a number field. The *class group* of K is the set of ideals of \mathcal{O}_K , modulo the equivalence relation

$$I \sim J \text{ iff } \alpha I = \beta J \text{ for some nonzero } \alpha, \beta \in R.$$

- Back to $K = \mathbb{Q}[i\sqrt{5}]$ and $\mathcal{O}_K = \mathbb{Z}[i\sqrt{5}]$

$$(2) \sim (3)$$

$$2(3) = (6) = 3(2)$$

$$(2) \not\sim (2, 1 + i\sqrt{5})$$

- That's it.

THE IDEAL CLASS GROUP

DEFINITION (IDEAL CLASS GROUP)

Let $K = \mathbb{Q}[\alpha]$ be a number field. The *class group* of K is the set of ideals of \mathcal{O}_K , modulo the equivalence relation

$$I \sim J \text{ iff } \alpha I = \beta J \text{ for some nonzero } \alpha, \beta \in R.$$

- Back to $K = \mathbb{Q}[i\sqrt{5}]$ and $\mathcal{O}_K = \mathbb{Z}[i\sqrt{5}]$

$$(2) \sim (3)$$

$$2(3) = (6) = 3(2)$$

$$(2) \not\sim (2, 1 + i\sqrt{5})$$

- That's it.
- The class group of $\mathbb{Q}[i\sqrt{5}]$ is \mathbb{Z}_2 .

THE CLASS NUMBER

DEFINITION (CLASS NUMBER)

The class number of a number field $K = \mathbb{Q}[\alpha]$ is the size of its ideal class group.

THE CLASS NUMBER

DEFINITION (CLASS NUMBER)

The class number of a number field $K = \mathbb{Q}[\alpha]$ is the size of its ideal class group.

- $\mathbb{Q}[i\sqrt{5}]$ has class number 2
- K has class number 1 iff \mathcal{O}_K is a UFD
- Measures "how far away" \mathcal{O}_K is from achieving unique factorization

THE CLASS NUMBER

DEFINITION (CLASS NUMBER)

The class number of a number field $K = \mathbb{Q}[\alpha]$ is the size of its ideal class group.

- $\mathbb{Q}[i\sqrt{5}]$ has class number 2
- K has class number 1 iff \mathcal{O}_K is a UFD
- Measures "how far away" \mathcal{O}_K is from achieving unique factorization

THEOREM

Class numbers are always finite.

THANK YOU!

THANK YOU!



Slides