Supplementary online content

Table S1. Assessment of quality of evidence for case-report and case series
Source: Murad, M. H., S. Sultan, S. Haffar and F. Bazerbachi (2018). Methodological quality and synthesis of case series and case reports. BMJ Evid Based Med 2018;23(2): 60-63.

Items 4-6 are relevant to cases of adverse drug events according to Murad et al above.

Study (Country)	Number of cases (age)	1. Does the patient(s) represent(s) the whole experience of the investigator (centre) or is the selection method unclear to the extent that other patients with similar presentation may not have been reported?	2. Was the exposure adequately ascertained?	3. Was the outcome adequately ascertained?	4. Were other alternative causes that may explain the observation ruled out?	5. Was there a challenge/rechallenge phenomenon?	6. Was there a dose–response effect?	7. Was follow-up long enough for outcomes to occur?	8. Is the case(s) described with sufficient details to allow other investigators to replicate the research or to allow practitioners make inferences related to their own practice?
Casalino (Italy)[1]	1 (48 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Cheema (Canada)[2]	1 (29 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Chen (China)[3]	1 (30 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Colavita (Italy)[4]	1 (65 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Daruich (France)[5]	1 (27 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Unsure
Khavand (Iran)[6]	1 (65 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes

Mendez Mangana (Spain)[7]	1 (31 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Navel (France)[8]	1 (63 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Pascual-Prieto (Spain)[9]	7 (age unstated)	Whole experience	Mostly yes	Yes	NA	NA	NA	Yes	Yes
Salducci (Italy)[10]	1 (72 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Scalinci (Italy)[11]	5 (37, 41, 43, 48, 65 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Unsure	Unsure
Stevens (USA)[12]	1 (74 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Unsure	Unsure
Wu (China)[13]	1 (2 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Ying (Malaysia)[14]	1 (54 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Hu (China)[15]	1 (70 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes
Huang (China)[16]	1 (22 yrs old)	Whole experience	Yes	Yes	NA	NA	NA	Yes	Yes

yrs – years

Note: Four case reports/series were not included:

- one detailed the COVID-19 outbreak in an ophthalmology department in Norway;
- one reported eye injury caused by improper use of UV light for disinfection at home during COVID-19 pandemic;
- one described mucous membrane pemphigoid and COVID-19 treated with high-dose intravenous immunoglobulins
- one described eye jury during lockdown.

Table S2. Assessment of quality of evidence for cross-sectional studies
Source: The Agency for Healthcare Research and Quality. Methodology Checklist for Cross-sectional study. Available at https://www.ncbi.nlm.nih.gov/books/NBK35156/ Accessed Jul 15, 2020.

Each item was assessed with Yes, No, Unclear and not applicable (NA).

Study (country, # Yes/11)	1. Define the source of information (survey, record, review)	2. List inclusion and exclusion criteria for exposed and unexposed subjects (cases and controls) or refer to previous publications	3. Indicate time period used for identifying patients	4. Indicate whether or not subjects were consecutive if not population-based	5. Indicate if evaluators of subjective components of study were masked to other aspects of the status of the participants	6. Describe any assessments undertaken for quality assurance purposes (e.g., test/retest of primary outcome measurements)	7. Explain any patient exclusions from analysis	8. Describe how confounding was assessed and/or controlled.	9. If applicable, explain how missing data were handled in the analysis	10. Summarize patient response rates and completeness of data collection	11. Clarify what follow-up, if any, was expected and the percentage of patients for which incomplete data or follow-up was obtained
Conjunctivitis or other ocular findings in COVID-19 patients											
Chen, (China, 10/11)[17]	Yes	Yes (hospitalized/ severe COVID-19 pts were excluded)	Yes (Feb 1 - Mar 1)	Yes	Unclear	Yes (2 researchers reviewed the data collection forms)	Yes	Yes	Yes	Yes	Yes
Güemes-Villahoz (Spain, 6/11)[18]	Yes	Yes (hospitalized pts with COVID-19 were included. ICU pts were excluded)	Yes (Apr 14-May 15)	Yes	No	Yes (2 ophthalmologists examined pts for diagnosis)	Yes	No	No	Unclear	NA
Bostanci Ceran (Turkey, 5/11)[19]	Yes	Yes (hospitalized pts with COVID-19 diagnosis	Yes (Mar 11-Apr 30)	Yes	No	No	No	No	No	Yes	Unclear

		were included)									
Hong (China, 5/11)[20]	Yes	Yes (hospitalized and discharged pts)	Yes (Jan 29-Feb 29)	Yes	No	No	No	No	NA	Yes	NA
Karimi (Iran, 5/11)[21]	Yes	Yes (hospitalized pts were included)	Yes (Mar 28-Apr 5)	Probably yes	No	No	No	No	Unclear	Yes	NA
Xia (China, 5/11)[22]	Yes	Yes	Yes (Jan 26-Feb 9)	Probably yes	No	Yes (2 samples of each pt, both tear sample and sputum sample from each pt)	No	No	NA	Unclear	NA
Xie (China, 5/11)[23]	Yes	Yes	Yes (Feb 12-Feb 28)	Yes	No	No	Yes	No	NA	unclear	NA
Kumar (India, 4/11)[24]	Yes	Yes (Critically ill pts were excluded)	Yes (Apr 2020)	Unclear	No	No	Yes	No	NA	No	NA
Lan (China, 4/11)[25]	Yes	Yes	Yes (Feb 6-Mar 6)	Yes	No	Unclear	No	No	Unclear	Unclear	NA
Wu (China, 4/11)[26]	Yes	Yes (hospitalized pts were included)	Yes (Feb 9-15)	Yes	Unclear	Unclear	No	No	Unclear	Unclear	Unclear
Zhang, China (3/11)[27]	Yes	Unclear	Yes (Dec 30-Feb 7)	Yes	No	No	No	No	Unclear	Unclear	NA
Zhou (China, 3/11)[28]	Yes	Yes	Yes (Jan 17-Feb 16)	Probably yes	No	No	No	No	Unclear	Unclear	NA

Deng (China, 1/11)[29]	Unclear	Yes (hospitalized pts)	No	Unclear	Unclear	No	No	No	NA	No	NA
Seah Yu (Singapore, 1/11)[30]	Unclear	Unclear	Unclear	Unclear	Unclear	Yes (tear samples and nasopharyngeal swabs)	Unclear	No	Unclear	Unclear	NA
Ocular injury or ED visits during lockdown											
Hamroush (UK, 4/11)[31]	Yes	Yes	Yes (1 wk up to Apr 15)	Yes	No	No	NA	No	NA	NA	NA
Pellegrini (Italy, 4/11, ocular trauma)[32]	Yes	Yes	Yes (Mar 10-Apr 10)	Yes	No	No	No	No	NA	Unclear	NA
Pellegrini (Italy, 4/11, ED visits)[33]	Yes	Yes	Yes (Mar 10-Apr 20)	Yes	No	Unclear	No	No	NA	unclear	NA

ED – emergency department; pt(s) – patient(s)

Note: Twelve studies were not included in this assessment because the focus of the studies was on:

- increased nonattendance of eye appointments during COVID-19 outbreak;
- perception of occupational risk in vitreoretinal surgery fellows;
- managing ocular allergy during COVID-19;
- survey ophthalmologists on COVID-19 guidelines;
- the impact of COVID-19 lockdown on ophthalmology training;
- contact lens wearing during COVID-19 lockdown;
- effect of COVID-19 lockdown on ophthalmologic practice;
- future use of personal protective equipment;
- symptomatic COVID-19 in ophthalmologists/nurses vs ophthalmologists/nurses without COVID-19;
- eye care practice patterns during COVID-19 pandemic;
- experience of eye patients to video-based telemedicine consultations during COVID-19;
- psychological impact of COVID-19 on ophthalmologists and ophthalmologists-in-training.

Table S3. Assessment of quality of evidence for systematic review and meta-analyses.

Source: Shea, B. J., J. M. Grimshaw, G. A. Wells, M. Boers, N. Andersson, C. Hamel, A. C. Porter, P. Tugwell, D. Moher and L. M. Bouter (2007). "Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews." BMC Med Res Methodol 2007;7: 10.

Each item was assessed with Yes, No, Can't answer and Not applicable.

Study (Country)	1. Was an 'a priori' design provided?	2. Was there duplicate study selection and data extraction?	3. Was a comprehensive literature search performed?	4. Was the status of publication (i.e. grey literature) used as an inclusion criterion?	5. Was a list of studies (included and excluded) provided?	6. Were the characteristics of the included studies provided?	7. Was the scientific quality of the included studies assessed and documented?	8. Was the scientific quality of the included studies used appropriately in formulating conclusions?	9. Were the methods used to combine the findings of studies appropriate?	10. Was the likelihood of publication bias assessed?	11. Was the conflict of interest stated?
Loffredo et al (Italy)[34]	Yes	Yes	Yes	Can't answer	Yes	Yes	No	No	Yes	Yes	No
Ulhaq et al (Indonesia)[35]	Yes	Can't answer	Yes	No	No	Yes	No	No	Yes	Yes	Yes
Liu et al (China)[36]	Yes	Can't answer	Yes	No	Can't answer	No	No	No	Yes	No	No

References

1. Casalino G, Monaco G, Di Sarro PP, David A, Scialdone A. Coronavirus disease 2019 presenting with conjunctivitis as the first symptom. *Eye (Lond).* 2020.
2. Cheema M, Aghazadeh H, Nazarali S, et al. Keratoconjunctivitis as the initial medical presentation of the novel coronavirus disease 2019 (COVID-19). *Can J Ophthalmol.* 2020.
3. Chen L, Liu M, Zhang Z, et al. Ocular manifestations of a hospitalised patient with confirmed 2019 novel coronavirus disease. *Br J Ophthalmol.* 2020.
4. Colavita F, Lapa D, Carletti F, et al. SARS-CoV-2 Isolation From Ocular Secretions of a Patient With COVID-19 in Italy With Prolonged Viral RNA Detection. *Ann Intern Med.* 2020.
5. Daruich A, Martin D, Bremond-Gignac D. Ocular manifestation as first sign of Coronavirus Disease 2019 (COVID-19): Interest of telemedicine during the pandemic context. *J Fr Ophtalmol.* 2020.
6. Khavandi S, Tabibzadeh E, Naderan M, Shoar S. Corona virus disease-19 (COVID-19) presenting as conjunctivitis: atypically high-risk during a pandemic. *Cont Lens Anterior Eye.* 2020.
7. Mendez Mangana C, Barraquer Kargacin A, Barraquer RI. Episcleritis as an ocular manifestation in a patient with COVID-19. *Acta Ophthalmol.* 2020.
8. Navel V, Chiambaretta F, Dutheil F. Haemorrhagic conjunctivitis with pseudomembranous related to SARS-CoV-2. *Am J Ophthalmol Case Rep.* 2020.
9. Pascual-Prieto J, Narvaez-Palazon C, Porta-Etessam J, Gomez-de-Liano R. COVID-19 epidemic: Should ophthalmologists be aware of oculomotor paresis? *Arch Soc Esp Oftalmol.* 2020.
10. Salducci M, La Torre G. COVID-19 emergency in the cruise's ship: a case report of conjunctivitis. *Clin Ter.* 2020.
11. Scalinci SZ, Trovato Battagliola E. Conjunctivitis can be the only presenting sign and symptom of COVID-19. *IDCases.* 2020.
12. Stevens DV, Tran AQ, Kim E. Complications of Orbital Emphysema in a COVID-19 Patient. *Ophthalmology.* 2020.
13. Wu P, Liang L, Chen C, Nie S. A child confirmed COVID-19 with only symptoms of conjunctivitis and eyelid dermatitis. *Graefes Arch Clin Exp Ophthalmol.* 2020.
14. Ying NY, Idris NS, Muhamad R, Ahmad I. Coronavirus Disease 2019 Presenting as Conjunctivitis. *Korean J Fam Med.* 2020.
15. Hu Y, Chen T, Liu M, et al. Positive detection of SARS-CoV-2 combined HSV1 and HHV6B virus nucleic acid in tear and conjunctival secretions of a non-conjunctivitis COVID-19 patient with obstruction of common lacrimal duct. *Acta Ophthalmol.* 2020.
16. Huang L, Zhang X, Zhang X, et al. Rapid asymptomatic transmission of COVID-19 during the incubation period demonstrating strong infectivity in a cluster of youngsters aged 16-23 years outside Wuhan and characteristics of young patients with COVID-19: A prospective contact-tracing study. *J Infect.* 2020.
17. Chen L, Deng C, Chen X, et al. Ocular manifestations and clinical characteristics of 535 cases of COVID-19 in Wuhan, China: a cross-sectional study. *Acta Ophthalmol.* 2020.
18. Guemes-Villahoz N, Burgos-Blasco B, Vilela AA, et al. SARS-CoV-2 RNA detection in tears and conjunctival secretions of COVID-19 patients with conjunctivitis. *J Infect.* 2020.
19. Bostanci Ceran B, Ozates S. Ocular manifestations of coronavirus disease 2019. *Graefes Arch Clin Exp Ophthalmol.* 2020.

20. Hong N, Yu W, Xia J, Shen Y, Yap M, Han W. Evaluation of ocular symptoms and tropism of SARS-CoV-2 in patients confirmed with COVID-19. *Acta Ophthalmol.* 2020.
21. Karimi S, Arabi A, Shahraki T, Safi S. Detection of severe acute respiratory syndrome Coronavirus-2 in the tears of patients with Coronavirus disease 2019. *Eye (Lond).* 2020.
22. Xia J, Tong J, Liu M, Shen Y, Guo D. Evaluation of coronavirus in tears and conjunctival secretions of patients with SARS-CoV-2 infection. *J Med Virol.* 2020.
23. Xie H-T, Jiang S-Y, Xu K-K, et al. SARS-CoV-2 in the ocular surface of COVID-19 patients. *Eye Vis (Lond).* 2020.
24. Kumar K, Prakash AA, Gangasagara SB, et al. Presence of viral RNA of SARS-CoV-2 in conjunctival swab specimens of COVID-19 patients. *Indian J Ophthalmol.* 2020.
25. Lan QQ, Zeng SM, Liao X, Xu F, Qi H, Li M. [Screening for novel coronavirus related conjunctivitis among the patients with corona virus disease-19]. *Zhonghua Yan Ke Za Zhi.* 2020.
26. Wu P, Duan F, Luo C, et al. Characteristics of Ocular Findings of Patients With Coronavirus Disease 2019 (COVID-19) in Hubei Province, China. *JAMA Ophthalmol.* 2020.
27. Zhang X, Chen X, Chen L, et al. The evidence of SARS-CoV-2 infection on ocular surface. *Ocul Surf.* 2020.
28. Zhou Y, Duan C, Zeng Y, et al. Ocular Findings and Proportion with Conjunctival SARS-COV-2 in COVID-19 Patients. *Ophthalmology.* 2020.
29. Deng C, Yang Y, Chen H, et al. Low risk of SARS-CoV-2 transmission through the ocular surface. *Acta Ophthalmol.* 2020.
30. Seah IYJ, Anderson DE, Kang AEZ, et al. Assessing Viral Shedding and Infectivity of Tears in Coronavirus Disease 2019 (COVID-19) Patients. *Ophthalmology.* 2020.
31. Hamroush A, Qureshi M, Shah S. Increased risk of ocular injury seen during lockdown due to COVID-19. *Cont Lens Anterior Eye.* 2020.
32. Pellegrini M, Roda M, Di Geronimo N, Lupardi E, Giannaccare G, Schiavi C. Changing trends of ocular trauma in the time of COVID-19 pandemic. *Eye (Lond).* 2020.
33. Pellegrini M, Roda M, Lupardi E, Di Geronimo N, Giannaccare G, Schiavi C. The impact of COVID-19 pandemic on ophthalmological emergency department visits. *Acta Ophthalmol.* 2020.
34. Loffredo L, Pacella F, Pacella E, Tiscione G, Oliva A, Violi F. Conjunctivitis and COVID-19: A meta-analysis. *J Med Virol.* 2020.
35. Ulhaq ZS, Soraya GV. The prevalence of ophthalmic manifestations in COVID-19 and the diagnostic value of ocular tissue/fluid. *Graefes Arch Clin Exp Ophthalmol.* 2020.
36. Liu M, Dai C, Lv X, Li B. Letter to the Editor: Are severe COVID-19 patients more susceptible to conjunctivitis? *J Med Virol.* 2020.