Supplementary materials

The effect of combination therapy on mortality and adverse events in patients with Staphylococcus aureus bacteraemia：A systematic review and meta-analysis of randomized controlled trials

Chao Ye • Chunjiang Wang • Zuojun Li • Xin Li • Juan Pan • Liang Liu • Zhaohui Wang

C. Ye

Department of Pharmacy, The Third Hospital of Changsha, Changsha, Hunan, People's Republic of China；

C.J. Wang

Department of Pharmacy, The Third Xiangya Hospital, Central South University, Changsha, Hunan, People's Republic of China；

Z.J. Li

Department of Pharmacy, The Third Xiangya Hospital, Central South University, Changsha, Hunan, People's Republic of China；

X. Li

Department of Pharmacy, The Third Hospital of Changsha, Changsha, Hunan, People's Republic of China；

J. Pan

Department of Pharmacy, Liuyang Hospital of Traditional Chinese Medicine, Changsha, Hunan, People's Republic of China;

L.Liu

Neurosurgery, People's Hospital of Ningxiang City Affiliated to Hunan University of Chinese Medicine, Changsha, Hunan, People's Republic of China;

Z.H. Wang (Corresponding)

Department of Pharmacy, People's Hospital of Ningxiang City Affiliated to Hunan University of Chinese Medicine, Changsha, Hunan, People's Republic of China.

Corresponding to : Department of Pharmacy, People's Hospital of Ningxiang City Affiliated to Hunan University of Chinese Medicine, Yihuan West Road, Ningxiang City, Hunan 410600, People's Republic of China. E-mail: wangzhaohuiry@163.com




Figure S1. Forest plot of the risk ratio (RR) for all-cause mortality in patients with Staphylococcus aureus bacteremia (SAB) at different recording time.




Figure S2. Forest plot of the risk ratio (RR) for any serious adverse events (SAEs) in patients with Staphylococcus aureus bacteremia (SAB)




Figure S3. Forest plot of the risk ratio (RR) for any adverse events (AEs) leading to treatment discontinuation in patients with Staphylococcus aureus bacteremia (SAB).

Table S1. Study characteristics.

Cheng MP[1], 2021	STAN:4 (3-8.5) COMBO:5(3-9)	STAN:1 (0-3.5) COMBO:2(0-4)	STAN:21.6 COMBO:26.4	STAN:7.8 COMBO:5.7	STAN:15.7 COMBO:7.7	Unknown	STAN:13.7 COMBO:9.4	Unknown	Unknown	STAN:76.5 COMBO:81.1
Pujol M[2], 2021	STAN:4 (2-5.8) COMBO:3(2-5.8)	Unknown	STAN:23.5 COMBO:13.5	Unknown	Unknown	STAN:9.9 COMBO:18.9	STAN:11.1 COMBO:12.2	STAN:48.1 COMBO:41.9	Unknown	Unknown
Tong SYC[3], 2020	STAN:5(2-7) COMBO:5(2-7)	STAN:1 (0-4) COMBO:2(1-4)	STAN:23 COMBO:28	STAN:6 COMBO:7	STAN:15 COMBO:18	Unknown	STAN:3 COMBO:5	STAN:12 COMBO:14	Unknown	Unknown
GeriakM[4], 2019	STAN:6 COMBO:5	Unknown	STAN:35 COMBO:53	STAN:26 COMBO:6	STAN:17 COMBO:29	STAN:17 COMBO:18	STAN:4 COMBO:18	STAN:13 COMBO:6	Unknown	Unknown
Thwaites GE [5],2018	STAN: 2 (0-3) COMBO: 1 (0-3)	STAN:2 (1-4) COMBO:2(1-4)	STAN:17 COMBO:19	Unknown	STAN:10 COMBO:10	Unknown	STAN:4 COMBO:5	STAN:17 COMBO:19	STAN:41 COMBO:38	Unknown
Davis JS[6], 2016	STAN: 3 (1-5) COMBO: 2(1-6)	STAN:1 (1-3) COMBO:1(0-2)	STAN:21 COMBO:23	STAN:10 COMBO:0	STAN:3 COMBO:19	STAN:7 COMBO:3	STAN:0 COMBO:3	STAN: 7 COMBO:19	Unknown	Unknown
Fowler VG[7], 2006	Unknown	Unknown	Unknown	STAN: 0 COMBO:0	STAN:17.5 COMBO:9.6	Unknown	STAN:75.0 COMBO:79.1	Unknown	Unknown	STAN:53 COMBO:50
Ruotsalainen E[8],2006	Unknown	Unknown	STAN:69* COMBO:59	STAN:35 COMBO:34	STAN:47 COMBO:37	STAN:8 COMBO:6	STAN:20 COMBO:16	STAN:8 COMBO:12	STAN:87 COMBO:81	Unknown
Ribera E[9], 1996	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown	STAN:100 COMBO:100	Unknown	Unknown	Unknown
Levine DP[10], 1991	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown	STAN:100 COMBO:100	Unknown	Unknown	Unknown
Vander Auwera P[11],1985	Unknown	Unknown	STAN:18.8 COMBO:46.2	STAN:12.5 COMBO:15.4	STAN:18.8 COMBO:15.4	Unknown	Unknown	STAN:25 COMBO:23.1	Unknown	Unknown

Vander Auwera P[12],1983	Unknown	Unknown	STAN:11.1 COMBO:0	STAN:11.1 COMBO:20	Unknown	Unknown	Unknown	STAN:30 COMBO:10	Unknown	Unknown
Korzeniowski O[13],1982	Unknown	Unknown	STAN:42.9 COMBO:44.2	Unknown	STAN:0 COMBO:4.7	Unknown	STAN:100 COMBO:100	STAN:5.7 COMBO:2.3	Unknown	Unknown
Abrams B[14], 1979	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown	STAN:100 COMBO:100	Unknown	Unknown	Unknown

COMBO: combination treatment ; STAN: standard therapy alone; IQR: interquartile range； *P=0.05

REFERENCES

1. Cheng MP, Lawandi A, Butler-Laporte G, et al. Adjunctive Daptomycin in the Treatment of Methicillin-susceptible Staphylococcus aureus Bacteremia: A Randomized, Controlled Trial. Clin Infect Dis 2021;72(9):e196–e203.
2. Pujol M, Miró JM, Shaw E, et al. Daptomycin Plus Fosfomycin Versus Daptomycin Alone for Methicillin-resistant Staphylococcus aureus Bacteremia and Endocarditis: A Randomized Clinical Trial.Clin Infect Dis 2021;72(9):1517–1525.
3. Tong SYC, Lye DC, Yahav D, et al. Effect of Vancomycin or Daptomycin With vs Without an Antistaphylococcal beta-Lactam on Mortality, Bacteremia, Relapse, or Treatment Failure in Patients With MRSA Bacteremia: A Randomized Clinical Trial. JAMA 2020;323(6):527–537.
4. Geriak M, Haddad F, Rizvi K, et al. Clinical Data on Daptomycin plus Ceftaroline versus Standard of Care Monotherapy in the Treatment of Methicillin-Resistant Staphylococcus aureus Bacteremia. Antimicrob Agents Chemother 2019;63(5):e02483–18.
5. Thwaites GE, Scarborough M, Szubert A, et al. Adjunctive rifampicin for Staphylococcus aureus bacteraemia (ARREST): a multicentre, randomised, double-blind, placebo-controlled trial. Lancet 2018; 391(10121): 668–78.
6. Davis JS, Sud A, O'Sullivan MVN, et al. Combination of Vancomycin and beta-Lactam Therapy for Methicillin-Resistant Staphylococcus aureus Bacteremia: A Pilot Multicenter Randomized Controlled Trial. Clin Infect Dis 2016;62(2):173–180.
7. Fowler VG Jr, Boucher HW, Corey GR, et al. Daptomycin versus standard therapy for bacteremia and endocarditis caused by Staphylococcus aureus. N Engl J Med 2006;355(7):653–65.
8. Ruotsalainen E, Järvinen A, Koivula I, et al. Levofloxacin does not decrease mortality in Staphylococcus aureus bacteraemia when added to the standard treatment: a prospective and randomized clinical trial of 381 patients. J Intern Med 2006;259(2):179–90.
9. Ribera E, Gómez-Jimenez J, Cortes E, et al. Effectiveness of cloxacillin with and without gentamicin in short-term therapy for right-sided Staphylococcus aureus endocarditis.A randomized, controlled trial.Ann Intern Med 1996;125(12):969–74.
10. Levine DP, Fromm BS, Reddy BR. Slow response to vancomycin or vancomycin plus rifampin in methicillin-resistant Staphylococcus aureus endocarditis. Ann Intern Med 1991;115(9):674–80.
11. Van der Auwera P, Klastersky J, Thys JP, et al. Double-blind, placebo-controlled study of oxacillin combined with rifampin in the treatment of staphylococcal infections. Antimicrob Agents Chemother 1985;28(4):467–72.
12. Van der Auwera P, Meunier-Carpentier F, Klastersky J. Clinical study of combination therapy with oxacillin and rifampin for staphylococcal infections. Rev Infect Dis 1983;5 Suppl 3:S515–22.
13. Korzeniowski O, Sande MA. Combination antimicrobial therapy for Staphylococcus aureus

endocarditis in patients addicted to parenteral drugs and in nonaddicts: A prospective study. Ann Intern Med 1982;97(4):496–503.

14. Abrams B, Sklaver A, Hoffman T, et al. Single or combination therapy of staphylococcal endocarditis in intravenous drug abusers. Ann Intern Med 1979;90(5):789–91.