Supplement Table 3. Description of interventions

Study	Intervention(s) (route, frequency, total dose/day)	Comparator(s) (route, frequency, total dose/day)
Choi et al [11], 1995	I1: KRG (tablet, NR, 1,800 mg)	C1: Placebo (NR in detail) C2: Trazodon (tablet, 25 mg daily at bedtime)
Choi et al [8], 1999	I1: KRG (tablet, 3 times, 1,800 mg)	C1: Placebo (tablet, 3 times, same shape)
Choi and Choi [9], 2001	I1: KRG (tablet, 3 times, 1,800 mg)	C1: Placebo (same shape)
Choi et al [10], 2003	I1: KRG (tablet, 3 times, 1,800 mg)	C1: Placebo (NR in detail)
de Andrade et al [12], 2007	I1: KRG (capsule, 3 times, 3,000 mg)	C1: Placebo (capsule containing starch with KRG flavour)
Ham et al [13], 2009	I1: KRG plus ginsenoside (capsule, 2 times, 800 mg)	C1: Placebo (capsule, microcrystalline cellulose 200 mg)
Hong et al [14], 2002	I1: KRG (NR, 3 times, 2,700 mg)	C1: Placebo (NR in detail)
Kim and Paick [15], 1999	I1: KRG (capsule, 3 times, 2,700 mg)	C1: Placebo (same shape and smell)
Kim et al [16], 2009	I1: TCMG (NR, 2 times, 2,000 mg)	C1: Placebo (NR in detail)

C: comparator, I: intervention, KRG: Korean red ginseng, TCMG: Tissue-cultured mountain ginseng, NR: not reported.