https://doi.org/10.5534/wjmh.210155

Supplement Table 1. Characteristics of included studies

Study	Intervention(s) and comparator(s)	Duration of intervention	Trial period	Country	Setting	Age in years (mean)	Baseline IELT in minutes (mean)	Number of participants with primary/secondary PE
Ahn 1996 [12]	I1: fluoxetine 20 mg daily for first 1 week and 40 mg daily for remaining 5 weeks after breakfast	6 weeks	NR	South Korea	Outpatient	39.8 (range 34–48)	0.78 (range 0.17–2.0)	NA/NA
	C1: placebo daily					39.8 (range 34–48)	0.78 (range 0.17–2.0)	NA/NA
Athanasios 2007 [13]	I1: duloxetine 20 mg daily for 1 week followed by 40 mg daily	12 weeks	NR	Greece	Academic	31.35 (SD 8.23)	0.63 (SD 0.27)	NA/NA
	C1: placebo daily					32.65 (SD 7.49)	0.58 (SD 0.30)	NA/NA
Atmaca 2002 [14]	I1: citalopram 20 mg daily up to 60 mg	8 weeks	NR	Turkey	Outpatient	Range 24–46	0.55 (SD 0.29)	NA/NA
	C1: placebo daily					Range 24–46	0.50 (SD 0.24)	NA/NA
Biri 1998 [15]	I1: sertraline 50 mg	4 weeks	1995–1997	Turkey	Outpatient	NR	0.68 (SD 0.21)	NA/NA
	C1: placebo daily					NR	0.72 (SD 0.33)	NA/NA
Buvat 2009 [16]	I1: dapoxetine 30 mg on-demand	24 weeks	2004–2006	France	Academic	39.6 (SD 9.53)	0.9 (SD 0.50)	NA/NA
	I2: dapoxetine 60 mg on-demand					40.5 (SD 9.62)	0.9 (SD 0.49)	NA/NA
	C1: placebo daily					40.1 (SD 9.98)	0.9 (SD 0.51)	NA/NA
Farnia 2009 [17]	I1: citalopram 20 mg on-demand	4 weeks	2006–2007	Iran	Outpatient	34.28 (SD 6.67)	1.11 (SD 0.61)	NA/NA
	C1: placebo daily					33.76 (SD 5.93)	1.10 (SD 0.56)	NA/NA
Gameel 2018 [18], 2013 [19], 2013 [20]	I1: paroxetine 20 mg on-demand+lubricating jelly	4 weeks	2009–2012	Egypt	Outpatient	NR	0.16 (SD 0.47)	NA/NA
	C1: placebo on-demand+lubricating jelly							
Gong 2011 [21]	I1: paroxetine 20 mg daily	30 days	NR	China	Likely outpatient	26.8 (SD 5.5)	0.89 (SD 0.21)	NA/NA
	C1: placebo daily					29.2 (SD 6.7)	0.97 (SD 0.18)	
Hamidi Madani 2016 [22], 2018 [23]	I1: tramadol 50 mg	12 weeks	NR	Iran	Outpatient	NR	NR	NA/NA
	I2: paroxetine 20 mg					NR	NR	NA/NA
	C1: placebo					NR	NR	NA/NA
Kara 1996 [24]	I1: fluoxetine 20 mg daily	NR	NR	Turkey	Outpatient	Range 15–50	0.42 (SD 0.21)	NA/NA
	C1: placebo daily					Range 15–50	0.5 (SD 0.14)	NA/NA
Kaufman 2009 [25]	I1: dapoxetine 60 mg on-demand	9 weeks	NR	USA and Canada	Outpatient	41.8 (SD 9.80)	NR	NA/NA
	C1: placebo on-demand					40.98 (SD 9.71)	NR	NA/NA
Khelaia 2015 [26], 2014 [27], 2012 [28]	I1: paroxetine 20 mg	4 weeks	NR	Georgia	Academic	22.7 (range 19–39)	NR	NA/NA
	I2: paroxetine 20 mg 2–3 hours before intercourse					22.7 (range 19–39)	NR	NA/NA
	C1: placebo					22.7 (range 19–39)	NR	NA/NA
Kim 1998 [29]	I1: fluoxetine 40 mg daily for 1 week then 80 mg for 3 weeks	16 weeks	NR	South Korea	Academic	44 (range 30–60)	0.77 (SD 0.68)	NA/NA
	I2: sertraline 100 mg for 1 week then 200 mg for 3 weeks					44 (range 30–60)	0.77 (SD 0.68)	NA/NA
	C1: placebo daily					44 (range 30–60)	0.77 (SD 0.68)	NA/NA

The World Journal of MEN's HEALTH

Supplement Table 1. Continued 1

Study	Intervention(s) and comparator(s)	Duration of intervention	Trial period	Country	Setting	Age in years (mean)	Baseline IELT in minutes (mean)	Number of participants with primary/secondary PE
Mattos 2008 [30]	I1: fluoxetine 90 mg daily	12 weeks	NR	Brazil	Academic	50 (SD 8.51)	0.94 (SD 0.31)	NA/NA
	I2: fluoxetine 90 mg daily+tadalafil 20 mg on-demand					42.81 (SD 7.73)	0.83 (SD 0.43)	NA/NA
	C1: placebo daily					45.93 (SD 9.96)	0.83 (SD 0.31)	NA/NA
	C2: placebo+tadalafil 20 mg on-demand					43.2 (SD 11.3)	0.83 (SD 0.32)	NA/NA
McMahon 1998 [31]	I1: sertraline 50 mg	12 weeks	NR	Australia	Academic	41 (range 19–70)	0.3	NA/NA
	C1: placebo daily					41 (range 19–70)	0.3	NA/NA
McMahon 1999 [32]	Study 1	17 weeks	NR	Australia	Academic	39.5	0.3	19/7
	I1: paroxetine 20 mg					39.5	0.3	19/7
	C1: paroxetine as needed 3–4 hours before planned sexual intercourse							
	Study 2					40.5	0.5	32/10
	I2: paroxetine 10 mg for 3 weeks then 20 mg paroxetine as needed for 4 weeks					40.5	0.5	32/10
	C2: placebo daily for 3 weeks then placebo daily for 4 weeks							
McMahon 2010 [33,34]	I1: dapoxetine 30 mg on-demand	12 weeks	2005–2006	Multicenter in Asia/Pacific	Academic	41.2 (SD 10.74)	3.9	92 (42.2%)/NA
	I2: dapoxetine 60 mg on-demand					41.0 (SD 10.78)	4.2	92 (42.2%)/NA
	C1: placebo daily					40.6 (SD 9.71)	2.4	96 (45.9%)/NA
McMahon 2013 [35]	I1: dapoxetine 30 mg on-demand, from week 4 up to 60 mg if tolerated+PDE5 inhibitor taken 1–3 hours prior to sexual intercourse	12 weeks	2010–2011	Australia	Academic	49.5 (SD 11.23)	NR	92 (42.2%)/NA
						47.9 (SD 11.96)	NR	96 (45.9%)/NA
	C1: placebo daily+PDE5 inhibitor taken 1–3 hours prior to intercourse							
Mendels 1995 [36]	I1: sertraline 50 mg daily that could be titrated up to 200 mg daily	10 weeks	NR	USA	Academic	NR	0.98 (SD 1.15)	NA/NA
	C1: placebo daily					NR	1.10 (SD 1.35)	NA/NA
Na 1996 [37]	I1: sertraline 50 mg at night that could be titrated up to 100 mg daily	6 weeks	NR	South Korea	Academic	NR	NR	NA/NA
	C1: placebo daily					NR	NR	
Novaretti 2002 [38]	I1: fluoxetine 20 mg	20 weeks	1998–2000	Brazil	Academic	37.4 (SD 10.7)	1.01 (SD 0.86)	NA/NA
	C1: placebo daily					37.4 (SD 10.7)	1.05 (SD 1.07)	NA/NA

https://doi.org/10.5534/wjmh.210155

Supplement Table 1. Continued 2

Study	Intervention(s) and comparator(s)	Duration of intervention	Trial period	Country	Setting	Age in years (mean)	Baseline IELT in minutes (mean)	Number of participants with primary/secondary PE
Pryor 2006 [39]	I1: dapoxetine 30 mg on-demand 1–3 hours before anticipated sexual activity	12 weeks	2003–2004	USA	Academic	40.3 (SD 9.10)	0.90 (SD 0.47)	563/227
	I2: dapoxetine 60 mg on-demand 1–3 hours before anticipated sexual activity					40.9 (SD 9.09)	0.92 (SD 0.50)	571/234
	C1: placebo on-demand 1–3 hours before anticipated sexual activity					40.3 (SD 9.55)	0.91 (SD 0.48)	560/248
Safarinejad 2006 [40]	I1: dapoxetine 60 mg daily	12 weeks	2003–2005	Iran	Academic	33.4 (range 20–50)	0.63	64 (61.5%)/NA
	I2: paroxetine 20 mg daily					34.6 (range 21–49)	0.52	63 (60.0%)/NA
	C1: placebo daily					34.3 (range 21–50)	0.57	11 (44.0%)/NA
Safarinejad 2006 [43]	I1: citalopram 30 mg	6 months	NR	Iran	Academic	32 (21–49)	0.53	10/16
	C1: placebo daily					34 (21–49)	0.47	11/14
Safarinejad 2007 [41]	I1: escitalopram 10 mg daily	12 weeks	2003–2005	Iran	Academic	33.5 (range 21–44)	NR	87 (70%)/NA
	C1: placebo daily					33.3 (range 19–46)	NR	88 (69.8%)/NA
Safarinejad 2008 [42]	I1: dapoxetine 30 mg daily	12 weeks	2004–2006	Iran	Academic	35.7 (range 21–54)	0.37	40 (37.7%)/NA
	C1: placebo daily					36.3 (range 19–56)	0.48	43 (40.6%)/NA
Shang 2012 [44]	I1: citalopram 20 mg daily	4 weeks	2011–2012	China	Academic	39.1 (SD 2.5)	0.91 (SD 0.18)	NA/NA
	C1: placebo daily					37.8 (SD 2.8)	0.95 (SD 0.17)	
Tuncel 2008 [45]	I1: sertraline 50 mg nightly for 2 months	8 weeks	NR	Turkey	Academic	36.9 (median) (SD 6.9)	NR	NA/NA
	C1: placebo daily					34.9 (median) (SD 9.0)	NR	NA/NA
Waldinger 1994 [46]	I1: paroxetine 20 mg daily for 1 week and then 40 mg daily from week 2–6	6 weeks	NR	The Netherlands	Outpatient	41 (range 27–48)	NR	7/8 (87.5%)/NA
	C1: placebo daily					38 (range 30–47)	NR	7/9 (77.7%)/NA
Waldinger 1999 [47]	I1: fluoxetine 20 mg daily	6 weeks	NR	The Netherlands	Outpatient	38 (SD 7.0)	0.3 (SD 0.22)	NA/NA
	I2: fluvoxamine 100 mg daily					44 (SD 10.0)	0.3 (SD 0.22)	NA/NA
	I3: paroxetine 20 mg daily					41 (SD 8.0)	0.3 (SD 0.22)	NA/NA
	I4: sertraline 50 mg daily					40 (SD 9.0)	0.3 (SD 0.22)	NA/NA
	C1: placebo daily					45 (SD 4.0)	0.3 (SD 0.22)	NA/NA
Yilmaz 1999 [48]	I1: fluoxetine 20 mg daily	1 month	1997–1997	Turkey	Academic	36.5 (range 22–56)	1.2 (SD 1.0)	NA/NA
	C1: placebo					37.3 (range 24–58)	1.1 (SD 1.1)	NA/NA

IELT: intravaginal ejaculatory latency time, PE: premature ejaculatione, NR: not reported, NA: not available, SD: standard deviation.