Supplementary - Database search algorithms

The search algorithms uesd in this study are presented below. All database were searched from January 1st 2011 until January 6th 2022. The number of retrieved items is presented next to each database.

Search for PubMed (n=1080):

((brugada) OR (brugada syndrome)) AND((prognosis) OR (risk assessment) OR (predictive value) OR (outcome)), 2011-01-01~2022-01-06

Search for Embase (n=1421):

(brugada or brugada syndrome).af. And (prognosis or risk assessment or predictive value or outcome).af. limit 3 to (human and english language and yr="2011 -Current")

Search for Medline (n=874):

(brugada or brugada syndrome).af. And (prognosis or risk assessment or predictive value or outcome).af. (english language and humans and yr="2011 -Current")

Search for Web of Science (n=1106):

(TS=(brugada OR brugada syndrome)) AND TS=(prognosis OR risk assessment OR predictive value OR outcome), 2011-01-01~2022-01-06

Supplementary Table S1 : Individual prognostic variables in the 3 risk models.

Model	Risk factor		Relative value	Categories
Delise Score	Spontaneous type 1 ECG		-	Individuals at higher risk are those with a baseline type 1 ECG pattern who have at least 2 of the following risk factors: - Syncope - Family history of SCD - Positive EPS A score > 2 showed a significantly higher event probability
	Syncope		-	
	Family history of SCD		-	
	Positive EPS		-	
Sieira Score	Pontaneous type 1 ECG		1	
	SCA		4	
	Syncope		2	
	Early familial SCA < 35yo		1	
	Inducible EPS		2	
	SND		3	
Shanghai Score	. ECG (12-lead/ambulatory)	Spontaneous type 1 ECG at nominal or high leads	3.5	- Moderate risk: 3.5 - High risk: 4 to 5 - Highest risk ≥5.5
		Fever-induced type 1 ECG at nominal or high leads	3	
		Type 2 or 3 ECG pattern that converts with drug provocation challenge	2	
	Clinical history	Unexplained cardiac arrest or documented VF/polymorphic VT	3	

	Nocturnal agonal respirations	2
	Suspected arrhythmic syncope	2
	Syncope of unclear mechanism/unclear etiology	1
	Atrial flutter/fibrillation in patients < 30 y without alternative etiology	0.5
	First- or second-degree relative with definite BrS	2
Family history	Suspicious SCD (fever, nocturnal, Brugada aggravating drugs) in a first- or second-degree relative	1
	Unexplained SCD at	0.5
Genetic test result	Probable pathogenic mutation in BrS susceptibility gene	0.5

Supplementary Figure S1 : Subgroup analysis of Sieira model between prospective and retrospective studies.