SUPPLEMENTAL MATERIAL

Table of Contents

Table S1. PRISMA 2020 item checklist
Table S2. Search Strategy
Table S3. Indirect health utility metrics
Table S4. Characteristics of studies included in systematic review
Table S5. Risk of bias assessment criteria
Table S6. Weighted baseline characteristics across studies
Table S7. Weighted baseline characteristics across studies for EQ-5D-3L
Table S8. Risk of bias assessment
Table S9. Summary statistics of risk of bias assessment
Table S10. Pooled EQ-5D-3L values for different time ranges and time points
Table S11. Sensitivity analyses
Table S12. Comparison of summary estimates using random effects and fixed effect meta-analysis
Figure S1. PRISMA flowchart
Figure S2. Characteristics of included papers
Figure S3. Random effects meta-analysis for EQ-5D-3L utility values
Figure S4. Fixed effects meta-analysis for EQ-5D-3L utility values
Figure S5. Random effects meta-analysis for EQ-5D-5L
Figure S6. Random effects meta-analysis for AQoL
Figure S7. Random effects meta-analysis for HUI2
Figure S8. Random effects meta-analysis for HUI3
Figure S9. Random effects meta-analysis for 15D
Figure S10. Random effects meta-analysis for SF-6D
Figure S11. Sex-stratified random effects meta-analysis for EQ-5D-3L
Figure S12. Age-stratified random effects meta-analysis for EQ-5D-3L
Figure S13. Stroke type-stratified random effects meta-analysis for EQ-5D-3L
Figure S14. Time-stratified random effects meta-analysis for EQ-5D-3L
Figure S15. Meta-regression analyses of mean EQ-5D-3L utility score by sex, age, and publication date
References – studies included in systematic review

Table S1. PRISMA 2020 item checklist

Section and Topic	Item #	Checklist item	
TITLE			Yes
Title	1	Identify the report as a systematic review.	
ABSTRACT			Reviewed/compliant
Abstract	2	See the PRISMA 2020 for Abstracts checklist (Table 2).	
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	Page 2
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	Page 2
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	Page 3
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	Page 2
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	Supplemental Table 1
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	Supplemental Table 1
Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	Page 2 and 3
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	Page 3
	10b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	Page 2 and 3
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results.	Page 4
Synthesis methods	13a	Describe the processes used to decide which studies were eligible for each synthesis.	
	13b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	Page 4
	13c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	
	13d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	
	13e	Describe any methods used to explore possible causes of heterogeneity among study results.	
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	

Section and Topic	Item #	Checklist item	
Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	Page 3 and Supp Table 2
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	Page 4
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram (see Figure 1).	Page 4 and Supp Figure 1
	16b	Cite studies that met many but not all inclusion criteria ('near-misses') and explain why they were excluded.	
Study characteristics	17	Cite each included study and present its characteristics.	Supp Table 3
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	Supp Table 5
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	Figure 1-3, Supp Figures 3-12
Results of syntheses	20a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	
	20b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	Page 5-6 Figure 1-3 Supp Table 6
	20c	Present results of all investigations of possible causes of heterogeneity among study results.	
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	Supp Table 5-6
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	Page 6
DISCUSSION			
Discussion	23a	Provide a general interpretation of the results in the context of other evidence.	
	23b	Discuss any limitations of the evidence included in the review.	Page 6-8
	23c	Discuss any limitations of the review processes used.	
	23d	Discuss implications of the results for practice, policy, and future research.	
OTHER INFORMATION			
Registration and protocol	24a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	
	24b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	Page 2
	24c	Describe and explain any amendments to information provided at registration or in the protocol.	
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	Acknowledgements
Competing interests	26	Declare any competing interests of review authors.	Disclosures
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	Page 4

Table S2. Search Strategy

Search Strategy 1: Global search (using Medline syntax)

#1	exp Stroke/ OR exp Cerebral Infarction/ or exp Brain Infarction/ or exp brain ischemia/ or cerebral hemorrhage/ OR (stroke* OR cerebral infarction OR brain infarction OR brain ischemia OR h?emorrhagic stroke* or cerebral h?emorrhage* or intracerebral h?emorrhage* or brain h?emorrhage*).tw,kf.
#2	exp "Quality of Life"/ OR (euroqol or euro qol or eq5d* eq-5d* or short form 6 dimension or short form six dimension or sf-6D* or hui or hui2 or hui3 or health utilit* or assessment of quality of life or aqol* or quality of well being or qwb or 15D).tw,kf.
#3	(#1) AND (#2)
#4	(#3) and NOT (address or autobiography or bibliography or biography or case reports or dataset or dictionary or directory or duplicate publication or editorial or "expression of concern" or festschrift or interactive tutorial or interview or lecture or legal case or legislation or news or newspaper article or patient education handout or periodical index or personal narrative or portrait or technical report or twin study or video-audio media or webcast or letter)
#5	1995 until day October 26, 2020

Table S3. Indirect health utility metrics

HRQoL questionnaire	Description	Range of possible scores
EuroQol five dimensions three levels (EQ-5D-3L)[22 (main text)]	Five dimensions (mobility, self-care, usual activities, pain/discomfort, anxiety/depression), each of which is assigned one of three levels, allowing 243 health states	-0.59 to 1.00 (Europe)
EuroQol five dimensions five levels (EQ-5D-5L)[23 (main text)]	Adaptation of EQ-5D-5L with 5 levels for each dimension, allowing 3,125 possible health states	-0.28 to 1.00 (UK)
Short form six dimensions (SF-6D)[26 (main text)]	Six dimensions (physical functioning, role limitations, social functioning, pain, mental health, vitality), with four to six levels, allowing 18 000 health states.	0.30 to 1.00 (UK)
Health utilities index mark 2 (HUI2)[25 (main text)]	Seven dimensions (sensory, mobility, emotion, cognitive, self-care, pain, fertility), with three to five levels, allowing 24 000 health states	-0.03 to 1.00 (Canada)
Health utilities index mark 3 (HUI3)[25 (main text)]	Eight dimensions (vision, hearing, speech, ambulation, dexterity, emotion, cognition, pain), with five to six levels, allowing 972 000 health states	-0.36 to 1.00 (Canada)
15D (15 dimensions)[28 (main text)]	Fifteen dimensions (mobility, vision, hearing, breathing, sleeping, eating, speech, elimination, usual activities, mental function, discomfort and symptoms, depression, distress, vitality, and sexual activity), with five grades of severity, allowing 3.1×10^{10} health states.	0 to 1 (Finland)
Assessment of Quality of Life (AQOL)[24 (main text)]	Five dimensions (illness, independent living, social relationships, physical senses, psychological well-being), with four levels, allowing for 1.1 billion health states.	0 to 1 (Australia)
Quality of Well-Being (QWB)*[27 (main text)]	Three dimensions (mobility, physical activity, social activity).	0 to 1 (USA)

*****No included studies in this systematic review had QWB

Table S4. Characteristics of studies included in systematic review

Author	Citation	Publication Date	Translated?	Country	Stroke type	Self-report or diagnosis	Percent ischemic	Percent left hemisphere	Location	Mean or median age	Percent female	Time since stroke	Mean or Median NIHSS	Percent hypertension	Percent diabetes	Percent smoking	Percent coronary disease	Percent prior stroke	Percent atrial fibrillation	Instrument used	If EQ, 3L or 5L?	Tariff used	Survey method
Adey-Wakeling	73	2016	No	Australia	B	D	88.3	NR	C	NR	48.5	At 12 months	NR	NR	NR	NR	NR	NR	NR	E	3	Australia	I
Appau	74	2019	No	Canada	B	D	NR	NR	C	71.5 (12.8)	NR	At 3 months	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Arrospide	75	2019	No	Spain	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	5	Spanish	I
Arwert	76	2017	No	Netherlands	B	D	83	50%	C	47.7 (9.7)	37	Mean 36 months (11.4)	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M/P
Barton	77	2008	No	UK	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E, 6D	3	UK	M
Broussy	78	2019	No	France	B	D	88	NR	C	69 (14)	38	At 12 months	Mean 4.7 (5.5)	61	16	32	NR	13	14	E	3	France	M/P
Burton	79	2014	No	UK	B	D	94	NR	C	70.78 (11.12)	50	Mean 27.5 months (6.6)	NR	NR	NR	NR	NR	NR	NR	E	3	UK	M
Bushnell	11	2014	No	USA	I	D	77.4	NR	C	65 (IQR 56-75)	46.3	At 3 and 12 months	Mean 3.78 (4.59)	79.2	27.9	24.7	24.2	23	10.8	E	3	USA	P
Cadilhac 1	80	2019	No	Australia	B	D	77.7	NR	C	78 (IQR 69-85) stroke unit, 76 (IQR 64-85) Non stroke unit	48.4	Median 101 days (97-107)	NR	NR	NR	NR	24.2	24	NR	E	3	Australia	I
Cadilhac 2	81	2010	No	Australia	B	D	NR	NR	C	NR	49.2	At 12 months and 5 years	NR	NR	NR	NR	NR	NR	NR	A		NR	?
Cao	82	2012	Y	China	B	D	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Chang	83	2016	No	Korea	B	D	80.1	NR	C	64.3 (12.8)	40.4	At 6 months	NR	54.4	23.2	27.6	6.4	NR	10.3	E	3	NR	I
Chen 1	84	2015	No	Taiwan	U	D	NR	50.7	R	52.8 (11.6)	26	Median 19.7 months (0.4-94)	NR	NR	NR	NR	NR	NR	NR	E	5	Japan	I
Chen 2	85	2018	No	USA	H	D	NR	NR	C	65 (IQR 57-75) resumption group/ 62 (IQR 53.5-73) no resumption group	38.8	At 3 months	NR	90.1	31.7	43.2	5.5	11.5	12.2	E	3	NR	I
Cheung	86	2019	No	Singapore	B	D	87	NR	C	62.7 (10.9)	32.4	At 3 months or 12 months	NR	NR	NR	NR	NR	NR	NR	H3	N/A	Canadian	NR
Cramm	87	2012	No	Netherlands	U	D	NR	NR	H	69.13 (14.24)	49.8	NR	NR	NR	NR	NR	NR	NR	NR	E	3	Netherland	I
Cup	88	2001	No	Netherlands	B	D	NR	NR	C	68 (15)	57.7	At 6 months	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I

Darlington 1	89	2009	No	Netherlands	B	D	79	NR	H/C	60.9 (16.9)	51	Mean 10.9 months (1.19)	NR	NR	NR	NR	NR	NR	NR	E	3	Netherland	I/P
Darlington 2	90	2007	No	Netherlands	B	D	79	31	H/R/N	60.9 (16.9)	51	Mean 10.9 months (1.19)	NR	NR	NR	NR	NR	NR	NR	E	3	Netherland	I/P
de Graaf	91	2020	No	Netherlands	U	D	93	46	C	68.8 (11.7)	40	At 3 months	NR	NR	NR	NR	NR	NR	NR	E	5	Netherland	P
Deb-Chatterji	92	2020	No	Germany	I	D	100	NR	C	76 (IQR 65-81)	51.8	At 3 months	Median 15 (IQR 10.25-19)	NR	NR	NR	NR	NR	NR	E	3	UK	I/P
Dewilde	93	2019	No	Belgium	B	D	NR	NR	C	68.7 (12.9)	41.1	Mean 13.8 (30.8)	NR	73	20.9	NR	NR	13.3	23.6	E	3	European	I/P
Dorman 1	94	2000	No	UK	B	D	NR	NR	C	NR	NR	median 72 weeks (IQR 43-104)	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Dorman 2	95	1997	No	UK	U	D	NR	NR	C	NR	NR	At least 3 months	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Du	96	2018	Y	China	I	D	100	NR	H and C	61.7 (12.6)	39.3	3 months	NR	NR	NR	NR	NR	NR	NR	E, 6D	3	NR	I
Edwards	97	2010	No	Canada	B	S	NR	NR	C	NR	46.8	NR	NR	NR	18.3	NR	33.4	NR	NR	H3	N/A	Canada	NR
Espuela	98	2017	Yes	Spain	B	D	NR	NR	C	71.3 (11.9)	36.3	>4 years		64.5	23.4	8.1	38.2	NR	NR	E	3	NR	I
Fischer	99	2008	No	Switzerland	I	D	100	NR	C	61 (13)	50.3	Mean 2.53 years (1.18)	Median 14 (range 3-38)	54	11	17	NR	13	NR	E	3	NR	M
Ghatnekar	100	2013	No	Sweden	B	D	87	NR	C	71 (13) 2006 cohort/ 74 (11) 2009 cohort	45.8	At 3 months	NR	56.8	17.8	15.1	NR	24.2	24.8	E	3	UK	I/P
Golicki 1	101	2014	No	Poland	B	D	92.9	NR	H/C	70.6 (11)	51.8	At 4 months	NR	NR	NR	NR	NR	NR	NR	E	3, 5	Poland	NR
Golicki 2	9	2014	No	Poland	B	D	87.4	NR	H	69 (12.9)	48.5	Median 8 days (during hospitalization)	NR	NR	NR	NR	NR	NR	NR	E	3, 5	Poland	I
Graessel	102	2014	No	Germany	B	D	83.3	47.4	R	68.7 (11)	42.2	NR	NR	NR	NR	NR	NR	NR	NR	E	3	European	P
Grasel	103	2019	Y	Germany	B	D	83.3	47.5	R/C	68.7 (11)	42.2	Discharge rehab/2.5/ 5 years	NR	NR	NR	NR	NR	NR	NR	E	3	European	P
Groeneveld 1	104	2018	No	Netherlands	B	D	77	47.9	R	60.2 (12.7)	41.8	Mean 29 days	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M
Groeneveld 2	105	2018	No	Netherlands	B	D	75.3	47.2	R (inpatient and outpatient)	57.3 (11.8)	53	NR	Median NIH 6 (3-12.5)	38.6	15.5	28.9	5.9	NR	NR	E	3	Netherlands	P/E
Guo	106	2017	No	Singapore	B	D	87.8	NR	C	64.1 (9.96)	29.5	3 months and 12 months	NR	NR	NR	NR	NR	NR	NR	E	3	Singapore	I
Haacke	107	2005	No	Germany	B	D	70	NR	C	71.7 (11.3)	54.5	At 4 years	NR	NR	NR	NR	NR	NR	NR	E, H2, H3	3	NR	I
Hansson	108	2012	No	Sweden	B	D	NR	NR	C	75.2 (11.8)	48.4	At 12 months	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Hokstad	109	2016	No	Norway	B	D	85.6	NR	C	76.8 (11.3)	51.9	At 3 months	Mean NIHS 7.9 (7.7)	NR	NR	NR	NR	27.7	NR	E	5	Denmark	I/P

Jeon	110	2017	No	Korea	B	D	67	NR	C	65 (11)	45.7	Mean 35.5 days (37.2)	Mean NIH 6.3 (4.8)	60.9	30.4	NR	NR	NR	NR	E	NR, assumed 3	NR	I
Katona	111	2015	No	Germany	B	D	82	NR	C	67.4 (11.1)	40	Mean 57 (28) days after discharge from rehab	NR	NR	NR	NR	NR	NR	NR	E	3	European	P
Katzan	112	2017	No	USA	I	D	100	NR	C	63.5 (14.4)	46	Median 58 days (IQR 32-258)	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Kelly	113	2014	No	USA	I	D	100	27%	O	55 (IQR 42-62)	64	Mean for early: 3months, for late: 9 months	Median 17 (IQR 16-22)	NR	NR	NR	NR	NR	NR	E	3	NR	I
Khiaocharoen	114	2012	No	Thailand	B	D	56.5	NR	H	60.9 (12.6)	42	<2 weeks	NR	NR	NR	NR	NR	NR	NR	E	3	UK	I
Kil	115	2008	Y	Korea	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	NR
Kim 1	116	2019	No	Korea	B	S	NR	NR	C	64.4 (SE 0.7)	44.3	NR	NR	81.1	56.7	15.2	NR	NR	NR	E	3	NR	NR
Kim 2	117	2018	No	Korea	B	S	NR	NR	C	67.4 (SE 0.53)	50.5	NR	NR	NR	NR	48	NR	NR	NR	E	3	Korea	NR
Kuo	118	2017	No	Japan	U	S	NR	NR	C	75.8 (6.58)	36.7	NR	NR	77.3	35.5	NR	NR	NR	NR	E	3	Japan	I
Kuroda 1	119	2007	Y	Japan	B	D	NR	NR	C	NR	NR	Mean 39.6 months	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M
Kuroda 2	120	2003	Y	Japan	B	D	75	NR	C	71.6 (9.9)	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M
Kuwano	121	2001	Y	Japan	B	D	NR	NR	C	64.3 (12.4)	NR	Two groups; short term (1-2.6 yrs) and long term (4.3-5 years)	NR	NR	NR	NR	NR	NR	NR	E	3	NR	NR
Kwon	122	2018	No	Korea	B	S	NR	NR	C	64.8 (0.6 SE)	42.6	NR	NR	70.1	32.1	15.3	NR	NR	NR	E	3	Korea	NR
Labberton	123	2020	No	Norway	B	D	61.4	NR	C	71 (12.5)	39.2	At 3 and 12 months	Median 3	NR	NR	NR	NR	NR	NR	E	3	UK	NR
Lannin	124	2017	No	Australia	B	D	84	NR	C	NR	NR	Median 156 days	NR	NR	NR	NR	NR	NR	NR	E	3	Australia	P
Leach	125	2010	No	Australia	B	D	NR	NR	C	67.6 (13.8)	49.7	At 7 years	Median 3 (IQR 1-7)	55.8	18.3	16.2	15.6	12.2	11.3	A	N/A	NR	I
Lee	126	2010	No	Taiwan	B	D	76.3	NR	R/O	64.8 (12.2)	36.6	Mean 4.3 years (4.2)	NR	NR	NR	NR	NR	NR	NR	E	3	UK/US	I
Leeds	127	2004	No	UK	U	D	NR	NR	R/C/N	79.9 (7.3)	74	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Lindgren	128	2007	No	Sweden	B	D	NR	NR	C	64.4 (9.3)	40.4	At 3/6/9/12 months	NR	NR	NR	NR	NR	NR	NR	E	3	UK	M
Lopez-Bastida	129	2012	No	Spain	U	D	NR	NR	C	67.1 (12.2)	43.3	1/2/3 years	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M
Lu	130	2016	No	Taiwan	B	D	NR	NR	R	65.3 (13.7)	37.9	NR	NR	NR	NR	NR	NR	NR	NR	E	3	Taiwan	I
Luengo-Fernandez	131	2013	No	UK	B	D	83	NR	C	75 (12)	51	1/6/12 months, 2/5 years	NR	60	10	NR	13	20	19	E	3	UK	I/P

Lunde	132	2012	No	Norway	U	D	NR	NR	C	68.7 (12.9)	36	Mean 12.2 (4.6)	NR	NR	NR	NR	NR	NR	NR	E, 15D	3	UK	M
Mahesh	133	2019	No	Sri Lanka	U	D	NR	NR	R/O	NR	NR	At 28-32 days	NR	NR	NR	NR	NR	NR	NR	E	3	Sri Lanka	I
Mar 1	134	2015	No	Spain	B	D	90.7	NR	H/C	72.1 (13.2)	45.2	At 3/12 months	NR	NR	NR	NR	NR	NR	50.2	E	3	NR	I
Mar 2	135	2005	No	Spain	B	D	57.8	NR	C	70.9 (SE 12.3)	38.5	At 1 year	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Mathias	136	1997	No	USA	B	D	NR	NR	C	69.2 (range 35-92)	61	Within 3 months	NR	NR	NR	NR	NR	NR	NR	H2	N/A	NR	I
McDonnell	137	2014	No	Australia	B	S	NR	NR	C	70 (10) exercise, 65 (7) no exercise	25.9	NR	NR	NR	NR	NR	NR	NR	NR	A	N/A	NR	I
Min	138	2015	No	Korea	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	Korea	M
Mittmann 1	139	2001	No	Canada	U	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	H3	N/A	NR	I/P
Mittmann 2	140	1999	No	Canada	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	H3	N/A	NR	I/P
Mulhern	141	2018	No	UK	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3, 5	UK	P
Oemrawsingh	142	2019	No	Netherlands	I	D	100	NR	C	74 (IQR 64-82)	43	At 3 months	Median 4 (IQR 2-12)	53	4	22	10	27	NR	E	5	Dutch	I/P
Olsson 1	143	2007	No	Sweden	B	D	70	46	C	50	48	Median 32 months (range 27-44)	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Olsson 2	144	2006	No	Sweden	B	D	71.2	44	R (outpatient0	51.3 (8.3)	46.2	Mean 180 days (range 22-473)	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Park	145	2013	No	Korea	B	D	NR	NR	C	69 (7)	38	At 3 months	NR	67	34	41	5	NR	2	E	3	Korean	I
Paul	146	2005	No	Australia	B	D	NR	NR	C	75.5 (13.8)	55%	At 5 years	NR	NR	NR	NR	NR	NR	NR	A	N/A	Australia	I
Peng	147	2019	No	Taiwan	B	D	80	NR	H	65.6 (IQR 57-75)	37.1	NR	NR	NR	NR	NR	NR	NR	NR	E	3	Taiwan	I
Pettersson	148	2007	No	Sweden	B	S	NR	NR	C	67 (range 43-85)	31	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Peters	149	2014	No	UK	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	UK	M
Phan 1	150	2020	No	Australia	B	D	NR	NR	C	71.2 (IQR 62-79.5) men/ 77.1 (IQR 65.7-84.4) women	44	Median 11.5 months (IQR 10.5-13.4)	NR	72.7	19.6	21.5	24.2	NR	34.7	E	3	NR	NR
Phan 2	151	2019	No	Australia	B	D	88.8	NR	C	68.6 (10) for EQ-5D/ 69 (14) for AQOL	48.3 for EQ-5D/ 49.2 for AQOL	At 1 year and 5 years	NR	60.7 (Oxford sample)	12.5	NR	11.5	NR	15.5	E, A	3	UK for EQ-5D/Australia for AQoL	I
Pickard 1	10	2005	No	Canada	I	D	NR	NR	H/C	67 (15)	48	Before discharge (95% within 2 weeks of stroke)	NR	NR	NR	NR	NR	NR	NR	E, H2, H3	3	NR	S

												and 6 months											
Pickard 2	53	2004	No	Canada	I	D	NR	NR	H/C	68.3 (14.6)	47%	At 1/3/6 months	NR	NR	NR	NR	NR	NR	NR	E, H3	3	UK	NR
Pinto	152	2011	No	Brazil	B	D	NR	NR	O	59.3 (13.3)	55.2	Mean 28 (36) months	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Price	153	2018	No	Australia	B	S	NR	34.5	C	62.3 (10.89)	48.3	Mean 3.8 years (4.1)	NR	NR	NR	NR	NR	NR	NR	A	N/A	NR	I
Ramirez-Moreno	154	2018	No	Spain	B	D	NR	NR	C/O	59.5 (8.2)	23.5	At 3 months	NR	58.7	30.4	64.8	NR	27.2	12	E	5	NR	I
Ran	155	2015	Y	China	I	D	NR	NR	NR	61.9 (12.3)	40.9	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	I
Rivero-Arias	156	2009	No	UK	B	D	NR	NR	C	72.8	53	NR	NR	NR	NR	NR	NR	NR	NR	E	3	UK	I
Saarni	157	2006	No	Finland	B	S	NR	NR	C	70	48	NR	NR	NR	NR	NR	NR	NR	NR	E, 15D	3	UK	I
Sallinen	158	2018	No	Finland	H	D	0	NR	C	70.5 (IQR 62-78)	48	At 3 months	Median 5	NR	13.7	NR	10.1	18.8	24.8	E, 15D	5	Crosswalk	I/M
Sanchez-Iriso	159	2017	No	Spain	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	5	Spain	NR
Sand	160	2015	No	Norway	I	D	NR	NR	C	71.8 (14.3) vision problems/ 66.5 (12.4) noo vision problems	37.3	Mean 372 days (range 185-757)	Mean 5.9 (6.4) vision problems, 3.8 (4.2) no vision problems	51.7	9.6	58.8	11.6	13.5	NR	E, 15D	3	NR	M
Sasaki	161	2018	No	Japan	B	D	NR	NR	O	69.5 (12)	27.3	Mean 6.3 months (6.2)	mean 1.7 (1.5)	90.1	22.7	NR	NR	NR	NR	E	3	Japan	S
Slaughter	162	2019	No	USA	H	D	NR	NR	C	63.9 (14.9)	38	At 3 months	Median 11 (IQR 3-22)	77.6	28	15.1	10.8	26.2	11.1	E	5	NR	P
Sturm 1	163	2004	No	Australia	B	D	85	NR	C	71 (95% CI 69-73)	49	Mean 737 days (range 646-898)	NR	NR	NR	NR	NR	NR	NR	A	N/A	NR	I
Sturm 2	164	2002	No	Australia	B	D	NR	NR	C	72 (range 28-89)	55	At 3 months	NR	NR	NR	NR	NR	NR	NR	A	N/A	NR	I
Szocs	165	2020	No	Budapest	I	D	NR	NR	C	68.5 (12.9)	44	At 3 months	Mean 7.88 (6.37)	NR	NR	NR	NR	NR	NR	E, 15D	5	NR	I/P
Teoh	166	2009	No	Australia	B	D	76.5	NR	C	67.5 (14.3)	32%	Mean 11.7 months (4.9)	NR	NR	NR	NR	NR	NR	NR	A	N/A	NR	NR
Tran	167	2015	No	Vietnam	B	D	72.7	NR	C	60.9 (12)	49.1	At 3 months	Mean 7.7 (0.6)	NR	NR	NR	NR	NR	NR	E	3	South Korea	S
Vahlberg	168	2013	No	Sweden	B	D	88	NR	C	74 (5.2)	29.2	1-3 years after	NR	64	16	15	12	21	22	E	3	NR	I
van Eeden	169	2015	No	Netherlands	B	D	92.9	38.2	H/C	66.8 (12.27)	35.2	At 2/6/12 months	Mean 2.6 (2.96)	NR	NR	NR	NR	NR	NR	E	3	Netherlands	I/O/M
Visser	170	2015	No	Cyprus	B	D	73.5	40.3	R	53.06 (10.19)	47	83.7% <1 year post stroke	NR	NR	NR	NR	NR	NR	NR	E	5	NR	NR
Wartenberg	171	2020	No	Germany	I	D	NR	NR	C	62.1 (12.6)	48.8	At 12 months	Median 19 (IQR 12-36)	74.4	27.9	NR	11.6	16.3	0	E	3	NR	M
White	172	2016	No	Australia	B	D	NR	NR	R/C	75 (12)	55	NR	NR	NR	NR	NR	NR	NR	NR	A	N/A	NR	I
Wu 1	173	2014	No	UK	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M

Wu 2	174	2015	No	UK	B	S	NR	NR	C	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	M
Xie	175	2006	No	USA	B	S	NR	NR	C	NR	56.1	NR	NR	67.9	27.4	23.6	33.9	NR	NR	E	3	USA	I
Yan	176	2015	Y	China	B	D	NR	NR	H	61.8 (12.67)	61	NR	NR	NR	NR	NR	NR	NR	NR	E	3	China	I
Yang	177	2017	No	Korea	B	D	64.7	NR	R (outpatient)	65.11 (2.39)	41.1	NR	NR	NR	NR	NR	NR	NR	NR	E	3	NR	NR
Yeoh 1	178	2018	No	Singapore	B	D	89.7	NR	H/C	62.2 (10.6)	34	At 3/12 months	NR	NR	NR	NR	NR	NR	NR	E	3	Singapore	I
Yeoh 2	179	2019	No	Singapore	B	D	90.9	NR	H/C	61.8 (10.3)	32.5	At 3/12 months	Mean 3.8 (3.5)	71.1	37.1	NR	NR	NR	NR	E	3	Singapore	I

Y = Yes; I=ischemic only; H=hemorrhagic only; B=Both ischemic and hemorrhagic; U=Unknown stroke type; S=Self-report of stroke; D=medical diagnosis of stroke; C=Community based; R=Rehabilitation facility; H=hospital; O=outpatient clinic; E=EQ-5D; A=AQoL; H2= HUI2; H3=HUI3; M=mail; P=phone; I=in-person interview; NR = not reported; N/A = Not applicable.
Numbers in brackets denote standard deviation unless specifically indicated otherwise.

Table S5. Risk of bias assessment criteria

Item	Description
Sample size	Very small <50 Small 50-99 Medium size 100-999 Large ≥1000
Respondent selection and recruitment*	Does this result in a population comparable to that being modelled? Is this sample broadly representative of stroke patients or skewed towards one subgroup? Was the selection of patients consecutive or population-based, or is there evidence of enrollment bias?
Inclusion/exclusion criteria*	Does this study exclude important groups, ie. very elderly, young adult, severe or mild strokes, or include only a narrow or select group of patients (ie. hemispheric infarcts).
Response rates to instrument used	Are response rates reported and if so, are the rates likely to be a threat to validity?
Loss to follow-up	How large is the loss to follow-up and are these reasons given? Are these likely to threaten the validity of the estimates?
Missing data	What are the levels of missing data and how are they dealt with? Could they threaten the validity of the estimates?
Proxy responses	Is the presence, proportion, and method of proxy responses reported?

*A high risk of bias does not necessarily imply methodological or quality concerns, as the objective of the study may have been to report utility in a specific sub-group (ie. patients with hemisphere stroke), but this value would be less representative of stroke patients as a whole.

Table S6. Baseline characteristics across all studies weighted by study size, among studies where variables were reported

Variable	Pooled mean (SD) weighted by study size	Number of studies for pooled mean
Mean Age	68.1 (5.7)	90
Percent Female	44.2 (6.2)	91
Mean Follow-up time	13.0 months (15.7)	69
Mean National Institutes of Health Stroke Scale score	6.5 (4.1)	19
Percent Ischemic stroke	85.5 (8.3)	50
Percent Left hemisphere	43.9 (5.2)	14
Percent Hypertension	68.8 (10.3)	27
Percent Diabetes	22.6 (9.2)	29
Percent Smoking	24.5 (9.3)	20
Percent Coronary artery disease	19.2 (8.9)	19
Percent Prior stroke	20.7 (5.9)	17
Percent Atrial fibrillation	24.1 (12.0)	17

Table S7. Baseline characteristics across studies reporting EQ-5D-3L weighted by study size, among studies where variables were reported

Variable	Pooled mean (SD) weighted by study size	Number of studies for pooled mean
Mean Age	68.2 (5.7)	65
Percent Female	44.3 (5.8)	64
Mean Follow-up time	10.9 months (11.1)	49
Mean National Institutes of Health Stroke Scale score	6.2 (4.3)	14
Percent Ischemic stroke	85.2 (8.3)	40
Percent Left hemisphere	43.6 (5.8)	10
Percent Hypertension	69.0 (10.3)	23
Percent Diabetes	22.7 (9.4)	23
Percent Smoking	24.7 (8.9)	17
Percent Coronary artery disease	19.6 (9.1)	14
Percent Prior stroke	20.5 (5.8)	12
Percent Atrial fibrillation	25.0 (12.1)	12

Table S8. Risk of bias assessment

Author	Sample size	Respondent selection/recruitment	Inclusion/Exclusion	Response rates	Loss to FU	Missing data	Proxy responses	Death
Adey-Wakeling	100-999	Low risk	Low risk	Low risk	Low risk	Medium risk*	?	Low risk
Appau	100-999	Medium risk	Low risk	?	?	?	?	?
Arrospide	100-999	Medium risk	Low risk	?	N/A	Low risk	?	N/A
Arwert	<50	Medium risk	Medium risk	High risk	?	?	?	N/A
Barton	50-99	Medium risk	Medium risk	Medium risk	N/A	Medium risk	?	N/A
Broussy	100-999	Medium risk	Low risk	Low risk	Medium risk	Low risk	Low risk	Low risk
Burton	50-99	Medium risk	Medium risk	High risk	N/A	Medium risk	Low risk	N/A
Bushnell	1000+	Low risk	Low risk	Low risk	Medium risk	Medium risk	Medium risk	Low risk
Cadilhac 1	100-999	Low risk	Low risk	Medium risk	Medium risk	Low risk	?	N/A
Cadilhac 2	100-999	Low risk	Low risk	?	?	?	?	N/A
Cao	100-999	Low risk	Medium risk	?	N/A	?	?	N/A
Chang	1000+	Low risk	Low risk	Low risk	Medium risk	Medium risk	?	N/A
Chen 1	50-99	Medium risk	Medium risk	Low risk	N/A	Low risk	?	N/A
Chen 2	100-999	Medium risk	Medium risk	Low risk	Medium risk	?	?	Low risk
Cheung	100-999	Medium risk	Low risk	High risk	Medium risk	?	?	N/A
Cramm	100-999	High risk	High risk	High risk	N/A	?	Low risk	N/A
Cup	<50	Medium risk	Medium risk	Low risk	N/A	?	?	N/A
Darlington 1	50-99	Medium risk	Medium risk	Medium risk	Medium risk	Medium risk	?	Low risk
Darlington 2	50-99	Medium risk	Medium risk	Medium risk	Medium risk	Medium risk	?	Low risk
de Graaf	100-999	Low risk	Low risk	?	N/A	?	?	N/A
Deb-Chatterji	100-999	High risk	High risk	Low risk	Low risk	Low risk	?	Low risk
Dewilde	100-999	Medium risk	Low risk	?	N/A	?	?	N/A
Dorman 1	100-999	Medium risk	Low risk	Low risk	N/A	?	Low risk	N/A

Dorman 2	100-999	Medium risk	Low risk	Low risk	N/A	?	Low risk	Medium risk
Du	100-999	Medium risk	Low risk	Low risk	N/A	?	?	N/A
Edwards	100-999	Medium risk	Low risk	?	N/A	?	?	N/A
Espuela	100-999	Medium risk	Medium risk	Medium risk	Medium risk	?	?	Medium risk
Fischer	100-999	Medium risk	High risk	Low risk	N/A	Low risk	Medium risk	N/A
Ghatnekar	100-999	Low risk	Low risk	?	N/A	?	Low risk	N/A
Golicki 1	100-999	Medium risk	Low risk	Low risk	Low risk	Low risk	Medium risk	N/A
Golicki 2	100-999	Medium risk	Medium risk	Low risk	N/A	Low risk	Medium risk	N/A
Graessel	100-999	High risk	Medium risk	Low risk	N/A	?	?	Low risk
Grasel	100-999	High risk	Medium risk	Low risk	N/A	?	?	Medium risk
Groeneveld	100-999	Medium risk	Medium risk	High risk	N/A	Medium risk	?	Low risk
Groeneveld	100-999	Medium risk	Medium risk	Medium risk	Low risk	?	?	Low risk
Guo	50-99	Medium risk	Low risk	Medium risk	Medium risk	?	Low risk	Low risk
Haacke	50-99	Medium risk	Low risk	Medium risk	Medium risk	?	Low risk	Medium risk
Hansson	100-999	Medium risk	Medium risk	Medium risk	Medium risk	?	Medium risk	Low risk
Hokstad	50-99	Medium risk	Low risk	Medium risk	Medium risk	Medium risk	?	Medium risk
Jeon	<50	High risk	Medium risk	Medium risk	N/A	?	?	N/A
Katona	100-999	High risk	Medium risk	Low risk	Medium risk	?	?	Medium risk
Katzan	1000+	Medium risk	Medium risk	Low risk	N/A	?	?	N/A
Kelly	<50	High risk	High risk	High risk	High risk	High risk	High risk	High risk
Khiaocharoen	100-999	Medium risk	Medium risk	Medium risk	N/A	?	?	N/A
Kil	<50	Medium risk	High risk	?	N/A	Medium risk	?	N/A
Kim 1	100-999	Medium risk	Low risk	?	N/A	?	?	N/A
Kim 2	100-999	Medium risk	Low risk	?	N/A	Low risk	?	N/A
Kuo	100-999	Medium risk	Low risk	?	N/A	?	Low risk	N/A
Kuroda 1	100-999	Medium risk	Low risk	Medium risk	Medium risk	?	Medium risk	Medium risk
Kuroda 2	100-999	Medium risk	Low risk	Medium risk	Medium risk	?	?	Medium risk
Kuwano	100-999	High risk	Medium risk	Medium risk	Medium risk	?	?	Medium risk

Kwon	100-999	Medium risk	Low risk	?	N/A	?	?	N/A
Labberton	100-999	Medium risk	Low risk	Medium risk	Medium risk	Low risk	Low risk	Medium risk
Lannin	1000+	Low risk	Low risk	Medium risk	Medium risk	?	?	Low risk
Leach	100-999	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	High risk
Lee	100-999	Medium risk	Low risk	Low risk	High risk	?	?	Medium risk
Leeds	50-99	High risk	Medium risk	Medium risk	Medium risk	Medium risk	?	Medium risk
Lindgren	100-999	Low risk	High risk	Medium risk	N/A	?	Medium risk	N/A
Lopez-Bastida	100-999	Low risk	Low risk	Medium risk	N/A	Medium risk	?	N/A
Lu	100-999	Medium risk	High risk	Low risk	Medium risk	Medium risk	?	?
Luengo-Fernandez	100-999	Low risk	Low risk	Medium risk	Medium risk	Medium risk*	?	N/A
Lunde	100-999	Medium risk	Low risk	Medium risk	N/A	Low risk	?	N/A
Mahesh	100-999	Medium risk	Medium risk	?	N/A	?	Low risk	N/A
Mar 1	100-999	Low risk	Low risk	Low risk	Low risk	?	?	Low risk
Mar 2	100-999	Medium risk	Low risk	High risk	Medium risk	?	?	Medium risk
Mathias	<50	Medium risk	High risk	Low risk	N/A	Low risk	Low risk	N/A
McDonnell	<50	High risk	High risk	Low risk	N/A	?	?	N/A
Min	1000+	Medium risk	Low risk	Low risk	N/A	Low risk	?	N/A
Mittmann 1	50-99	Medium risk	Medium risk	?	N/A	High risk	?	N/A
Mittmann 2	50-99	Medium risk	Medium risk	?	N/A	?	?	N/A
Mulhern	50-99	Medium risk	Medium risk	?	N/A	?	?	N/A
Oemrawsingh	1000+	Medium risk	Low risk	?	High risk	High risk	?	?
Olsson 1	50-99	High risk	High risk	?	N/A	?	?	Low risk
Olsson 2	50-99	High risk	High risk	?	N/A	?	?	N/A
Park	100-999	High risk	High risk	Medium risk	N/A	?	?	N/A
Paul	100-999	Low risk	Low risk	Medium risk	Medium risk	?	Medium risk	High risk
Peng	1000+	Medium risk	High risk	?	?	?	?	?
Peterrsson	<50	High risk	High risk	Low risk	Medium risk	?	?	N/A

Peters	50-99	Medium risk	Low risk	High risk	N/A	?	?	N/A
Phan 1	1000+	Low risk	Low risk	Medium risk	Medium risk	Medium risk*	Medium risk	Medium risk
Phan 2	1000+	Low risk	Low risk	Medium risk	Medium risk	Low risk*	?	Medium risk
Pickard 1	50-99	Medium risk	Medium risk	Medium risk	Medium risk	Low risk	Low risk	Low risk
Pickard 2	100-999	Medium risk	Medium risk	Medium risk	Medium risk	?	Low risk	Medium risk
Pinto	50-99	Medium risk	High risk	?	N/A	?	?	N/A
Price	<50	Medium risk	Medium risk	?	N/A	?	?	N/A
Ramirez-Moreno	50-99	High risk	High risk	Medium risk	?	?	?	?
Ran	100-999	Medium risk	Medium risk	Low risk	?	?	?	?
Rivero-Arias	1000+	Low risk	Low risk	Medium risk	Medium risk	Medium risk	?	Low risk
Saarni	100-999	Medium risk	Low risk	Low risk	N/A	?	?	N/A
Sallinen	100-999	Medium risk	Medium risk	Medium risk	Medium risk	Low risk	Medium risk	Medium risk
Sanchez-Iriso	100-999	Medium risk	Low risk	?	N/A	?	?	N/A
Sand	100-999	Medium risk	Low risk	Medium risk	Medium risk	?	?	?
Sasaki	<50	High risk	High risk	High risk	N/A	?	?	N/A
Slaughter	100-999	Medium risk	Medium risk	Medium risk	Medium risk	?	Medium risk	Medium risk
Sturm 1	100-999	Low risk	Low risk	Medium risk	Medium risk	?	Medium risk	Medium risk
Sturm 2	100-999	Low risk	Low risk	Medium risk	Medium risk	?	Medium risk	Medium risk
Szocs	100-999	Medium risk	Low risk	Medium risk	Medium risk	Medium risk*	?	Medium risk
Teoh	100-999	Medium risk	Medium risk	High risk	N/A	?	?	Low risk
Tran	100-999	Medium risk	Medium risk	Medium risk	?	?	Low risk	?
Vahlberg	100-999	Medium risk	High risk	Medium risk	N/A	?	?	N/A
van Eeden	100-999	Medium risk	Low risk	Low risk	N/A	Medium risk*	?	Low risk
Visser	100-999	High risk	High risk	Medium risk	N/A	?	?	N/A
Wartenberg	<50	High risk	High risk	?	N/A	?	?	N/A

White	100-999	Medium risk	Medium risk	Low risk	Medium risk	?	?	Medium risk
Wu 1	100-999	Medium risk	Low risk	High risk	N/A	?	?	N/A
Wu 2	100-999	Medium risk	Low risk	High risk	N/A	?	?	N/A
Xie	1000+	Medium risk	Low risk	?	N/A	Low risk	Medium risk	N/A
Yan	100-999	Medium risk	Medium risk	?	Medium risk	?	?	Medium risk
Yang	<50	High risk	High risk	?	N/A	?	?	N/A
Yeoh 1	100-999	Medium risk	Low risk	?	Medium risk	Medium risk*	?	Low risk
Yeoh 2	100-999	Medium risk	Low risk	Medium risk	Medium risk	Low risk	Low risk	Low risk

? = information not provided; N/A = follow-up not applicable, typically due to cross-sectional design
***8 studies used imputation for missing data: Adey-Wakeling, Luengo-Fernandez, Phan 1, Phan 2, Pickard 1, Szocs, van Eeden, Yeoh 1**

Table S9. Summary statistics of risk of bias assessment

Category	Percent with high risk	Percent with medium risk	Percent with low risk	Percent with missing information
Sample size	10.8	79.3	9.91	0
Respondent selection	17.1	64.9	18.0	0
Inclusion/Exclusion	18.0	33.3	48.7	0
Response rates	10.8	36.0	27.0	26.1
Loss to follow-up*	2.7	36.0	5.4	6.3
Missing data	2.7	17.1	17.1	63.1
Proxy responses	0.90	12.6	15.3	71.2

***Question not applicable in 49.6% due to cross-sectional nature of study**

Table S10. Pooled EQ-5D-3L values for different time ranges and time points

Time category	Pooled health utility value (95% CI)	Number of studies	Patient N
Ranges			
Prior to acute care discharge (hospital or in-patient rehabilitation)	0.45 (0.33-0.58)	16	4764
Prior to hospital discharge	0.39 (0.23-0.54)	10	3517
Prior to in-patient rehabilitation discharge	0.57 (0.47-0.67)	6	1247
≥ 3 months	0.65 (0.63-0.67)	73	52614
3 to <12 months	0.66 (0.63-0.68)	54	48020
12 months and over	0.66 (0.62-0.69)	31	7610
Specific time points			
3 months	0.65 (0.61-0.70)	20	11624
12 months	0.65 (0.59- 0.71)	17	4917
5 years	0.70 (0.64-0.76)	6	2455

Table S11. Sensitivity analyses

Utility metric and sensitivity analysis	Pooled health utility value	Number of studies
EQ-5D-3L		
Exclude studies with similar/overlapping cohorts	0.66 (0.64-0.68)	65
Exclude studies that assigned subjects who died to a utility of 0	0.66 (0.64-0.68)	70
Include only studies assigning subjects who died to a utility of 0	0.50 (0.33-0.67)	3
Exclude studies with >1 high risk of bias category	0.68 (0.65-0.70)	50
EQ-5D-5L		
Exclude studies with similar/overlapping cohorts	0.68 (0.60-0.76)	9
Exclude studies that assigned subjects who died to a utility of 0	N/A	
Include only studies assigning subjects who died to a utility of 0	N/A	
Exclude studies with >1 high risk of bias category	0.66 (0.59-0.74)	9
AQOL		
Exclude studies with similar/overlapping cohorts	0.54 (0.41-0.66)	7
Exclude studies that assigned subjects who died to a utility of 0	N/A	
Include only studies assigning subjects who died to a utility of 0	N/A	
Exclude studies with >1 high risk of bias category	0.49 (0.44-0.54)	5

*Sensitivity analyses not performed for certain categories above and for HUI2, HUI3, and 15D due to small number of studies

Table S12. Comparison of summary estimates using random effects and fixed effect meta-analysis

	Random effects	**Fixed effect**
Main (EQ-5D-3L)	0.65 (0.63-0.67)	0.73 (0.73-0.73)
Ranges (EQ-5D-3L)		
Prior to acute care discharge (hospital or in-patient rehabilitation)	0.45 (0.33-0.58)	0.40 (0.39-0.40)
Prior to hospital discharge	0.39 (0.23-0.54)	0.28 (0.27-0.29)
Prior to in-patient rehabilitation discharge	0.57 (0.47-0.67)	0.59 (0.58-0.61)
≥3 months	0.65 (0.63-0.67)	0.73 (0.73-0.73)
3 to <12 months	0.66 (0.63-0.68)	0.73 (0.73-0.73)
12 months and over	0.66 (0.62-0.69)	0.71 (0.70-0.71)
Specific time points (EQ-5D-3L)		
3 months	0.65 (0.61-0.70)	0.76 (0.76-0.77)
12 months	0.65 (0.59- 0.71)	0.78 (0.77-0.78)
5 years	0.70 (0.64-0.76)	0.72 (0.71-0.74)
Other metrics		
EQ-5D-5L	0.68 (0.61-0.76)	0.72 (0.71-0.73)
AQOL	0.51 (0.42-0.61)	0.40 (0.39-0.41)
HUI2	0.65 (0.62-0.68)	0.65 (0.62-0.68)
HUI3	0.64 (0.54-0.73)	0.71 (0.69-0.72)
15D	0.81 (0.78-0.84)	0.82 (0.82-0.83)
SF-6D	0.70 (0.63-0.78)	0.73 (0.72-0.75)

Figure S1. PRISMA flowchart




Figure S2. General characteristics of the included papers, showing frequency of studies with each health utility instrument (A), frequency of studies by country (B), frequency of studies by year from 1995 to October 2020 (C), and distribution of EQ-5D-3L values from 3 months onwards (D).




Figure S3. Random effects meta-analysis for EQ-5D-3L utility values




Figure S4. Fixed effect meta-analysis for EQ-5D-3L utility values




Figure S5. Random effects meta-analysis for EQ-5D-5L




Figure S6. Random effects meta-analysis for AQoL




Figure S7. Random effects meta-analysis for HUI2




Figure S8. Random effects meta-analysis for HUI3




Figure S9. Random effects meta-analysis for 15D




Figure S10. Random effects meta-analysis for SF-6D




Figure S11. Sex-stratified random effects meta-analysis for EQ-5D-3L




Figure S12. Age-stratified random effects meta-analysis for EQ-5D-3L




Figure S13. Stroke type-stratified random effects meta-analysis for EQ-5D-3L




Figure S14. Random effects meta-analysis for EQ-5D-3L stratified from time from stroke





Figure S15. Meta-regression analyses of mean EQ-5D-3L utility score across percentage females, mean/median age, and publication date. Higher percentage female in the study is associated with lower health utility.



Supplemental References – studies included in systematic review

9. Golicki D, Niewada M, Buczek J, Karlińska A, Kobayashi A, Janssen MF, Pickard AS. Validity of EQ-5D-5L in stroke. Qual Life Res. 2015;24:845–850 (Golicki 2)

10. Pickard A, Johnson JA, Feeny DH. Responsiveness of generic health-related quality of life measures in stroke. Qual Life Res. 2005;14:207–219

11. Bushnell CD, Reeves MJ, Zhao X, Pan W, Prvu-Bettger J, Zimmer L, Olson D, Peterson E. Sex differences in quality of life after ischemic stroke. Neurology. 2014;82:922–931

47. Lannin NA, Anderson CS, Kim J, Kilkenny M, Bernhardt J, Levi C, Dewey HM, Bladin C, Hand P, Castley H, Hill K, Faux S, Grimley R, Grabsch B, Middleton S, Donnan G, Cadilhac DA. Treatment and Outcomes of Working Aged Adults with Stroke: Results from a National Prospective Registry. Neuroepidemiology. 2017;49:113–120

53. Pickard AS, Johnson JA, Feeny DH, Shuaib A, Carriere KC, Nasser AM. Agreement Between Patient and Proxy Assessments of Health-Related Quality of Life After Stroke Using the EQ-5D and Health Utilities Index. Stroke. 2004;35:607–612 (Pickard 2)

70. Mulhern B, Feng Y, Shah K, Janssen MF, Herdman M, van Hout B, Devlin N. Comparing the UK EQ-5D-3L and English EQ-5D-5L Value Sets. Pharmacoeconomics. 2018;36:699–713

73. Adey-Wakeling Z, Liu E, Crotty M, Leyden J, Kleinig T, Anderson CS, Newbury J. Hemiplegic Shoulder Pain Reduces Quality of Life After Acute Stroke: A Prospective Population-Based Study. American Journal of Physical Medicine & Rehabilitation. 2016;95:758–763

74. Appau A, Lencucha R, Finch L, Mayo N. Further validation of the Preference-Based Stroke Index three months after stroke. Clin Rehabil. 2019;33:1214–1220

75. Arrospide A, Machón M, Ramos-Goñi JM, Ibarrondo O, Mar J. Inequalities in health-related quality of life according to age, gender, educational level, social class, body mass index and chronic diseases using the Spanish value set for Euroquol 5D-5L questionnaire. Health Qual Life Outcomes. 2019;17:69

76. Arwert HJ, Schults M, Meesters JJL, Wolterbeek R, Boiten J, Vliet Vlieland T. Return to Work 2–5 Years After Stroke: A Cross Sectional Study in a Hospital-Based Population. J Occup Rehabil. 2017;27:239–246

77. Barton GR, Sach TH, Doherty M, Avery AJ, Jenkinson C, Muir KR. An assessment of the discriminative ability of the EQ-5Dindex, SF-6D, and EQ VAS, using sociodemographic factors and clinical conditions. Eur J Health Econ. 2008;9:237–249

78. Broussy S, Saillour-Glenisson F, García-Lorenzo B, Rouanet F, Lesaine E, Maugeais M, Aly F, Glize B, Salamon R, Sibon I. Sequelae and Quality of Life in Patients Living at Home 1 Year After a Stroke Managed in Stroke Units. Front. Neurol 2019;10:907

79. Burton CR, Fargher E, Plumpton C, Roberts GW, Owen H, Roberts E. Investigating preferences for support with life after stroke: a discrete choice experiment. BMC Health Serv Res. 2014;14:63

80. Cadilhac DA, Dewey HM, Vos T, Carter R, Thrift AG. The health loss from ischemic stroke and intracerebral hemorrhage: evidence from the North East Melbourne Stroke Incidence Study (NEMESIS). 2010;8.

81. Cadilhac DA, Kilkenny MF, Lannin NA, Dewey HM, Levi CR, Hill K, Grabsch B, Grimley R, Blacker D, Thrift AG, et al. Outcomes for Patients With In-Hospital Stroke: A Multicenter Study From the Australian Stroke Clinical Registry (AuSCR). Journal of Stroke and Cerebrovascular Diseases. 2019;28:1302–1310 (Cadilhac 2)

82. Cao Y, Tang X, Yang L, Li N, Wu Y, Fan W, Liu J, Yu L, Xu H, Liu W, et al. [Influence of chronic diseases on health related quality of life in middle-aged and elderly people from rural communities: application of EQ-5D scale on a Health Survey in Fangshan, Beijing]. Zhonghua Liu Xing Bing Xue Za Zhi. 2012;33:17–22

83. Chang WH, Sohn MK, Lee J, Kim DY, Lee S-G, Shin Y-I, Oh G-J, Lee Y-S, Joo MC, Han EY, et al. Predictors of functional level and quality of life at 6 months after a first-ever stroke: the KOSCO study. J Neurol. 2016;263:1166–1177

84. Chen P, Lin K-C, Liing R-J, Wu C-Y, Chen C-L, Chang K-C. Validity, responsiveness, and minimal clinically important difference of EQ-5D-5L in stroke patients undergoing rehabilitation. Qual Life Res. 2016;25:1585–1596

85. Chen C-J, Ding D, Buell TJ, Testai FD, Koch S, Woo D, Worrall BB, for the ERICH Investigators. Restarting antiplatelet therapy after spontaneous intracerebral hemorrhage: Functional outcomes. Neurology. 2018;91:e26–e36 (Chen 2)

86. Cheung YB, Tan HX, Luo N, Wee HL, Koh GCH. Mapping the Shah-modified Barthel Index to the Health Utility Index Mark III by the Mean Rank Method. Qual Life Res. 2019;28:3177–3185

87. Cramm JM, Strating MMH, Nieboer AP. Satisfaction with care as a quality-of-life predictor for stroke patients and their caregivers. Qual Life Res. 2012;21:1719–1725

88. Cup EHC, Scholte op Reimer WJM, Thijssen MCE, van Kuyk-Minis MAH. Reliability and validity of the Canadian Occupat189ional Performance Measure in stroke patients. Clin Rehabil. 2003;17:402–409

89. Darlington A, Dippel D, Ribbers G, van Balen R, Passchier J, Busschbach J. A prospective study on coping strategies and quality of life in patients after stroke, assessing prognostic relationships and estimates of cost-effectiveness. J Rehabil Med. 2009;41:237–241

90. Darlington A-SE, Dippel DWJ, Ribbers GM, van Balen R, Passchier J, Busschbach JJV. Coping Strategies as Determinants of Quality of Life in Stroke Patients: A Longitudinal Study. Cerebrovasc Dis. 2007;23:401–407 (Darlington 2)

91. de Graaf J, Kuijpers M, Visser-Meily J, Kappelle L, Post M. Validity of an enhanced EQ-5D-5L measure with an added cognitive dimension in patients with stroke. Clin Rehabil. 2020;34:545–550

92. Deb-Chatterji M, Konnopka A, Flottmann F, Leischner H, Fiehler J, Gerloff C, Thomalla G. Patient-reported, health-related, quality of life after stroke thrombectomy in clinical practice. Neurology. 2020;95:e1724–e1732

93. Dewilde S, Annemans L, Lloyd A, Peeters A, Hemelsoet D, Vandermeeren Y, Desfontaines P, Brouns R, Vanhooren G, Cras P, et al. The combined impact of dependency on caregivers, disability, and coping strategy on quality of life after ischemic stroke. Health Qual Life Outcomes. 2019;17:31

94. Dorman P. Are the modified “simple questions” a valid and reliable measure of health related quality of life after stroke? Journal of Neurology, Neurosurgery & Psychiatry. 2000;69:487–493.

95. Dorman PJ, Waddell F, Slattery J, Dennis M, Sandercock P. Are Proxy Assessments of Health Status After Stroke With the EuroQol Questionnaire Feasible, Accurate, and Unbiased? Stroke. 1997;28:1883–1887. (Dorman 2)

96. Du X-D, Zhu P, Li M-E, Wang J, Meng H-D, Zhu C-R. [Health Utility of Patients with Stroke Measured by EQ-5D and SF-6D]. Sichuan Da Xue Xue Bao Yi Xue Ban. 2018;49:252–257

97. Edwards JD, Koehoorn M, Boyd LA, Levy AR. Is Health-Related Quality of Life Improving After Stroke?: A Comparison of Health Utilities Indices Among Canadians With Stroke Between 1996 and 2005. Stroke. 2010;41:996–1000

98. López Espuela F, Portilla Cuenca JC, Leno Díaz C, Párraga Sánchez JM, Gamez-Leyva G, Casado Naranjo I. Sex differences in long-term quality of life after stroke: Influence of mood and functional status. Neurologia (Engl Ed). 2020;35:470–478

99. Fischer U, Anca D, Arnold M, Nedeltchev K, Kappeler L, Ballinari P, Schroth G, Mattle HP. Quality of Life in Stroke Survivors after Local Intra-Arterial Thrombolysis. Cerebrovasc Dis. 2008;25:438–444

100. Ghatnekar O, Eriksson M, Glader E-L. Mapping health outcome measures from a stroke registry to eq-5d weights. Health Qual Life Outcomes. 2013;11:34

101. Golicki D, Niewada M, Karlińska A, Buczek J, Kobayashi A, Janssen MF, Pickard AS. Comparing responsiveness of the EQ-5D-5L, EQ-5D-3L and EQ VAS in stroke patients. Qual Life Res. 2015;24:1555–1563

102. Graessel E, Schmidt R, Schupp W. Stroke patients after neurological inpatient rehabilitation: a prospective study to determine whether functional status or health-related quality of life predict living at home 2.5 years after discharge. International Journal of Rehabilitation Research 2014;37:212–219

103. Gräßel E, Schupp W, Schmidt R. Schlaganfallpatienten nach stationärer neurologischer Rehabilitation: Eine prospektive Studie zur Ermittlung von Prädiktoren für das Überleben zuhause bis 5 Jahre nach Entlassung. Rehabilitation. 2019;58:296–303

104. Groeneveld IF, Goossens PH, van Braak I, van der Pas S, Meesters JJL, Rambaran Mishre RD, Arwert HJ, Vliet Vlieland TPM. Patients' outcome expectations and their fulfilment in multidisciplinary stroke rehabilitation. Annals of Physical andRehabilitation Medicine. 2019;62:21–27

105. Groeneveld IF, Goossens PH, van Meijeren-Pont W, Arwert HJ, Meesters JJL, Rambaran Mishre AD, Van Vree F, Vliet Vlieland TPM. Value-Based Stroke Rehabilitation: Feasibility and Results of Patient-Reported Outcome Measures in the First Year After Stroke. Journal of Stroke and Cerebrovascular Diseases. 2019;28:499–512 (Groeneveld 2)

106. Guo YE, Togher L, Power E, Heard R, Luo N, Yap P, Koh GCH. Sensitivity to change and responsiveness of the Stroke and Aphasia Quality-of-Life Scale (SAQOL) in a Singapore stroke population. Aphasiology. 2017;31:427–446

107. Haacke C, Althaus A, Spottke A, Siebert U, Back T, Dodel R. Long-Term Outcome After Stroke: Evaluating Health-Related Quality of Life Using Utility Measurements. Stroke. 2006;37:193–198

108. Hansson EE, Beckman A, Wihlborg A, Persson S, Troein M. Satisfaction with rehabilitation in relation to self-perceived quality of life and function among patients with stroke - a 12 month follow-up: Rehabilitation in Malmö. Scandinavian Journal of Caring Sciences. 2013;27:373–379

109. Hokstad A, Indredavik B, Bernhardt J, Langhammer B, Gunnes M, Lundemo C, Bovim M, Askim T. Upright activity within the first week after stroke is associated with better functional outcome and health-related quality of life: A Norwegian multi-site study. J Rehabil Med. 2016;48:280–286

110. Jeon H, Sohn MK, Jeon M, Jee S. Clinical Characteristics of Sleep-Disordered Breathing in Subacute Phase of Stroke. Ann Rehabil Med. 2017;41:556

111. Katona M, Schmidt R, Schupp W, Graessel E. Predictors of health-related quality of life in stroke patients after neurological inpatient rehabilitation: a prospective study. Health Qual Life Outcomes. 2015;13:58

112. Katzan IL, Thompson NR, Lapin B, Uchino K. Added Value of Patient-Reported Outcome Measures in Stroke Clinical Practice. JAHA [Internet]. 2017 [cited 2021 Feb 15];6 Available from: https://wwwahajournalsorg/doi/101161/JAHA1165356

113. Kelly ML, Rosenbaum BP, Kshettry VR, Weil RJ. Comparing clinician- and patient-reported outcome measures after hemicraniectomy for ischemic stroke. Clinical Neurology and Neurosurgery. 2014;126:24–29

114. Khiaocharoen O, Pannarunothai S, Riewpaiboon W, Ingsrisawang L, Teerawattananon Y. Economic Evaluation of Rehabilitation Services for Inpatients with Stroke in Thailand: A Prospective Cohort Study. Value Health Reg Issues. 2012;1:29–35

115. Kil S-R, Lee S-I, Yun S-C, An H-M, Jo M-W. The Decline of Health-Related Quality of Life Associated with Some Diseases in Korean Adults. J Prev Med Public Health. 2008;41:434

116. Kim Y, Moon H. Association between quality of life and sleep time among community-dwelling stroke survivors: Findings from a nationally representative survey. Geriatr. Gerontol Int 2019;19:1226–1230

117. Kim Y, Kim M, Park H-S, Cho I-H, Paik JK. Association of the Anxiety/Depression with Nutrition Intake in Stroke Patients. Clin Nutr Res. 2018;7:11 (Kim 2)

118. Kuo L-M, Tsai W-C, Chiu M-J, Tang L-Y, Lee H-J, Shyu Y-IL. Cognitive dysfunction predicts worse health-related quality of life for older stroke survivors: a nationwide population-based survey in Taiwan. Aging & Mental Health. 2019;23:305–310

119. Kuroda A, Kanda T. [Correlation between QOL utility score and VAS score of EuroQol in stroke patients]. Nihon Ronen Igakkai Zasshi. 2007;44:264–266

120. Kuroda A, Kanda T, Asai N. [Health-related quality of life assessed by EuroQol in caregivers of home care stroke patients]. Nihon Ronen Igakkai Zasshi. 2003;40:381–389 (Kuroda 2)

121. Kuwano M, Kanda T, Shimizu K, Asai N, Kuwano M. [Health-related quality of life assessed by EuroQol in home care patients with stroke]. Nihon Ronen Igakkai Zasshi. 2001;38:831–833

122. Kwon S, Park J-H, Kim W-S, Han K, Lee Y, Paik N-J. Health-related quality of life and related factors in stroke survivors: Data from Korea National Health and Nutrition Examination Survey (KNHANES) 2008 to 2014. PLoS ONE. 2018;13:e0195713

123. Labberton AS, Augestad LA, Thommessen B, Barra M. The association of stroke severity with health-related quality of life in survivors of acute cerebrovascular disease and their informal caregivers during the first year post stroke: a survey study. Qual Life Res. 2020;29:2679–2693

124. Leach MJ, Gall SL, Dewey HM, Macdonell RAL, Thrift AG. Factors associated with quality of life in 7-year survivors of stroke. Journal of Neurology, Neurosurgery & Psychiatry. 2011;82:1365–1371

125. Lee H-Y, Hwang J-S, Jeng J-S, Wang J-D. Quality-Adjusted Life Expectancy (QALE) and Loss of QALE for Patients With Ischemic Stroke and Intracerebral Hemorrhage: A 13-Year Follow-Up. Stroke. 2010;41:739–744

126. Leeds L, Meara J, Hobson P. The impact of discharge to a care home on longer term stroke outcomes. Clin Rehabil. 2004;18:924–928

127. Lindgren P, Glader E-L, Jönsson B. Utility loss and indirect costs after stroke in Sweden. Eur J Cardiovasc Prev Rehabil. 2008;15:230–233.

128. Lopez-Bastida J, Moreno JO, Cerezo MW, Perez LP, Serrano-Aguilar P, Montón-Álvarez F. Social and economic costs and health-related quality of life in stroke survivors in the Canary Islands, Spain. 2012;9.

129. Lu W-S, Huang S-L, Yang J-F, Chen M-H, Hsieh C-L, Chou C-Y. Convergent validity and responsiveness of the EQ-5D utility weights for stroke survivors. J Rehabil Med. 2016;48:346–351.

130. Luengo-Fernandez R, Gray AM, Bull L, Welch S, Cuthbertson F, Rothwell PM, Oxford Vascular Study. Quality of life after TIA and stroke: ten-year results of the Oxford Vascular Study. Neurology. 2013;81:1588–1595

131. Lunde L. Can EQ-5D and 15D be used interchangeably in economic evaluations? Assessing quality of life in post-stroke patients. Eur J Health Econ. 2013;14:539–550

132. Mahesh PKB, Gunathunga MW, Jayasinghe S, Arnold SM, Senanayake S, Senanayake C, De Silva LSD, Kularatna S. Construct validity and reliability of EQ-5D-3L for stroke survivors in a lower middle income setting. Ceylon Med. J 2019;64:52

133. Mar J, Masjuan J, Oliva-Moreno J, Gonzalez-Rojas N, Becerra V, Casado MÁ, Torres C, Yebenes M, Quintana M, et al, on behalf of CONOCES Investigators Group, . Outcomes measured by mortality rates, quality of life and degree of autonomy in thefirst year in stroke units in Spain. Health Qual Life Outcomes. 2015;13:36

134. Mar J, Begiristain JM, Arrazola A. Cost-Effectiveness Analysis of Thrombolytic Treatment for Stroke. Cerebrovasc Dis. 2005;20:193–200 (Mar 2)

135. Mathias SD, Bates MM, Pasta DJ, Cisternas MG, Feeny D, Patrick DL. Use of the Health Utilities Index With Stroke Patients and Their Caregivers. Stroke. 1997;28:1888–1894

136. McDonnell MN, Mackintosh SF, Hillier SL, Bryan J. Regular group exercise is associated with improved mood but not quality of life following stroke. PeerJ. 2014;2:e331

137. Min K -b, Min J -y. Health-related quality of life is associated with stroke deficits in older adults. Age and Ageing. 2015;44:700–704

138. Mittmann N, Chan D, Trakas K, Risebrough N. Health Utility Attributes for Chronic Conditions: Disease Management and Health Outcomes. 2001;9:11–21.

139. Mittmann N, Trakas K, Risebrough N, Liu BA. Utility Scores for Chronic Conditions in a Community-Dwelling Population: PharmacoEconomics. 1999;15:369–376. (Mittmann 2)

140. Oemrawsingh A, van Leeuwen N, Venema E, Limburg M, de Leeuw F-E, Wijffels MP, de Groot AJ, Hilkens PHE, Hazelzet JA, Dippel DWJ, et al. Value-based healthcare in ischemic stroke care: case-mix adjustment models for clinical and patient-reported outcomes. BMC Med Res Methodol. 2019;19:229

141. Olsson BG, Stibrant Sunnerhagen K. Functional and Cognitive Capacity and Health-Related Quality of Life 2 Years After Day Hospital Rehabilitation for Stroke: A Prospective Study. Journal of Stroke and Cerebrovascular Diseases. 2007;16:208–215

142. Olsson BG, Sunnerhagen KS. Effects of Day Hospital Rehabilitation After Stroke. Journal of Stroke and Cerebrovascular Diseases. 2006;15:106–113 (Olsson 2)

143. Park JH, Kim BJ, Bae H-J, Lee J, Lee J, Han M-K, O KY, Park SH, Kang Y, Yu K-H, et al. Impact of Post-Stroke Cognitive Impairment with No Dementia on Health-Related Quality of Life. J Stroke. 2013;15:49

144. Paul SL, Sturm JW, Dewey HM, Donnan GA, Macdonell RAL, Thrift AG. Long-Term Outcome in the North East Melbourne Stroke Incidence Study: Predictors of Quality of Life at 5 Years After Stroke. Stroke. 2005;36:2082–2086

145. Peng L-N, Chen L-J, Lu W-H, Tsai S-L, Chen L-K, Hsiao F-Y. Post-acute care regains quality of life among middle-aged and older stroke patients in Taiwan. Archives of Gerontology and Geriatrics. 2019;83:271–276

146. Peters M, Crocker H, Jenkinson C, Doll H, Fitzpatrick R. The routine collection of patient-reported outcome measures (PROMs) for long-term conditions in primary care: a cohort survey. BMJ Open. 2014;4:e003968

147. Pettersson I, Ahlström G, Törnquist K. The Value of an Outdoor Powered Wheelchair With Regard to the Quality of Life of Persons With Stroke: A Follow-Up Study. Assistive Technology. 2007;19:143–153

148. Phan HT, Gall SL, Blizzard CL, Lannin NA, Thrift AG, Anderson CS, Kim J, Grimley RS, Castley HC, Kilkenny MF, et al. Sex differences in quality of life after stroke were explained by patient factors, not clinical care: evidence from the Australian Stroke Clinical Registry. Eur J Neurol. 2021;28:469–478

149. Phan HT, Blizzard CL, Reeves MJ, Thrift AG, Cadilhac DA, Sturm J, Heeley E, Otahal P, Rothwell P, Anderson CS, et al. Sex Differences in Long-Term Quality of Life Among Survivors After Stroke in the INSTRUCT. Stroke. 2019;50:2299–2306 (Phan 2)

150. Pinto EB, Maso I, Pereira JL, Fukuda TG, Seixas JC, Menezes DF, Cincura C, Neville IS, Jesus PA, Oliveira-Filho J. Differential aspects of stroke and congestive heart failure in quality of life reduction: a case series with three comparison groups. Health Qual Life Outcomes. 2011;9:65

151. Price R, Choy NL. Investigating the Relationship of the Functional Gait Assessment to Spatiotemporal Parameters of Gait and Quality of Life in Individuals With Stroke. Journal of Geriatric Physical Therapy. 2019;42:256–264

152. Ramírez-Moreno JM, Muñoz-Vega P, Alberca SB, Peral-Pacheco D. Health-Related Quality of Life and Fatigue After Transient Ischemic Attack and Minor Stroke. Journal of Stroke and Cerebrovascular Diseases. 2019;28:276–284

153. Ran M, Liu B, Chen L, Zhu C. [Assessing quality of life of patients with stroke using EQ-5D and SF-12]. Sichuan Da Xue Xue Bao Yi Xue Ban. 2015;46:94–98

154. Rivero-Arias O, Ouellet M, Gray A, Wolstenholme J, Rothwell PM, Luengo-Fernandez R. Mapping the Modified Rankin Scale (mRS) Measurement into the Generic EuroQol (EQ-5D) Health Outcome. Med Decis Making. 2010;30:341–354

155. Saarni SI, Härkänen T, Sintonen H, Suvisaari J, Koskinen S, Aromaa A, Lönnqvist J. The impact of 29 chronic conditions on health-related quality of life: a general population survey in Finland using 15D and EQ-5D. Qual Life Res. 2006;15:1403–1414

156. Sallinen H, Sairanen T, Strbian D. Quality of life and depression 3 months after intracerebral hemorrhage. Brain Behav. 2019;9:e01270

157. Sánchez-Iriso E, Errea Rodríguez M, Cabasés Hita JM. Valuing health using EQ-5D: The impact of chronic diseases on the stock of health. Health Economics. 2019;28:1402–1417

158. Sand KM, Wilhelmsen G, Naess H, Midelfart A, Thomassen L, Hoff JM. Vision problems in ischaemic stroke patients: effects on life quality and disability. Eur J Neurol. 2016;23:1–7

159. Sasaki S, Kanai M, Shinoda T, Morita H, Shimada S, Izawa KP. Relation between health utility score and physical activity in community-dwelling ambulatory patients with stroke: a preliminary cross-sectional study. Topics in Stroke Rehabilitation. 2018;25:475–479

160. Slaughter KB, Meyer EG, Bambhroliya AB, Meeks JR, Ahmed W, Bowry R, Behrouz R, Mir O, Begley C, Tyson JE, et al. Direct Assessment of Health Utilities Using the Standard Gamble Among Patients With Primary Intracerebral Hemorrhage. Circ: Cardiovascular Quality and Outcomes [Internet]. 2019 [cited 2021 Feb 15];12 Available from: https://wwwahajournalsorg/doi/101161/CIRCOUTCOMES1195606

161. Sturm JW, Donnan GA, Dewey HM, Macdonell RAL, Gilligan AK, Srikanth V, Thrift AG. Quality of Life After Stroke: The North East Melbourne Stroke Incidence Study (NEMESIS). Stroke. 2004;35:2340–2345

162. Sturm JW, Osborne RH, Dewey HM, Donnan GA, Macdonell RAL, Thrift AG. Brief Comprehensive Quality of Life Assessment After Stroke: The Assessment of Quality of Life Instrument in the North East Melbourne Stroke Incidence Study (NEMESIS). Stroke. 2002;33:2888–2894 (Sturm 2)

163. Szőcs I, Dobi B, Lám J, Orbán-Kis K, Häkkinen U, Belicza É, Bereczki D, Vastagh I. Health related quality of life and satisfaction with care of stroke patients in Budapest: A substudy of the EuroHOPE project. PLoS ONE. 2020;15:e0241059

164. Teoh V, Sims J, Milgrom J. Psychosocial Predictors of Quality of Life in a Sample of Community-Dwelling Stroke Survivors: A Longitudinal Study. Topics in Stroke Rehabilitation. 2009;16:157–166

165. Tran PL, Leigh Blizzard C, Srikanth V, Hanh VTX, Lien NTK, Thang NH, Gall SL. Health-related quality of life after stroke: reliability and validity of the Duke Health Profile for use in Vietnam. Qual Life Res. 2015;24:2807–2814

166. Vahlberg B, Cederholm T, Lindmark B, Zetterberg L, Hellström K. Factors Related to Performance-Based Mobility and Self-reported Physical Activity in Individuals 1-3 Years after Stroke: A Cross-sectional Cohort Study. Journal of Stroke andCerebrovascular Diseases. 2013;22:e426–e434

167. van Eeden M, van Heugten C, van Mastrigt GAPG, van Mierlo M, Visser-Meily JMA, Evers SMAA. The burden of stroke in the Netherlands: estimating quality of life and costs for 1 year poststroke. BMJ Open. 2015;5:e008220

168. Visser MM, Heijenbrok-Kal MH, Spijker A van't, Oostra KM, Busschbach JJ, Ribbers GM. Coping, Problem Solving, Depression, and Health-Related Quality of Life in Patients Receiving Outpatient Stroke Rehabilitation. Archives of Physical Medicine and Rehabilitation. 2015;96:1492–1498

169. Wartenberg KE, Henkner J, Brandt S, Zierz S, Müller TJ. Effect of Recanalization on Cerebral Edema, Long-Term Outcome, and Quality of Life in Patients with Large Hemispheric Infarctions. Journal of Stroke and Cerebrovascular Diseases. 2020;29:105358

170. White J, Magin P, Attia J, Sturm J, McElduff P, Carter G. Predictors of health-related quality of life in community-dwelling stroke survivors: a cohort study. FAMPRJ. 2016;33:382–387

171. Wu M, Brazier J, Relton C, Cooper C, Smith C, Blackburn J. Examining the incremental impact of long-standing health conditions on subjective well-being alongside the EQ-5D. Health Qual Life Outcomes. 2014;12:61

172. Wu M, Brazier JE, Kearns B, Relton C, Smith C, Cooper CL. Examining the impact of 11 long-standing health conditions on health-related quality of life using the EQ-5D in a general population sample. Eur J Health Econ. 2015;16:141–151 (Wu2)

173. Xie J, Wu EQ, Zheng Z-J, Croft JB, Greenlund KJ, Mensah GA, Labarthe DR. Impact of Stroke on Health-Related Quality of Life in the Noninstitutionalized Population in the United States. Stroke. 2006;37:2567–2572

174. Yan P, Zhan F, Hou L, Guo J, He L, Liu D, Zhu C. [Lesion Locations and Quality of Life in Patients with Ischemic Stroke]. Sichuan Da Xue Xue Bao Yi Xue Ban. 2015;46:860–865

175. Yang Y-N, Kim B-R, Uhm KE, Kim SJ, Lee S, Oh-Park M, Lee J. Life Space Assessment in Stroke Patients. Ann Rehabil Med. 2017;41:761

176. Yeoh YS, Koh GC-H, Tan CS, Lee KE, Tu TM, Singh R, Chang HM, De Silva DA, Ng YS, Ang YH, et al. Can acute clinical outcomes predict health-related quality of life after stroke: a one-year prospective study of stroke survivors. Health QualLife Outcomes. 2018;16:221

177. Yeoh YS, Koh GC-H, Tan CS, Tu TM, Singh R, Chang HM, De Silva DA, Ng YS, Ang YH, Yap P, et al. Health-related quality of life loss associated with first-time stroke. PLoS ONE. 2019;14:e0211493 (Yeoh 2)