Appendix 1 Searching strategies

MEDLINE (PubMed)

Search date: March 16, 2022

Results: 1,908

#1: "hypertension, pregnancy induced"[MeSH Terms] OR ("hypertension"[All Fields] AND "pregnancy induced"[All Fields]) OR "pregnancy-induced hypertension"[All Fields] OR ("gestational"[All Fields] AND "hypertension"[All Fields]) OR "gestational hypertension"[All Fields] OR ("hypertension, pregnancy induced"[MeSH Terms] OR ("hypertension"[All Fields] AND "pregnancy induced"[All Fields]) OR "pregnancy-induced hypertension"[All Fields] OR ("pregnancy"[All Fields] AND "induced"[All Fields] AND "hypertension"[All Fields]) OR "pregnancy induced hypertension"[All Fields]) OR ("pre eclampsia"[MeSH Terms] OR "pre eclampsia"[All Fields] OR "preeclampsia"[All Fields]) OR ("eclampsia"[MeSH Terms] OR "eclampsia"[All Fields] OR "eclampsias"[All Fields])

#2: "postpartum period"[MeSH Terms] OR ("postpartum"[All Fields] AND "period"[All Fields]) OR "postpartum period"[All Fields] OR "postpartum"[All Fields]

#3: "cardiovascular diseases"[MeSH Terms] OR ("cardiovascular"[All Fields] AND "diseases"[All Fields]) OR "cardiovascular diseases"[All Fields] OR ("cardiovascular"[All Fields] AND "disease"[All Fields]) OR "cardiovascular disease"[All Fields]

#4: #1 AND #2 AND #3

Cochrane library

Search date: March 16, 2022

Results: 72

#1: “gestational hypertension OR pregnancy induced hypertension OR preeclampsia OR eclampsia”

#2: “postpartum”

#3: “cardiovascular disease”

#4: (#1) AND (#2) AND (#3)

Web of Science

Search date: March 16, 2022

Results: 256

#1: ALL=(“gestational hypertension”) OR ALL=(“pregnancy induced hypertension”) OR ALL=(“preeclampsia”) OR ALL=(“eclampsia”)

#2: ALL=(“postpartum”)

#3: ALL=(“cardiovascular disease”)

#4: #1 AND #2 AND #3

#5: #1 AND #2 AND #3 AND DOCUMENT TYPES: (ARTICLE)

Scopus

Search date: March 16, 2022

Results: 2,495

#1: ("gestational hypertension" OR "pregnancy induced hypertension" OR "preeclampsia" OR "eclampsia") AND "postpartum" AND "cardiovascular disease" AND (LIMIT-TO (SRCTYPE , "j")) AND (LIMIT-TO (DOCTYPE , "ar"))

Appendix 2 Criteria for the Newcastle-Ottawa Scale regarding star allocation to assess quality of studies

Criteria	**Acceptable (star given)**	**Unacceptable (star not given)**
Selection		
Representativeness of the exposed cohort	Population- or hospital-based	Selected group of women
Selection of the non-exposed cohort	Drawn from the same source as the exposed cohort	Drawn from a different source
Ascertainment of exposure	Secure record or structured interview	Written self-report
Demonstration that outcome of interest was not present at the start of the study	Demonstrated in exclusion criteria or analyzed in separated group	Did not demonstrate
Comparability		
Comparability of cohorts on the basis of the design or analysis	Matched or adjusted for age, BMI, and smoking	Did not match or adjust for age, BMI, and smoking
	Matched or adjusted for any additional factor	Did not match or adjust for any additional factor
Outcome		
Assessment of outcome	Directly measured or record linkage	Self-report or no information regarding definition of outcome
Follow-up long enough for outcomes to occur	Follow-up ≥ 5 years	Follow-up < 5 years or no information regarding follow-up duration
Adequacy of follow up of cohorts	Follow-up rate ≥ 80 %	Follow-up rate < 80 % or no information regarding follow-up rate

BMI = body mass index

Review: HDP and risk of any cardiovascular diseases in later life
Comparison: HDP vs. Normotensive
Outcome: Composite cardiovascular and cerebrovascular diseases



Review: HDP and risk of mortality in later life
Comparison: HDP vs. Normotensive
Outcome: Mortality




Standard Error
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0.01
0.10
1.00
10.00
100.00
1000.00
Risk Ratio

Supplementary Table 1 Quality assessment of included studies

Author, year	Study population	Ascertainment of exposure	Ascertainment of outcome	Completeness of follow-up	Degree of adjustment*
Kestenbaum, 2003[61]	All singleton births recorded in Washington State Birth Events Record Database (BERD), 1987–1998	ICD codes and diagnoses in birth certificate	ICD codes during hospitalization	No information	Age, parity, year of delivery, race/ethnicity, GDM, smoking, cesarean section, type of insurance used for the index pregnancy
Lykke, 2009[34]	All singleton deliveries in Denmark, January 1978 – October 2007	ICD codes in National Patient Registry	ICD codes in National Patient Registry	99.2% complete data	Age at delivery, year of delivery, preterm delivery, SGA, placental abruption, stillbirth, DM subsequent to the index pregnancy
Edlow, 2009[62]	Women enrolled in the Preeclampsia: Mechanisms and Consequences (PMC) study at the Hospital of the University of Pennsylvania, January 2006 – October 2006	Maternal criteria for preeclampsia	Telephone interviews by the primary author	70% of preeclampsia and 59% of controls completed the follow-up study	Race, BMI, parity, chronic hypertension
Garovic, 2010[63]	Women participating in the Family Blood Pressure Program (FBPP) study which consists of four different research networks in United States	Interviews by trained examiners	Self-reports and blood pressure measurement	No information	Race, network, family history of cardiovascular disease, education, DM, smoking, BMI, hypertension, dyslipidemia

Mongraw-Chaffin, 2010[64]	Women enrolled in the original Child Health and Development Studies (CHDS) cohort at the East Bay Area of California, 1959–1967	Medical records abstraction	Results from the California Vital Status matches to CHDS files	No information	IUGR child, preexisting hypertension, age, BMI, current smoking at enrollment
Melchiorre, 2011[35]	Women visiting Fetal-Maternal Medicine Unit of St George's Hospital, United Kingdom, January 2008 – December 2010	Diagnoses in medical records	No definition of outcome (hypertension)	83.1% of preeclampsia and 100% of controls complete data	Age, ethnicity, gestational age at echocardiography
Drost, 2012[36]	Women registered in the obstetric database at the Isala Klinieken in Zwolle, Netherlands, 1991–2007	Diagnoses in medical records	Blood pressure measurement	83.9% of preeclampsia and 73.7% of controls complete data	Age, years postpartum, current smoking
Zhao, 2012[75]	Female employees giving birth at the Kailuan Medical Group, China, October 1976 – December 2008	Delivery data filled by trained medical staff	Diagnoses in medical records	83.7% of preeclampsia and 83.3% of controls complete data	Childbearing age, age at the 2006-2007 physical examination, BMI, blood pressure, fasting blood glucose, total cholesterol, triglycerides, LDL-C, HDL-C, hs-CRP
Collen, 2013[37]	Women giving birth at the Sahlgrenska University Hospital and Östra Hospital, Sweden, 1969–1973	No information	No definition of outcome (hypertension)	94.6% complete data	Age
Callaway, 2013[87]	Women delivering a single live birth at a major public hospital in South Brisbane, Australia, 1981–1983	Diagnoses recorded by obstetricians	Blood pressure measurement	No information	Age, ethnicity, education, alcohol intake, number of cigarettes smoked, exercise, BMI

Shalom, 2013[88]	Women giving birth at the Soroka University Medical Center, Israel, January 1988 – December 1998	Diagnoses in obstetric database	Diagnoses in Demog-ICD9 database	No information	None
Östlund, 2013[38]	Women giving birth at the Karolinska University Hospital, Sweden	Diagnoses in hospital records	No definition of outcome (hypertension)	88.9% of preeclampsia and 94.1% of controls complete data	Age, parity, date of delivery
Kvehaugen, 2014[39]	Women participating in the phase 2 of the Nord-Trøndelag Health Study (HUNT), Norway, 1995–1997	Diagnoses in Medical Birth Registry of Norway (MBRN)	Self-reported questionnaires and blood pressure measurement	No information	Age
Zhou, 2014[76]	Women delivering at the Guangdong Women and Children's Hospital Affiliated to Guangzhou Medical University, China, January 2008 – June 2012	Diagnoses in medical records	Blood pressure measurement	82.8% of preeclampsia and 79.4% of controls complete data	Prenatal, postpartum, and prenatal and postpartum difference in: BMI, blood pressure, fasting blood glucose, fasting insulin, total cholesterol, triglycerides, HDL-C, LDL-C
Ghossein-Doha, 2014[40]	Women participating in a previous study in the Netherlands, 1996–1999	Diagnoses in medical records	Blood pressure measurement	80% of preeclampsia and 61% of controls complete data	Parity
Yeh, 2014[77]	Random selection of women from the National Health Insurance	ICD codes in medical records	ICD codes in medical records	No information	Age, date of delivery

	Research Database (NHIRD) in Taiwan, January 1997 – December 2009				
Breetveld, 2014[41]	Preeclampsia from the 6-week postpartum screening and controls from advertisement, tertiary referral hospital in the Netherlands	Diagnoses in pregnancy charts	No definition of outcome (hypertension)	No information	None
Ehrenthal, 2015[65]	Women visiting postpartum service of an academic community hospital, 2011–2012	Diagnoses in inpatient records	Blood pressure measurement	No information	Age, BMI, race, family history of cardiovascular disease, tobacco use, insurance
Behrens, 2016[42]	Women in Denmark with at least one pregnancy, 1978–2012	ICD codes in National Patient Register and Medical Birth Register	ICD codes in National Patient Register or Causes of Death Register	97.5% complete data	Maternal birth year, parity, multiple pregnancy, stillbirth
Cain, 2016[66]	Singleton first birth to women aged 15–49 years in Florida, January 2004 – December 2007	ICD codes in Florida's Statewide maternal and infant linked database	ICD codes in hospital discharge data	No information	Age, race/ethnicity, nativity, education, income, history of hyperlipidemia, migraine, and lupus, pre-pregnancy BMI, GDM, tobacco use, drug use, infant sex
Nelander, 2016[43]	All twins in the nationwide Swedish Twin Register born in 1958 or earlier, Sweden	Telephone interview	ICD codes in National Patient Register and Cause of Death	No information	Age at interview, current BMI, education, current smoking

			Register		
Pérez-Adan, 2016[44]	Women giving birth in the Complexo Hospitalario Universitario de Ourense, Spain, January 2000 – December 2010	Diagnoses in hospital records	Diagnoses in hospital records	90.3% complete data	None
Grandi, 2017[45]	First birth for women aged 15–45 years in United Kingdom, January 1990 – December 2013	Read codes in United Kingdom's Clinical Practice Research Datalink (CPRD)	Read codes in United Kingdom's Clinical Practice Research Datalink (CPRD)	No information	Age, smoking, BMI, excessive alcohol use, year of cohort entry, region of residence, multiple gestation at first pregnancy, depression, dyslipidemia, venous thromboembolism, polycystic ovary syndrome, GDM, DM, renal disease, migraines, family history of hypertension and cardiovascular disease, number of distinct drug classes prescribed, use of statin, aspirin, anti-depressant medications, NSAIDs, oral contraceptives, and anti-migraine medications
Timpka, 2017[46]	Female registered nurses aged 25–42 years in 1989, United States	Self-reported questionnaires	Self-reports questionnaires	No information	Age, race/ethnicity, parity, BMI at age 18, updated smoking status, alcohol use, NSAIDs use, menopausal status, history of GDM, updated BMI, physical

					activity, DASH diet, sodium/potassium intake
Dunietz, 2017[67]	Women receiving prenatal care from 52 clinics in five Michigan communities, 1998–2004	Diagnoses in medical records	Blood pressure measurement	52.1% complete data	Race/ethnicity, pre-pregnancy BMI, age, parity, Medicaid insurance coverage at the time of the Pregnancy Outcomes and Community Health (POUCH) Study pregnancy, time between the POUCH Study pregnancy and follow-up
Best, 2017[68]	American Indian women participating in a previous study, January 1995 – December 2012	Diagnoses in medical records	Diagnoses in medical records	No information	Systolic blood pressure, current age, current BMI
Ghossein-Doha, 2017[47]	Preeclampsia from postpartum screening and healthy parous controls from advertisement	Diagnoses in pregnancy charts	Blood pressure measurement	No information	Parity
Bokslag, 2018[48]	Women giving birth at two tertiary medical centers in the Netherlands, 1998–2005	Diagnoses in medical records	Blood pressure measurement	97.8% of preeclampsia and 91.8% of controls complete data	Age, date of delivery, smoking status, blood pressure, BMI, educational level
Li, 2018[78]	Women participating in the Growing Up in Singapore Towards Healthy Outcomes (GUSTO) cohort	Self-reported questionnaires	Blood pressure measurement	90% complete data	Age, ethnicity, college education, pre-pregnancy BMI, parity
Chen,	Women aged 20–50 years admitted	ICD codes in	ICD codes in	No information	Age and year of delivery, parity,

2018[79]	for delivery in Taiwan, January 2000 – December 2013	medical records	medical records		gestational age, gestational number, length of follow-up, comorbidities (DM, hypertension, dyslipidemia, coronary artery disease, chronic kidney disease, COPD, cerebrovascular disease)
Bergen, 2018[49]	Women participating in the Generation R Study, Netherlands	Diagnoses in hospital charts	Blood pressure measurement	97.7% complete data	Age at intake, visit interval, ethnicity, educational level, smoking, subsequent pregnancies between index and follow-up, child's sex, BMI at follow-up
Theilen, 2018[69]	Women giving birth in Utah, 1939–2012	Diagnoses in birth certificate data	Utah Population Database, Utah death certificates, Social Security Death Index	No information	5-year age groups, year of childbirth, parity, infant sex, gestational age at delivery, parental education, race/ethnicity, marital status
Basit, 2018[50]	Women in Denmark with at least one pregnancy, 1978–2015	ICD codes in National Patient Register	ICD codes in National Patient Register	No information	Age, parity, maternal birth year, region in which the child was delivered
Kuo, 2018[80]	Random selection of women from the National Health Insurance Research Database (NHIRD) in Taiwan, January 1996 – December 2010	ICD codes in medical records	ICD codes in medical records	No information	Age, date of delivery
Egeland,	Women participating in the	Diagnoses in	Pharmacologically	No information	Age at delivery, age, BMI,

2018[51]	Norwegian Mother and Child Cohort Study (MoBa) from all regions in Norway, 1999–2008	Medical Birth Registry of Norway (MBRN)	treated hypertension in Norwegian Prescription Database		educational level, physical activity, daily smoking, alcohol consumption frequency, duration of lifetime oral contraceptive use, mid-pregnancy poor diet quality, total energy intake, multiple birth pregnancies at delivery
Haas, 2019[70]	First births in 8 United States sites (Case Western University; Columbia University; Indiana University; University of Pittsburgh; Northwestern University; University of California at Irvine; University of Pennsylvania; and University of Utah), 2010–2013	Diagnoses in medical records	Blood pressure measurement	79.2% complete data	Age, BMI, type of health insurance at Nulliparous Pregnancy Outcomes Study—Monitoring Mothers-to-be Heart Health Study (nuMoM2b) enrollment, race/ethnicity, smoking during the 3 months before pregnancy
Haug, 2019[52]	Women from the Nord-Trøndelag Health Study (HUNT) with at least one birth, Norway, 1967–2012	Diagnoses in Medical Birth Registry of Norway (MBRN)	Diagnoses in hospital records	91.7% complete data	Age, maternal birth year, highest educational level, ever daily smoking, parity before age 40 years, family history of coronary heart disease
Amiri, 2019[81]	Women aged 20–50 years with at least one pregnancy at the beginning of the Tehran Lipid and Glucose Study (TLGS), Iran, 1998–2014	Self-reports and diagnoses hospital records	Blood pressure measurement	95.5% of preeclampsia and 90.0% of controls complete data	None

Smith, 2019[71]	Pre-Eclampsia New Emerging Team (PE-NET) cohort from preeclampsia and normotensive women at the Kingston or Ottawa General Hospitals; Maternal Health Clinic (MHC) cohort from women referred with hypertensive pregnancy, November 2011 – January 2017	Diagnoses in prenatal records	Blood pressure measurement	No information	Age
Honigberg, 2019[53]	Women aged 40–69 years with at least one pregnancy at the baseline of the UK Biobank study visit, 2006–2010	ICD codes in hospital records or self-reports at enrollment	ICD codes in hospital records and primary care diagnoses	99.7% complete data	Age, race, BMI, ever-smoking, prevalent hypertension, hyperlipidemia, and DM
Ernawati, 2019[82]	Women with severe preeclampsia who carried out pregnancy termination in dr. Soetomo General Hospital of Surabaya, Indonesia, 2013 - January 2014	Diagnoses in medical records	Blood pressure measurement	54.8% of early-onset and 39.0% of late-onset complete data	None
Osoti, 2019[89]	Women aged 28 years or older delivering at Kenyatta National Hospital, Kemya	Diagnoses in inpatient records	Blood pressure measurement	91% complete data	Age, level of education, BMI, hormonal contraception, breastfeeding, marital and employment status
Leon, 2019[54]	Completed pregnancy registered in United Kingdom's Clinical Practice Research Datalink (CPRD), January 1997 – December 2016	Read and ICD codes in linked participant records	Read and ICD codes in linked participant records	No information	Ethnicity, age, pre-pregnancy diabetes and hypertension, index of multiple deprivation, cluster term

Arnaout, 2019[72]	First birth to women aged 18 years or older in California, 2005–2009	ICD codes in Healthcare Cost and Utilization Project (HCUP) database	ICD codes in HCUP database	99% complete data	Age, race, insurance status, median household income, chronic kidney disease, pre-existing diabetes, obesity, drug abuse, smoking, multiple gestations
Huang, 2020[83]	Random selection of women from the National Health Insurance Research Database (NHIRD) in Taiwan, January 2000 – December 2013	ICD codes in medical records	ICD codes in medical records	No information	Index day, age with propensity scores
Scheres, 2020[55]	Women in Denmark with at least one pregnancy, 1999–2012	Diagnoses in Dutch Perinatal Registry (Perined)	Diagnoses in anticoagulation clinics	93% complete data	Number of pregnancies, age at start of follow-up, self-reported ancestry
Wagata, 2020[84]	Women aged 20 years or older in Miyagi and Iwate Prefectures, Japan, May 2013 – March 2016	Self-reported questionnaires	Blood pressure measurement	99.9% complete data	Age, BMI, family history of hypertension, alcohol consumption
Garovic, 2020[73]	All live birth or stillborn deliveries in Olmsted County, Minnesota, January 1976 – December 1982	Rochester Epidemiology Project (REP) medical record-linkage system	Diagnoses in inpatient and outpatient visits to REP-affiliated providers	98.7% complete data	Date of delivery, age, parity at index pregnancy, education, smoking, obesity
Bergman, 2020[56]	All deliveries in Sweden, January 1973 – December 2010	ICD codes in Swedish Medical	ICD codes in MBR	Complete follow-up	Age at delivery, chronic hypertension before first birth,

		Birth Register (MBR)			level of education, time period of birth
Moe, 2020[57]	Women aged 18 years or older delivering a singleton baby at Oslo University Hospital, Norway, September 2014 – April 2018	Diagnoses in medical records	Blood pressure measurement	Complete follow-up	None
Sharma, 2021[58]	Women with pregnancies in the Helsingborg area who delivered at the hospital in the Helsingborg birth cohort (HbgBC) from Sweden linked to registries of the National Board of Health and Welfare in Sweden, February 1964 – January 1967	ICD codes in national registers	ICD codes in national registers	Complete follow-up	Age at delivery, maternal weight at first antenatal visit, smoking in pregnancy, parity
Mooij, 2021[90]	Women who were treated in Ndala Hospital for severe preeclampsia and eclampsia, 2011–2012	Diagnoses in medical records	Blood pressure measurement	Complete follow-up	None
Ntlemo, 2021[91]	Women delivered at two tertiary-level hospitals in Pretoria, February 2019 – April 2020	Daily morning ward rounds	Blood pressure measurement	95.8% complete data	None
Nuckols, 2021[74]	Women between 18 and 40 years of age, 10–36 months postpartum	Diagnoses in medical records	Blood pressure measurement	Complete follow-up	None
Oliver-Williams, 2022[59]	Women who had one or more singleton live births in England recorded in national medical records in all National Health Service	ICD codes in medical records	ICD codes in medical records	No information	Maternal age at delivery, socioeconomic status, ethnicity, hospital-recorded diabetes, diagnosed prior to pregnancy,

	hospitals identified from the Hospital Episode Statistics (HES) database, 1997–2015				year of delivery
Park, 2022[85]	Women who gave birth and recorded in the Korean National Health Insurance Database by the Health Insurance Review & Assessment Service, January 2007 – December 2015	ICD codes in medical records	ICD codes in medical records	Complete follow-up	Age, primipara, cesarean section, and multiple pregnancy in all data sets; body mass index; systolic and diastolic blood pressures; fasting blood glucose; aspartate aminotransferase levels; alanine aminotransferase levels; total cholesterol levels; and current smoking
Hung, 2022[86]	Matching 1:4 selection of women aged 18–45 years without a previous history of stroke registered in the National Health Insurance Research Database (NHIRD) in Taiwan, 2000–2017	ICD codes in medical records	ICD codes in medical records	No information	Age, delivery type, gestation number, hospital level, delivery season, living area, family income level and all comorbidities
Kennedy, 2022[92]	Women gave birth to a live baby at St George Hospital, Sydney, completed at least the 6-month postpartum assessment, and enroll in the subsequent pregnancy component of the P4 study	Diagnoses in inpatient records	Blood pressure measurement	No information	Age and ethnicity, physiological measures (SBP, DBP, fat mass percentage) and interpregnancy factors of BMI change, inter-pregnancy interval and breastfeeding duration
van Baar,	Nulliparous women delivered in the	ICD codes in the	ICD codes in the	96.9% Complete	None

2022[60]	Netherlands and registered in the Perinatal Registry of the Netherlands (PERINED) database and the National Death Registry of the Netherlands (Central Bureau for Statistics, CBS), 1995–2018	Perinatal Registry of the Netherlands (PERINED) database	Perinatal Registry of the Netherlands (PERINED) database	data	

*List of covariates that were adjusted for or matched

Abbreviation: ICD = International Classification of Diseases, GDM = gestational diabetes mellitus, SGA = small-for-gestational age, DM = diabetes mellitus, BMI = body mass index, IUGR = intrauterine growth restriction, LDL-C = low-density lipoprotein cholesterol, HDL-C = high-density lipoprotein cholesterol, hs-CRP = high-sensitivity C-reactive protein, NSAID = non-steroidal anti-inflammatory drug, DASH = diet foods for high blood pressure, COPD = chronic obstructive pulmonary disease; SBP = systolic blood pressure; DBP = diastolic blood pressure

Supplementary Table 2 Risk of bias assessment of included studies

Author, year	Representative of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Outcome of interest was not present at start of study	Comparability of cohorts	Assessment of outcome	Follow-up long enough	Adequacy of follow-up	Total score
Kestenbaum, 2003[61]	*	*	*	*	*	*	*		7
Lykke, 2009[34]	*	*	*	*	*	*	*	*	8
Edlow, 2009[62]	*	*	*		*				4
Garovic, 2010[63]	*	*	*	*	**		*		7
Mongraw-Chaffin, 2010[64]	*	*	*	*	**	*	*		8
Melchiorre, 2011[35]	*	*	*		*			*	5
Drost, 2012[36]	*	*	*		*	*	*		6
Zhao, 2012[75]	*	*	*	*	*	*	*	*	8
Collen, 2013[37]	*	*			*		*	*	5
Callaway, 2013[87]	*	*	*	*	**	*	*		8
Shalom, 2013[88]	*	*	*	*		*	*		6
Östlund,	*	*	*	*	**		*	*	8

2013[38]									
Kvehaugen, 2014[39]	*	*	*	*	*		*		6
Zhou, 2014[76]	*	*	*	*	*	*			6
Ghossein-Doha, 2014[40]	*	*	*	*	*	*	*		7
Yeh, 2014[77]	*	*	*	*	*	*	*		7
Breetveld, 2014[41]	*	*	*	*			*		5
Ehrenthal, 2015[65]	*	*	*	*	**	*			7
Behrens, 2016[42]	*	*	*	*	*	*	*	*	8
Cain, 2016[66]	*	*	*	*	**	*			7
Nelander, 2016[43]		*	*		**	*	*		6
Pérez-Adan, 2016[44]	*	*	*		*	*	*	*	7
Grandi, 2017[45]	*	*	*	*	**	*	*		8
Timpka, 2017[46]		*		*	**				4
Dunietz, 2017[67]	*	*	*		*	*	*		6

Best, 2017[68]	*	*	*	*	**	*	*		8
Ghossein-Doha, 2017[47]	*	*	*	*		*	*		6
Bokslag, 2018[48]	*	*	*	*	**	*	*	*	9
Li, 2018[78]		*		*	*	*	*	*	6
Chen, 2018[79]	*	*	*	*	*	*	*		7
Bergen, 2018[49]	*	*	*	*	**	*	*	*	9
Theilen, 2018[69]	*	*	*	*	*	*	*		7
Basit, 2018[50]	*	*	*	*	*	*	*		7
Kuo, 2018[80]	*	*	*	*	*	*	*		7
Egeland, 2018[51]	*	*	*	*	**	*	*		8
Haas, 2019[70]	*	*	*	*	**	*	*		8
Haug, 2019[52]	*	*	*	*	*	*	*	*	8
Amiri, 2019[81]	*	*		*		*	*	*	6
Smith, 2019[71]	*		*		*	*			4
Honigberg, 2019[53]	*	*	*	*	**	*	*	*	9
Ernawati, 2019[82]	*	*	*	*	*	*	*		7
Osoti, 2019[89]	*	*	*	*	*	*		*	7

Leon, 2019[54]	*	*	*	*	*	*	*		7
Arnaout, 2019[72]	*	*	*	*	**	*		*	8
Huang, 2020[83]	*	*	*	*	*	*	*		7
Scheres, 2020[55]	*	*	*	*	*	*	*	*	8
Wagata, 2020[84]	*	*			*	*	*	*	6
Garovic, 2020[73]	*	*	*	*	**	*	*	*	9
Bergman, 2020[56]	*	*	*	*	*	*	*	*	8
Moe, 2020[57]	*	*	*	*	*	*		*	7
Sharma, 2021[58]	*	*	*	*	**	*	*	*	9
Mooij, 2021[90]		*	*	*		*	*	*	6
Ntlemo, 2021[91]	*	*	*	*		*		*	6
Nuckols, 2021[74]	*	*	*	*		*		*	6
Oliver-Williams, 2022[59]	*	*	*	*	**	*	*		8
Park, 2022[85]	*	*	*	*	**	*	*	*	9

Hung, 2022[86]	*	*	*	*	*	*	*		7
Kennedy, 2022[92]	*	*	*	*	**	*			7
van Baar, 2022[60]	*	*	*	*		*	*	*	7

Supplementary Table 3 Summary of findings table for each outcome

Outcomes	RR (95% CI)	Certainty of evidence
Prior HDP vs normotensive pregnancies		
Hypertension at any time (n= 37)		⨁⨁⊖⊖ Low[1,2,3,4]
Hypertension <= 5 years (n = 7)	5.34 (2.74, 10.39)	⨁⊖⊖⊖ Very low[1,3,4]
Hypertension 6-10 years (n = 6)	4.22 (2.19, 8.10)	⨁⊖⊖⊖ Very low[1,3,4]
Hypertension 11-15 years (n = 7)	3.27 (2.02, 5.30)	⨁⨁⊖⊖ Low[1,2,3,4]
Hypertension >=15 years (n = 2)	1.79 (1.22, 2.61)	⨁⨁⊖⊖ Low[1,4]
Hypertension unspecified (n =15)	3.10 (1.81. 5.29)	⨁⨁⊖⊖ Low[1,3]
Ischemic heart disease (n = 10)	2.06 (1.38, 3.08)	⨁⨁⊖⊖ Low[1,3]
Heart failure (n = 7)	2.53 (1.28, 5.00)	⨁⊖⊖⊖ Very low[1,3,4]
Venous thromboembolism (n = 4)	1.34 (0.80, 2.24)	⨁⊖⊖⊖ Very low[1,3,4,5]
Peripheral vascular disease (n = 3)	1.23 (0.26, 5.76)	⨁⊖⊖⊖ Very low[1,4,5]
Stroke (n = 12)	1.59 (1.08, 2.33)	⨁⨁⊖⊖ Low[1,3]
Dementia (n = 3)	1.37 (0.70, 2.71)	⨁⊖⊖⊖ Very low[1,4,5]
Composite cardiovascular and cerebrovascular diseases (n = 8)	1.91 (1.18, 3.09)	⨁⊖⊖⊖ Very low[1,3,4,5]
Mortality (n = 2)	2.15 (0.70, 6.58)	⨁⊖⊖⊖ Very low[1,3,4,5]
Preeclampsia with severe features vs normotensive pregnancies		
Hypertension at any time (n = 3)	6.67 (1.51, 29.40)	⨁⊖⊖⊖ Very low[1,3,4,5]
Ischemic heart disease (n = 2)	2.11 (0.04, 113.97)	⨁⊖⊖⊖ Very low[1,3,4,6]
Venous thromboembolism (n = 2)	2.00 (0.41, 9.78)	⨁⊖⊖⊖ Very low[1,4,6]
Preeclampsia without severe features vs normotensive pregnancies		
Hypertension at any time (n = 2)	7.21 (0, 17565)	⨁⊖⊖⊖ Very low[1,3,4,6]
Ischemic heart disease (n = 2)	1.92 (1.09, 3.38)	⨁⊖⊖⊖ Very low[1,4,5]
Venous thromboembolism (n = 2)	1.64 (1.55, 1.73)	⨁⨁⊖⊖ Low[1,4]
Early onset preeclampsia vs normotensive pregnancies		
Hypertension at any time (n = 2)	2.53 (1.93, 3.32)	⨁⨁⊖⊖ Low[1,4]
The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the		

comparison group and the relative effect of the intervention (and its 95% CI). CI: Confidence interval; RR: Relative Risk

GRADE Working Group grades of evidence

- High quality: Further research is very unlikely to change our confidence in the estimate of effect.
- Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
- Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.
- Very low quality: We are very uncertain about the estimate.

Footnotes

[1]We downgraded (1) level for serious limitation in study design.
[2]We upgraded (1) level for magnitude of effect.
[3]We downgraded (1) level for serious inconsistency.
[4]We downgraded (1) level for serious publication bias due to p value <0.05.
[5]We downgraded (1) level for serious imprecision due to wide confidence interval.
[6]We downgraded (2) level for very serious imprecision due to wide confidence interval.