






Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 0.0% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 5.5% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.0% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 11.7% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 9.0% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 7.3% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 91 91 39.5% -7.25 [-15.39, 0.89]
Heterogeneity: Tau² = 57.65; Chi² = 13.25, df = 4 (P = 0.01); I² = 70%
Test for overall effect: Z = 1.75 (P = 0.08)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 11.8% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 13.8% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 13.6% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 12.6% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 8.7% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 122 122 60.5% -2.68 [-5.73, 0.37]
Heterogeneity: Tau² = 4.56; Chi² = 6.52, df = 4 (P = 0.16); I² = 39%
Test for overall effect: Z = 1.72 (P = 0.08)
Total (95% CI) 213 213 100.0% -4.81 [-8.82, -0.80]
Heterogeneity: Tau² = 26.05; Chi² = 29.99, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.35 (P = 0.02)
Test for subgroup differences: Chi² = 1.06, df = 1 (P = 0.30), I² = 5.8%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]



Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.6% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 0.0% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.3% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 12.2% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 9.4% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 7.7% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 83 83 37.2% -8.45 [-17.05, 0.15]
Heterogeneity: Tau² = 57.63; Chi² = 12.15, df = 4 (P = 0.02); I² = 67%
Test for overall effect: Z = 1.93 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 12.3% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.2% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 14.1% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 13.1% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 9.2% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 122 122 62.8% -2.68 [-5.73, 0.37]
Heterogeneity: Tau² = 4.56; Chi² = 6.52, df = 4 (P = 0.16); I² = 39%
Test for overall effect: Z = 1.72 (P = 0.08)
Total (95% CI) 205 205 100.0% -5.07 [-9.21, -0.92]
Heterogeneity: Tau² = 27.08; Chi² = 30.34, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.39 (P = 0.02)
Test for subgroup differences: Chi² = 1.53, df = 1 (P = 0.22), I² = 34.8%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]



Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.6% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 5.8% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 0.0% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 12.3% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 9.4% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 7.7% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 82 82 36.9% -8.09 [-16.97, 0.79]
Heterogeneity: Tau² = 62.24; Chi² = 12.60, df = 4 (P = 0.01); I² = 68%
Test for overall effect: Z = 1.79 (P = 0.07)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 12.4% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.3% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 14.1% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 13.1% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 9.2% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 122 122 63.1% -2.68 [-5.73, 0.37]
Heterogeneity: Tau² = 4.56; Chi² = 6.52, df = 4 (P = 0.16); I² = 39%
Test for overall effect: Z = 1.72 (P = 0.08)
Total (95% CI) 204 204 100.0% -4.95 [-9.12, -0.78]
Heterogeneity: Tau² = 27.24; Chi² = 30.37, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.33 (P = 0.02)
Test for subgroup differences: Chi² = 1.28, df = 1 (P = 0.26), I² = 21.6%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]



Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 0.5% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 2.3% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 2.6% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 0.0% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 5.7% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 3.7% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 64 64 14.8% -3.60 [-9.11, 1.90]
Heterogeneity: Tau² = 0.00; Chi² = 0.55, df = 4 (P = 0.97); I² = 0%
Test for overall effect: Z = 1.28 (P = 0.20)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 12.5% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 26.7% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 24.6% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 16.1% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 5.3% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 122 122 85.2% -2.68 [-5.73, 0.37]
Heterogeneity: Tau² = 4.56; Chi² = 6.52, df = 4 (P = 0.16); I² = 39%
Test for overall effect: Z = 1.72 (P = 0.08)
Total (95% CI) 186 186 100.0% -2.73 [-4.85, -0.61]
Heterogeneity: Tau² = 0.00; Chi² = 7.19, df = 9 (P = 0.62); I² = 0%
Test for overall effect: Z = 2.53 (P = 0.01)
Test for subgroup differences: Chi² = 0.08, df = 1 (P = 0.77), I² = 0%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]



Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.7% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 6.1% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.6% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 12.6% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 0.0% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 8.0% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 80 80 35.0% -8.94 [-17.58, -0.29]
Heterogeneity: Tau² = 52.54; Chi² = 9.91, df = 4 (P = 0.04); I² = 60%
Test for overall effect: Z = 2.03 (P = 0.04)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 12.7% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.7% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 14.5% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 13.5% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 9.5% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 122 122 65.0% -2.68 [-5.73, 0.37]
Heterogeneity: Tau² = 4.56; Chi² = 6.52, df = 4 (P = 0.16); I² = 39%
Test for overall effect: Z = 1.72 (P = 0.08)
Total (95% CI) 202 202 100.0% -5.16 [-9.44, -0.88]
Heterogeneity: Tau² = 28.09; Chi² = 30.26, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.36 (P = 0.02)
Test for subgroup differences: Chi² = 1.79, df = 1 (P = 0.18), I² = 44.0%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]



Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.6% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 5.9% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.4% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 12.4% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 9.6% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 0.0% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 80 80 36.0% -8.73 [-17.42, -0.05]
Heterogeneity: Tau² = 56.04; Chi² = 11.18, df = 4 (P = 0.02); I² = 64%
Test for overall effect: Z = 1.97 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 12.5% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.5% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 14.3% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 13.3% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 9.4% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 122 122 64.0% -2.68 [-5.73, 0.37]
Heterogeneity: Tau² = 4.56; Chi² = 6.52, df = 4 (P = 0.16); I² = 39%
Test for overall effect: Z = 1.72 (P = 0.08)
Total (95% CI) 202 202 100.0% -5.13 [-9.34, -0.92]
Heterogeneity: Tau² = 27.49; Chi² = 30.28, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.39 (P = 0.02)
Test for subgroup differences: Chi² = 1.66, df = 1 (P = 0.20), I² = 39.7%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]


Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.7% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 6.1% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.6% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 13.1% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 10.0% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 8.1% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 96 96 45.6% -7.69 [-15.26, -0.11]
Heterogeneity: Tau² = 50.23; Chi² = 13.31, df = 5 (P = 0.02); I² = 62%
Test for overall effect: Z = 1.99 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 0.0% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 15.4% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 15.2% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 14.1% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 9.7% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 95 95 54.4% -3.45 [-6.78, -0.12]
Heterogeneity: Tau² = 4.43; Chi² = 4.92, df = 3 (P = 0.18); I² = 39%
Test for overall effect: Z = 2.03 (P = 0.04)
Total (95% CI) 191 191 100.0% -5.73 [-9.94, -1.53]
Heterogeneity: Tau² = 25.53; Chi² = 26.83, df = 9 (P = 0.001); I² = 66%
Test for overall effect: Z = 2.67 (P = 0.008)
Test for subgroup differences: Chi² = 1.01, df = 1 (P = 0.32), I² = 0.6%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]


Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.6% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 6.0% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.6% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 13.5% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 10.1% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 8.2% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 96 96 46.1% -7.69 [-15.26, -0.11]
Heterogeneity: Tau² = 50.23; Chi² = 13.31, df = 5 (P = 0.02); I² = 62%
Test for overall effect: Z = 1.99 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 13.6% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 0.0% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 15.9% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 14.6% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 9.9% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 95 95 53.9% -3.91 [-6.94, -0.88]
Heterogeneity: Tau² = 1.32; Chi² = 3.46, df = 3 (P = 0.33); I² = 13%
Test for overall effect: Z = 2.53 (P = 0.01)
Total (95% CI) 191 191 100.0% -5.81 [-9.95, -1.67]
Heterogeneity: Tau² = 23.39; Chi² = 23.18, df = 9 (P = 0.006); I² = 61%
Test for overall effect: Z = 2.75 (P = 0.006)
Test for subgroup differences: Chi² = 0.82, df = 1 (P = 0.36), I² = 0%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]


Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 2.0% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 6.8% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 7.3% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 13.0% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 10.4% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 8.7% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 96 96 48.2% -7.69 [-15.26, -0.11]
Heterogeneity: Tau² = 50.23; Chi² = 13.31, df = 5 (P = 0.02); I² = 62%
Test for overall effect: Z = 1.99 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 13.1% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.8% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 0.0% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 13.7% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 10.2% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 90 90 51.8% -1.98 [-5.70, 1.74]
Heterogeneity: Tau² = 5.78; Chi² = 5.06, df = 3 (P = 0.17); I² = 41%
Test for overall effect: Z = 1.04 (P = 0.30)
Total (95% CI) 186 186 100.0% -5.00 [-9.70, -0.29]
Heterogeneity: Tau² = 34.62; Chi² = 30.34, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.08 (P = 0.04)
Test for subgroup differences: Chi² = 1.76, df = 1 (P = 0.19), I² = 43.0%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]


Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.9% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 6.5% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 7.1% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 13.0% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 10.2% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 8.5% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 96 96 47.2% -7.69 [-15.26, -0.11]
Heterogeneity: Tau² = 50.23; Chi² = 13.31, df = 5 (P = 0.02); I² = 62%
Test for overall effect: Z = 1.99 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 13.1% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.9% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 14.8% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 0.0% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 10.0% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 102 102 52.8% -1.82 [-5.11, 1.47]
Heterogeneity: Tau² = 3.77; Chi² = 4.53, df = 3 (P = 0.21); I² = 34%
Test for overall effect: Z = 1.08 (P = 0.28)
Total (95% CI) 198 198 100.0% -4.82 [-9.34, -0.30]
Heterogeneity: Tau² = 31.31; Chi² = 29.99, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.09 (P = 0.04)
Test for subgroup differences: Chi² = 1.94, df = 1 (P = 0.16), I² = 48.4%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]


Nitrate
Placebo
Mean Difference
Mean Difference
Study or Subgroup
Mean
SD
Total
Mean
SD
Total
Weight
IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 Clinic
Baião et al. 2019 145.2664577 22.63323 5 159.5611285 27.39812 5 1.7% -14.29 [-45.44, 16.85]
Broxterman et al. 2019 [1] 129 16 13 132 20 13 6.1% -3.00 [-16.92, 10.92]
Broxterman et al. 2019 [2] 136 16 14 141 19 14 6.6% -5.00 [-18.01, 8.01]
Kapil et al. 2015 130.0296736 13.05638 32 148.5756677 11.72107 32 12.6% -18.55 [-24.63, -12.47]
Siervo et al. 2020 [1] 141.377 13.1148 16 144.26 12.5902 16 9.8% -2.88 [-11.79, 6.03]
Siervo et al. 2020 [2] 146.0328 13.1148 16 148.7869 18.0984 16 8.0% -2.75 [-13.71, 8.20]
Subtotal (95% CI) 96 96 44.7% -7.69 [-15.26, -0.11]
Heterogeneity: Tau² = 50.23; Chi² = 13.31, df = 5 (P = 0.02); I² = 62%
Test for overall effect: Z = 1.99 (P = 0.05)
1.1.2 Ambulatory
Bondonno et al. 2015 128.4 11.3755 27 127.4 11.1227 27 12.7% 1.00 [-5.00, 7.00]
Gilchrist et al. 2013 135.4155 7.33524355 27 134.9570201 8.02292264 27 14.6% 0.46 [-3.64, 4.56]
Kapil et al. 2015 133 8.3 32 137.8 9.1 32 14.5% -4.80 [-9.07, -0.53]
Kerley et al. 2018 132 9 20 138 8 20 13.5% -6.00 [-11.28, -0.72]
Siervo et al. 2020 135.9374 12.2058 16 141.4497 14.1745 16 0.0% -5.51 [-14.68, 3.65]
Subtotal (95% CI) 106 106 55.3% -2.40 [-5.84, 1.05]
Heterogeneity: Tau² = 6.23; Chi² = 6.10, df = 3 (P = 0.11); I² = 51%
Test for overall effect: Z = 1.36 (P = 0.17)
Total (95% CI) 202 202 100.0% -4.90 [-9.18, -0.63]
Heterogeneity: Tau² = 28.08; Chi² = 30.32, df = 9 (P = 0.0004); I² = 70%
Test for overall effect: Z = 2.25 (P = 0.02)
Test for subgroup differences: Chi² = 1.55, df = 1 (P = 0.21), I² = 35.6%
-20 -10 0 10 20
Favours [Nitrate] Favours [Placebo]