Supplementary appendix G. Quality of life among frail TAVI recipients

Study	QoL measures	Baseline (SD)	Post-TAVI (SD)	Change in QoL post-TAVI	30-day (SD)	Change in QoL at 30 days	12-months (SD)
Okoh, 2017	KCCQ overall score					-5.86 (p=.62)	
	KCCQ physical limitation					-8.34 (p=0.46)	
	KCCQ symptoms					-5.68 (p=0.65)	
	KCCQ quality of life					6.36 (p=0.63)	
	KCCQ social limitation					-7.57 (p=0.58)	
Kobe, 2016 Subgroup of frail patients with a frailty score 4-6	SF-36 physical functioning	55.8 (15.3)			49.6 (22.2)		
	SF-36 role physical	37.9 (32.8)			33.8 (43.1)		
	SF-36 bodily pain	88.1 (23.2)			74.4 (23.7)		
	SF-36 role emotional	88.2 (23.6)			38.3 (42.3)		
	SF-36 general health	73.4 (18.7)			62 (15.5)		
	SF-36 vitality	45.5 (24.6)			43.4 (15.5)		
	SF-36 social functioning	91.5 (17.8)			68.8 (26.4)		
	SF-36 mental health	80.4 (16.8)			67.2 (23.2)		
	SF-36 physical component summary	42.9 (6.54)			42.3 (8.02)		
	SF-36 mental component summary	55 (8.79)			42.8 (9.86)		
Kobe, 2016 Subgroup of frail patients with a frailty score 6-14	SF-36 physical functioning	40.9 (20.5)			46.5 (30.2)		
	SF-36 role physical	25.6 (33.2)			27.5 (39.7)		
	SF-36 bodily pain	69.1 (35.6)			60.4 (34.8)		
	SF-36 role emotional	82.5 (32.9)			45 (46.2)		
	SF-36 general health	63.1 (22.7)			50.6 (20.9)		
	SF-36 vitality	35.4 (25.8)			44.1 (24.9)		
	SF-36 social functioning	79.7 (28.7)			67.9 (23.9)		
	SF-36 mental health	75 (16.3)			66 (23.3)		
	SF-36 physical component summary	36.1 (9.57)			38.6 (8.68)		
	SF-36 mental component summary	53 (9.23)			45.2 (11.6)		
Goudzwaard, 2020	EQ-5D index	0.55 (0.26)					0.44 (0.33)
	EQ-VAS	58.4 (16.8)					63.4 (14.6)
Patel, 2020	KCCQ 12 score	36 (25)	75 (70-80)	37 (31-43)			
Morris, 2020	KCCQ 12 score	37	66				

KCCQ, Kansas City Cardiomyopathy Questionnaire. SF-36, Short Form-36 questionnaire. EQ-5D, Euro-Qol-5-dimensions. VAS, visual analogue scale.