Supplementary Table 5: Demographics, salient clinical manifestations, investigations, management, and clinical outcomes in a comparative analysis

Parameters	**Pediatrics (≤18 years)**	**Adults (>18 years)**	**OR (95%CI)**	**P value**
No. of cases (%)	10 (14.1)	61 (85.9)	NA	NA
Age (Median with IQR) in years	11 (5-17)	58 (47-74)	NA	NA
Male (%)	3 (30)	41 (67.2)	0.209 (0.049-0.895)	0.035*
Clinical manifestations				
Severe COVID illness[#]	0	16/52	NA	NA
ICU requirement	1 (10)	23/58	0.169(0.020-1.426)	0.102
Altered mental status	5 (50)	48 (78.7)	0.245 (0.061-0.984)	0.047*
Movement disorder	6 (60)	22 (36.1)	2.659 (0.676-10.452)	0.161
Myoclonus	1 (10)	13 (21.3)	NA	NA
Ataxia	3 (30)	6 (9.8)	11.276 (0.560-227.189)	0.114
Seizures	6 (60)	18 (29.5)	11.078(.481-255.002)	0.133
GTCS	1 (10)	9 (14.8)	0.633(0.023-17.438)	0.787
Status epilepticus	0	10 (16.4)	NA	NA
Focal weakness	0	7 (11.5)	NA	NA
Behavioural and psychiatric symptoms	5 (50)	20 (32.8)	3.971 (0.270-58.456)	0.315
Visual Hallucinations	1 (10)	5 (8.2)	NA	NA
Eye movement abnormalities	0	8 (13.1)	NA	NA
Cognitive symptoms	2 (20)	8 (13.1)	0.688 (0.0.16-29.918)	0.846
Insomnia	2 (20)	6 (9.8)	NA	NA
Speech abnormality	3 (30)	14 (23.0)	NA	NA
Mutism	2 (20)	6 (9.8)	NA	NA
Investigations				
Elevation of at least one inflammation marker	3/5 (60)	20/24 (83.3)	0.300(0.037-2.417)	0.258
MRI abnormality	2/10 (20)	35/54 (64.8)	0.136(0.026-0.705)	0.017*

EEG abnormality	6/6 (100)	30/37 (81.1)	--	--
Timelines				
Latency to neurological symptoms: Median (IQR) days	0 (0-5)	2 (0-13)	NA	0.477
Latency to relapse or non-remission of neurological symptoms: Median (IQR) days	14 (4-17)	12 (3-22)	NA	0.728
Duration at presentation: Median (IQR) days	6 (4-14)	7 (3-14)	NA	0.859
Time to initiation of immunotherapy from diagnosis: Median (IQR) days	7 (3-21)	18 (9-41)	NA	0.160
Diagnosis at second admission (%)	2 (20)	4 (6.6)	4.800 (0.938-24.568)	0.06
Treatment				
At least one form of immunotherapy	9 (90)	38/47 (80.9)	NA	NA
Parenteral steroids[&]	8 (80)	31/47 (66.0)	1.290(0.132-12.658)	0.827
IVIg	6 (60)	17/47 (36.2)	3.00(0.529-17.020)	0.215
IVIg 2nd course	1 (10)	2/47 (4.3)	--	--
Plasma exchange	0	8/47 (17)	--	--
Combination[$]	5 (50)	16/47 (34.0)	1.458(0.294-7.231)	0.644
No immunotherapy	1 (10)	8/47 (17)	--	--
Clinical outcomes				
Immunotherapy responders	8/9 (88.9)	34/39 (87.1)	1.176(0.120-11.531)	0.889
Latency to therapeutic response in median (IQR) in days	7 (3-14)	10 (5-22)	NA	0.364
Sequelae[#]	3/10 (30)	19/50 (38)	0.699(0.161-3.035)	0.633
Expired	0	4 (6.6)	--	--

© One of the studies doesn't have information of sex (J. Valadez-Calderon et al 2021(35))

*One patient had both Anti-GAD65 and Anti-NMDAR positivity;

#Includes 4 deaths

$ Combination therapy included use of steroids or IVIg or PLEX in combination. One of the patients was given IVIg with Tocilizulab (Fiere-alvarez. 2020 (57))

&Two patients were given dexamethasone instead of Methylprednisolone (Panariello et al,2020 (17) , Urso et al, 2020 (40))

GTCS: Generalized tonic clonic seizures, "--": Not available/applicable