Supplementary File 3. Characteristics and Reference List of Included Studies

Supplementary Table 1. Detailed Characteristics of Studies Included in the Systematic Review

First Author	Publication Year	Country	Study Design	Patient Numbers	Age*	Male/ Female	SBE Procedures	Oral/Anal Route	Numbers of Diagnosis	Numbers of Findings	Total Enterscopy/ Total Attempts	Adverse Events
Vargo[1]	2007	USA	Case series	20	65.9±19.5	9/11	20	19/1	11	11	NA	0
Kawamura[2]	2008	Japan	Prospective case series	27	56.6（24-83）	20/7	37	22/15	NA	11	1/8	1
Ohtsuka[3]	2008	Japan	Case series	30	19-78***	21/9	48	14/34	NA	NA	5/7	0
Tsujikawa[4]	2008	Japan	Case series	41	48.9（23-85）	24/17	78	38/14	22	22	6/24	2
Ramchandani[5]	2009	India	Prospective case series	106	40.1±14.6	65/41	131	95/36	68	68	5/20	8
Vere[6]	2009	Romnia	Case series	21	50.28±17.48	11/10	21	NA	16	16	NA	NA
Aktas[7]	2010	Netherlands	Prospective cohort	105	51**	53/52	166	105/61	NA	65	4/99	14
Barth[8]	2010	USA	Retrospective case series	6	13**	3/3	8	5/3	3	3	0/2	1
Frantz[9]	2010	USA	Retrospective case series	38	62±17	22/16	38	38/0	18	18	NA	0
Khashab[10]	2010	USA	Retrospective cohort	41	55±16	15/26	52	52/0	NA	NA	NA	2
May[11]	2010	Germany	RCT	50	56±18	33/17	NA	NA	21	21	11/50	2
Trifan[12]	2010	Romania	Case series	3	52±11	2/1	3	3/0	3	3	NA	0
Upchurch[13]	2010	USA	Retrospective case series	161	64（23-88）	78/83	172	143/29	NA	NA	NA	1
Dulic-Lakovic[14]	2011	Austria	Retrospective case series	NA	NA	NA	78	70/8	NA	NA	NA	0
Riccioni[15]	2011	Italy	Prospective case series	70	51.6（26-84）	36/34	73	63/10	40	43	NA	NA
Takano[16]	2011	Japan	RCT	18	64.9±14.7	13/5	32	15/17	11	11	0/14	1
Alderlieste[17]	2012	Netherlands	Prospective case series	10	55(31-64)	4/6	10	10/0	9	9	NA	1
Chalazan[18]	2012	USA	Diagnostic test	14	NA	NA	14	14/0	NA	14	NA	NA
Nardo[19]	2012	Italy	Prospective cohort	30	13.9**	17/13	36	23/13	28	28	NA	0
Dutta[20]	2012	India	Retrospective case series	84	42±15	61/23	90	48/42	27	27	0/6	1
Efthymiou[21]	2012	Australia	RCT	50	67（51,72）	19/31	53	32/21	20	NA	NA	1
Gong[22]	2012	China	Retrospective case series	67	39.7（12-73）	47/20	80	42/38	46	46	4/13	0
Kushnir[23]	2013	USA	Retrospective cohort	147	69.6±12.7	69/78	150	146/4	90	90	NA	1
Lenz[24]	2013	Germany	RCT	107	56.6**	53/54	169	98/71	75	NA	NA	0
Lenz[25]	2013	Germany	Retrospective cohort	298	55±19.1	152/146	515	277/238	184	NA	NA	2
Luo[26]	2013	China	Retrospective case series	70	NA	38/32	82	38/44	NA	58	NA	0
Manno[27]	2013	Italy	Case series	111	61.0±17.9	62/49	131	99/32	82	82	8/17	1

Supplementary File 3. Characteristics and Reference List of Included Studies

Continued

First Author	Publication Year	Country	Study Design	Patient Numbers	Age*	Male/ Female	SBE Procedures	Oral/Anal Route	Numbers of Diagnosis	Numbers of Findings	Total Enterscopy/ Total Attempts	Adverse Events
Prachayakul[28]	2013	Thailand	Retrospective cohort	116	58.1±17.7	58/58	145	NA	NA	NA	1/16	15
Saigusa[29]	2013	Japan	Retrospective case series	11	NA	10/1	11	0/11	NA	8	NA	NA
Baijal[30]	2014	India	Retrospective case series	40	47.9±13.5	30/10	48	33/15	22	22	NA	2
Li[31]	2014	China	RCT	214	40.85**	125/89	365	186/179	113	113	56/151	2
Sethi[32]	2014	USA	Retrospective cohort	150	63**	90/60	170	152/18	NA	NA	1/6	1
Banerjee[33]	2015	USA	Retrospective case series	NA	NA	NA	46	37/9	NA	NA	NA	0
Davis-Yadley[34]	2015	USA	Retrospective cohort	428	63±15.5	225/203	439	390/49	247	270	NA	10
Holman[35]	2015	USA	Retrospective cohort	125	63.1**	66/59	125	116/9	72	85	NA	NA
Kav[36]	2015	Turkey	Prospective diagnostic test	33	35.7±12.9	8/25	33	33/0	22	22	NA	NA
Minamino[37]	2015	Japan	Prospective case series	16	45.6（25-61）	11/5	16	0/16	6	6	NA	NA
Oliva[38]	2015	Italy	Prospective case series	22	12.5±3.9	14/8	27	16/11	19	19	2/7	0
Pinho[39]	2015	Portugal	Retrospective case series	NA	NA	NA	351	239/112	NA	NA	NA	2
Tao[40]	2015	China	Retrospective case series	186	45.87±15.77	102/84	196	118/78	NA	129	NA	NA
Tsujikawa[41]	2015	Japan	Case control study	84	52.9(17-86)	57/27	84	84/0	NA	NA	NA	3
Christian[42]	2016	USA	Retrospective case series	136	57.5±16.2	67/69	136	0/136	73	69	NA	2
Lin[43]	2016	China	Retrospective case series	128	58.22±20.69	101/99	200	101/99	NA	NA	NA	19
Ma[44]	2016	China	Retrospective case series	353	42.1±15.8	235/118	419	98/321	246	246	NA	0
Ooka[45]	2016	Japan	Retrospective case series	91	68±17	45/46	91	43/48	67	67	NA	3
Takenaka[46]	2016	Japan	Prospective diagnostic test	200	NA	137/63	200	0/200	NA	94	NA	NA
Haanstra[47]	2017	Netherlands	Prospective cohort	3	46**	0/3	3	0/3	2	2	NA	0
Kim[48]	2017	Korea	Retrospective cohort	44	50.1±22	38/27	65	39/26	38	38	2/16	2
Lu[49]	2017	China	Retrospective cohort	79	52.24**	42/37	79	29/50	48	48	NA	9
Marques[50]	2017	Portugal	Retrospective case series	168	53.3±17.6	108/60	197	122/75	NA	NA	4/6	3
Xu[51]	2017	China	Case control study	132	63.92**	66/66	188	77/111	83	99	22/46	NA
Holleran[52]	2018	Italy	Retrospective case series	52	44.3**	41/11	56	11/45	35	35	NA	NA
Lopez[53]	2018	Spain	Prospective case series	39	74（18-89）	24/15	42	NA	NA	24	NA	0
Moran[54]	2018	USA	RCT	17	55.8±20.3	3/14	17	17/0	7	7	NA	8

Supplementary File 3. Characteristics and Reference List of Included Studies

[*]Values of age were presented as median (interquartile range) or mean±s.d. unless indicated otherwise.

[**] Values of age were presented as mean.

[***] Value of age was presented as range.

RCT, Randomized Controlled Trial; NA: Not Available.

Supplementary File 3. Characteristics and Reference List of Included Studies

List of studies included in the systematic review

1. Vargo J, Upchurch B, Dumot J, et al. Clinical utility of the olympus single balloon enteroscope: the initial U.S. experience. Gastrointest Endosc. 2007;65:AB90

2. Kawamura T, Yasuda K, Tanaka K, et al. Clinical evaluation of a newly developed single-balloon enteroscope. Gastrointest Endosc. 2008;68:1112-6.

3. Ohtsuka K, Kashida H, Kodama K, et al. Diagnosis and treatment of small bowel diseases with a newly developed single balloon endoscope. Dig Endosc 2008;20:134-7.

4. Tsujikawa T, Saitoh Y, Andoh A, et al. Novel single-balloon enteroscopy for diagnosis and treatment of the small intestine: preliminary experiences. Endoscopy. 2008;40:11-5.

5. Ramchandani M, Reddy DN, Gupta R, et al. Diagnostic yield and therapeutic impact of single-balloon enteroscopy: series of 106 cases. J Gastroenterol Hepatol. 2009;24:1631-8.

6. Vere CC, Foarfa C, Streba CT, et al. Videocapsule endoscopy and single balloon enteroscopy: novel diagnostic techniques in small bowel pathology. Rom J Morphol Embryol. 2009;50:467-74.

7. Aktas H, de Ridder L, Haringsma J, et al. Complications of single-balloon enteroscopy: a prospective evaluation of 166 procedures. Endoscopy. 2010;42:365-8.

8. Barth BA, Channabasappa N. Single-balloon enteroscopy in children: initial experience at a pediatric center. J Pediatr Gastroenterol Nutr. 2010;51:680-4.

9. Frantz DJ, Dellon ES, Grimm IS, et al. Single-balloon enteroscopy: results from an initial experience at a U.S. tertiary-care center. Gastrointest Endosc. 2010;72:422-6.

10. Khashab MA, Lennon AM, Dunbar KB, et al. A comparative evaluation of single-balloon enteroscopy and spiral enteroscopy for patients with mid-gut disorders. Gastrointest Endosc. 2010;72:766-72.

11. May A, Farber M, Aschmoneit I, et al. Prospective multicenter trial comparing push-and-pull enteroscopy with the single- and double-balloon techniques in patients with small-bowel disorders. Am J Gastroenterol. 2010;105:575-81.

12. Trifan A, Singeap AM, Cojocariu C, et al. Single-balloon enteroscopy following videocapsule endoscopy for diagnosis of small bowel tumors: preliminary experiences. Chirurgia (Bucur). 2010;105:211-7.

13. Upchurch BR, Sanaka MR, Lopez AR, et al. The clinical utility of single-balloon enteroscopy: a single-center experience of 172 procedures. Gastrointest Endosc. 2010;71:1218-23.

14. Dulic-Lakovic E, Dulic M, Hubner D, et al. Bleeding Dieulafoy lesions of the small bowel: a systematic study on the epidemiology and efficacy of enteroscopic treatment. Gastrointest Endosc. 2011;74:573-80.

15. Riccioni ME, Urgesi R, Cianci R, et al. Single-balloon push-and-pull enteroscopy system: does it work?

A single-center, 3-year experience. Surg Endosc. 2011;25:3050-6.

16. Takano N, Yamada A, Watabe H, et al. Single-balloon versus double-balloon endoscopy for achieving total enteroscopy: a randomized, controlled trial. Gastrointest Endosc. 2011;73:734-9.

17. Alderlieste YA, Rauws EA, Mathus-Vliegen EM, et al. Prospective enteroscopic evaluation of jejunal polyposis in patients with familial adenomatous polyposis and advanced duodenal polyposis. Fam Cancer. 2012;12:51-6.

18. Chalazan B, Gostout CJ, Song LM, et al. Use of Capsule small bowel transit time to determine the optimal enteroscopy approach. Gastroenterology Res. 2012;5:39-44.

19. Di Nardo G, Oliva S, Aloi M, et al. Usefulness of single-balloon enteroscopy in pediatric Crohn's disease. Gastrointest Endosc. 2012;75:80-6.

20. Dutta AK, Sajith KG, Joseph AJ, et al. Learning curve, diagnostic yield and safety of single balloon enteroscopy. Trop Gastroenterol. 2012;33:179-84.

21. Efthymiou M, Desmond PV, Brown G, et al. SINGLE-01: a randomized, controlled trial comparing the efficacy and depth of insertion of single- and double-balloon enteroscopy by using a novel method to determine insertion depth. Gastrointest Endosc. 2012;76:972-80.

22. Gong W, Zhi FC, Bai Y, et al. Single-balloon enteroscopy for small bowel diseases: results from an initial experience at a single Chinese center. J Dig Dis. 2012;13:609-13.

23. Kushnir VM, Tang M, Goodwin J, et al. Long-term outcomes after single-balloon enteroscopy in patients with obscure gastrointestinal bleeding. Dig Dis Sci. 2013;58:2572-9.

24. Lenz P, Meister T, Manno M, et al. CO2 insufflation during single-balloon enteroscopy: a multicenter randomized controlled trial. Endoscopy. 2013;46:53-8.

25. Lenz P, Roggel M, Domagk D. Double- vs. single-balloon enteroscopy: single center experience with emphasis on procedural performance. Int J Colorectal Dis. 2013;28:1239-46.

26. Luo YH, You JY, Liu L, et al. Clinical application of single-balloon enteroscopy in children with small intestinal bleeding. Zhongguo Dang Dai Er Ke Za Zhi. 2013;15:546-9.

27. Manno M, Riccioni ME, Cannizzaro R, et al. Diagnostic and therapeutic yield of single balloon enteroscopy in patients with suspected small-bowel disease: results of the Italian multicentre study. Dig Liver Dis. 2013;45:211-5.

28. Prachayakul V, Deesomsak M, Aswakul P, et al. The utility of single-balloon enteroscopy for the diagnosis and management of small bowel disorders according to their clinical manifestations: a retrospective review. BMC Gastroenterol. 2013;13:103.

29. Saigusa N, Yokoyama T, Shinozaki M, et al. Anorectal fistula is an early manifestation of Crohn's disease that occurs before bowel lesions advance: a study of 11 cases. Clin J Gastroenterol. 2013;6:309-14.

30. Baijal R, Kumar P, Gupta DT, et al. Single-balloon enteroscopy: a single-center experience of 48 procedures. Indian J Gastroenterol. 2014;33(1):55-8.

31. Li X, Zhao YJ, Dai J, et al. Carbon dioxide insufflation improves the intubation depth and total enteroscopy rate in single-balloon enteroscopy: a randomised, controlled, double-blind trial. Gut. 2014;63:1560-5.

32. Sethi S, Cohen J, Thaker AM, et al. Prior capsule endoscopy improves the diagnostic and therapeutic yield of single-balloon enteroscopy. Dig Dis Sci. 2014;59:2497-502.

33. Banerjee N, Presta M, Charous M, et al. Revenue from single-balloon enteroscopy is driven by anesthesia: experience from a tertiary care facility. Surg Endosc. 2015;30:1635-9.

34. Davis-Yadley AH, Lipka S, Rodriguez AC, et al. The safety and efficacy of single balloon enteroscopy in the elderly. Therap Adv Gastroenterol. 2015;9:169-79.

35. Holman N, Wallace K, Moore JM, et al. Open-access single balloon enteroscopy: a tertiary care experience. South Med J. 2015;108:739-43.

36. Kav T, Sokmensuer C, Sivri B. Enteroscopic findings of Celiac Disease and their correlation with mucosal histopathologic changes. Comput Biol Med. 2015;65:315-9.

37. Minamino H, Machida H, Tominaga K, et al. Rectal biopsy, rather than ileal, is appropriate to confirm the diagnosis of early gastrointestinal graft-versus-host disease. Scand J Gastroenterol. 2015;50:1428-34.

38. Oliva S, Pennazio M, Cohen SA, et al. Capsule endoscopy followed by single balloon enteroscopy in children with obscure gastrointestinal bleeding: A combined approach. Digest Liver Dis. 2015;47:125-30.

39. Pinho R, Mascarenhas-Saraiva M, Mao-de-Ferro S, et al. Multicenter survey on the use of device-assisted enteroscopy in Portugal. United European Gastroenterol J. 2015;4:264-74.

40. Tao Z, Liu GX, Cai L, et al. Characteristics of small intestinal diseases on single-balloon enteroscopy: a single-center study conducted over 6 years in China. Medicine (Baltimore). 2015;94:e1652.

41. Tsujikawa T, Bamba S, Inatomi O, et al. Factors affecting pancreatic hyperamylasemia in patients undergoing peroral single-balloon enteroscopy. Dig Endosc. 2015;27:674-8.

42. Christian KE, Kapoor K, Goldberg EM. Performance characteristics of retrograde single-balloon endoscopy: A single center experience. World J Gastrointest Endosc. 2016;8:501-7.

43. Lin M, Chen P, Shih Y, et al. Outcome and safety of anterograde and retrograde single-balloon enteroscopy: clinical experience at a tertiary medical center in Taiwan. PLOS One. 2016;11:e161188.

44. Ma J, Wang Y, Xu X, et al. Capsule endoscopy and single-balloon enteroscopy in small bowel diseases: Competing or complementary? World J Gastroenterol. 2016;22:10625.

45. Ooka S, Kobayashi K, Kawagishi K, et al. Roles of capsule endoscopy and single-balloon enteroscopy in diagnosing unexplained gastrointestinal bleeding. Clin Endosc. 2016;49:56-60.

46. Takenaka K, Ohtsuka K, Kitazume Y, et al. Magnetic resonance evaluation for small bowel strictures in Crohn's disease: comparison with balloon enteroscopy. J Gastroenterol. 2016;52:879-88.

47. Haanstra JF, Al-Toma A, Dekker E, et al. Incidence of small bowel neoplasia in Lynch syndrome assessed by video capsule endoscopy. Endosc Int Open. 2017;5:E622-6.

48. Kim TJ, Kim ER, Chang DK, et al. Comparison of the efficacy and safety of single- versus double-balloon enteroscopy performed by endoscopist experts in single-balloon enteroscopy: a single-center experience and meta-analysis. Gut Liver. 2017;11:520-7.

49. Lu Z, Qi Y, Weng J, et al. Efficacy and safety of single-balloon versus double-balloon enteroscopy: a single-center retrospective analysis. Med Sci Monit. 2017;23:1933-9.

50. Marques M, Santos-Antunes J, Coelho R, et al. Single-balloon enteroscopy efficacy and degree of concordance with noninvasive evaluation of small bowel. Endosc Int Open. 2017;5:E96-102.

51. Xu N, Yu Z, Cao X, et al. Characteristics of nonsteroidal anti-inflammatory drugs (NSAIDs)-induced small bowel injury identified by single-balloon endoscopy or capsule endoscopy. Med Sci Monit. 2017;23:5237-45.

52. Holleran G, Valerii G, Tortora A, et al. The use of single balloon enteroscopy in Crohn's disease and its impact on clinical outcome. Scand J Gastroenterol. 2018:1-5.

53. Lopez RL, Alvarez B, Gonzalez RA, et al. Viability of single balloon enteroscopy performed under endoscopist-directed sedation. Rev Esp Enferm Dig. 2018;110:240-5.

54. Moran RA, Barola S, Law JK, et al. A randomized controlled trial comparing the depth of maximal insertion between anterograde single-balloon versus spiral enteroscopy. Clin Med Insights Gastroenterol. 2018;11:419549375.